



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:51 AM GMT

PDB ID : 3JSE
Title : Crystal structure of archaeal 20S proteasome in complex with mutated P26 activator
Authors : Stadtmueller, B.M.; Whitby, F.G.; Hill, C.P.
Deposited on : 2009-09-10
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>6%</div> <div>82%</div> <div>17%</div> </div>
1	B	227	<div> <div>5%</div> <div>84%</div> <div>15%</div> </div>
1	C	227	<div> <div>5%</div> <div>85%</div> <div>14%</div> </div>
1	D	227	<div> <div>6%</div> <div>86%</div> <div>14%</div> </div>
1	E	227	<div> <div>7%</div> <div>81%</div> <div>19%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	227	 6% 81% 19%
1	G	227	 5% 81% 18%
2	H	203	 % 83% 15% •
2	I	203	 % 84% 14% •
2	J	203	 2% 81% 17% •
2	K	203	 2% 85% 13% •
2	L	203	 % 82% 15% ••
2	M	203	 2% 82% 16% •
2	N	203	 3% 86% 12% •
3	O	228	 7% 78% 18% •
3	P	228	 9% 82% 14% •
3	Q	228	 8% 82% 13% •
3	R	228	 8% 81% 14% •
3	S	228	 7% 78% 18% •
3	T	228	 7% 81% 14% •
3	U	228	 9% 81% 14% ••

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 35056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	B	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	C	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	D	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	E	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	F	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			
1	G	227	Total	C	N	O	S	0	0	0
			1768	1123	299	343	3			

- Molecule 2 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	I	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	J	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	K	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	L	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	M	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			
2	N	203	Total	C	N	O	S	0	0	0
			1557	985	264	297	11			

- Molecule 3 is a protein called Proteasome activator protein PA26.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	O	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	P	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	Q	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	R	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	S	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	T	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0
3	U	218	Total 1683	C 1056	N 296	O 325	S 6	0	0	0


There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	49	VAL	THR	VARIANT	UNP Q9U8G2
O	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
P	49	VAL	THR	VARIANT	UNP Q9U8G2
P	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
Q	49	VAL	THR	VARIANT	UNP Q9U8G2
Q	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
R	49	VAL	THR	VARIANT	UNP Q9U8G2
R	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
S	49	VAL	THR	VARIANT	UNP Q9U8G2
S	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
T	49	VAL	THR	VARIANT	UNP Q9U8G2
T	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2
U	49	VAL	THR	VARIANT	UNP Q9U8G2
U	230	PHE	VAL	ENGINEERED	UNP Q9U8G2
O	226	THR	SER	ENGINEERED	UNP Q9U8G2


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha





• Molecule 1: Proteasome subunit alpha








- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta





- Molecule 3: Proteasome activator protein PA26









- Molecule 3: Proteasome activator protein PA26





- Molecule 3: Proteasome activator protein PA26




- Molecule 3: Proteasome activator protein PA26




- Molecule 3: Proteasome activator protein PA26





- Molecule 3: Proteasome activator protein PA26





● Molecule 3: Proteasome activator protein PA26



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	254.70Å 126.10Å 180.39Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	19.86 – 2.90 19.86 – 2.90	Depositor EDS
% Data completeness (in resolution range)	93.0 (19.86-2.90) 93.0 (19.86-2.90)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.88Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.212 , 0.241 0.218 , 0.246	Depositor DCC
R_{free} test set	2328 reflections (2.03%)	DCC
Wilson B-factor (Å ²)	61.7	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 28.9	EDS
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 117219 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	35056	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/1792	0.73	0/2416
1	B	0.80	0/1792	0.75	0/2416
1	C	0.81	1/1792 (0.1%)	0.74	0/2416
1	D	0.83	0/1792	0.74	0/2416
1	E	0.83	1/1792 (0.1%)	0.77	1/2416 (0.0%)
1	F	0.80	0/1792	0.77	1/2416 (0.0%)
1	G	0.82	0/1792	0.75	1/2416 (0.0%)
2	H	0.79	0/1576	0.79	1/2129 (0.0%)
2	I	0.81	0/1576	0.81	2/2129 (0.1%)
2	J	0.79	0/1576	0.79	0/2129
2	K	0.77	0/1576	0.79	1/2129 (0.0%)
2	L	0.82	0/1576	0.80	2/2129 (0.1%)
2	M	0.80	0/1576	0.82	2/2129 (0.1%)
2	N	0.79	0/1576	0.81	2/2129 (0.1%)
3	O	0.75	0/1707	0.71	0/2307
3	P	0.76	0/1707	0.70	0/2307
3	Q	0.76	0/1707	0.72	0/2307
3	R	0.77	0/1707	0.71	0/2307
3	S	0.75	1/1707 (0.1%)	0.70	0/2307
3	T	0.73	0/1707	0.70	0/2307
3	U	0.79	1/1707 (0.1%)	0.70	0/2307
All	All	0.79	4/35525 (0.0%)	0.75	13/47964 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	U	35	GLU	CG-CD	5.69	1.60	1.51
3	S	35	GLU	CG-CD	5.38	1.60	1.51
1	E	151	CYS	CB-SG	-5.28	1.73	1.81
1	C	132	TYR	CE2-CZ	-5.17	1.31	1.38

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	203	LEU	CA-CB-CG	5.86	128.77	115.30
2	L	157	ARG	NE-CZ-NH2	-5.57	117.52	120.30
2	M	203	LEU	CA-CB-CG	5.56	128.08	115.30
1	E	147	ARG	NE-CZ-NH1	5.54	123.07	120.30
2	M	157	ARG	NE-CZ-NH2	-5.49	117.56	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1768	0	1800	23	0
1	B	1768	0	1800	25	0
1	C	1768	0	1800	20	0
1	D	1768	0	1800	13	0
1	E	1768	0	1800	27	0
1	F	1768	0	1800	30	0
1	G	1768	0	1800	25	0
2	H	1557	0	1609	20	0
2	I	1557	0	1609	16	0
2	J	1557	0	1609	21	0
2	K	1557	0	1609	18	0
2	L	1557	0	1609	22	0
2	M	1557	0	1609	21	0
2	N	1557	0	1609	17	0
3	O	1683	0	1702	31	0
3	P	1683	0	1702	24	0
3	Q	1683	0	1702	22	0
3	R	1683	0	1702	24	1
3	S	1683	0	1702	29	1
3	T	1683	0	1702	27	1
3	U	1683	0	1702	24	1
All	All	35056	0	35777	395	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 395 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:8:TYR:OH	1:F:9:ASP:OD2	1.96	0.84
3:T:89:ARG:HD3	3:U:203:THR:HG21	1.64	0.79
3:T:142:MET:HE2	3:U:192:LEU:HD21	1.68	0.75
1:F:8:TYR:OH	1:G:9:ASP:OD2	2.01	0.73
3:O:68:LEU:HD21	3:O:149:LEU:HD21	1.70	0.72

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:R:14:ASP:OD1	3:U:45:GLU:OE1[4_555]	1.45	0.75
3:S:45:GLU:OE1	3:T:14:ASP:OD1[4_555]	1.80	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/227 (99%)	209 (93%)	14 (6%)	2 (1%)	21	57
1	B	225/227 (99%)	209 (93%)	14 (6%)	2 (1%)	21	57
1	C	225/227 (99%)	211 (94%)	12 (5%)	2 (1%)	21	57
1	D	225/227 (99%)	209 (93%)	14 (6%)	2 (1%)	21	57
1	E	225/227 (99%)	210 (93%)	13 (6%)	2 (1%)	21	57
1	F	225/227 (99%)	213 (95%)	10 (4%)	2 (1%)	21	57
1	G	225/227 (99%)	212 (94%)	11 (5%)	2 (1%)	21	57
2	H	201/203 (99%)	193 (96%)	7 (4%)	1 (0%)	34	71
2	I	201/203 (99%)	193 (96%)	7 (4%)	1 (0%)	34	71
2	J	201/203 (99%)	194 (96%)	6 (3%)	1 (0%)	34	71

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	K	201/203 (99%)	194 (96%)	6 (3%)	1 (0%)	34	71
2	L	201/203 (99%)	194 (96%)	6 (3%)	1 (0%)	34	71
2	M	201/203 (99%)	192 (96%)	8 (4%)	1 (0%)	34	71
2	N	201/203 (99%)	192 (96%)	8 (4%)	1 (0%)	34	71
3	O	214/228 (94%)	207 (97%)	7 (3%)	0	100	100
3	P	214/228 (94%)	208 (97%)	6 (3%)	0	100	100
3	Q	214/228 (94%)	208 (97%)	6 (3%)	0	100	100
3	R	214/228 (94%)	207 (97%)	7 (3%)	0	100	100
3	S	214/228 (94%)	208 (97%)	6 (3%)	0	100	100
3	T	214/228 (94%)	209 (98%)	5 (2%)	0	100	100
3	U	214/228 (94%)	208 (97%)	6 (3%)	0	100	100
All	All	4480/4606 (97%)	4280 (96%)	179 (4%)	21 (0%)	34	71

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	ILE
1	B	64	ILE
1	D	64	ILE
1	E	64	ILE
1	F	64	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/188 (100%)	176 (94%)	12 (6%)	22	53
1	B	188/188 (100%)	176 (94%)	12 (6%)	22	53
1	C	188/188 (100%)	176 (94%)	12 (6%)	22	53
1	D	188/188 (100%)	175 (93%)	13 (7%)	19	48
1	E	188/188 (100%)	175 (93%)	13 (7%)	19	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	188/188 (100%)	175 (93%)	13 (7%)	19	48
1	G	188/188 (100%)	174 (93%)	14 (7%)	17	44
2	H	170/170 (100%)	159 (94%)	11 (6%)	21	52
2	I	170/170 (100%)	160 (94%)	10 (6%)	24	58
2	J	170/170 (100%)	159 (94%)	11 (6%)	21	52
2	K	170/170 (100%)	159 (94%)	11 (6%)	21	52
2	L	170/170 (100%)	159 (94%)	11 (6%)	21	52
2	M	170/170 (100%)	159 (94%)	11 (6%)	21	52
2	N	170/170 (100%)	159 (94%)	11 (6%)	21	52
3	O	179/187 (96%)	173 (97%)	6 (3%)	44	79
3	P	179/187 (96%)	173 (97%)	6 (3%)	44	79
3	Q	179/187 (96%)	173 (97%)	6 (3%)	44	79
3	R	179/187 (96%)	173 (97%)	6 (3%)	44	79
3	S	179/187 (96%)	173 (97%)	6 (3%)	44	79
3	T	179/187 (96%)	175 (98%)	4 (2%)	60	88
3	U	179/187 (96%)	173 (97%)	6 (3%)	44	79
All	All	3759/3815 (98%)	3554 (94%)	205 (6%)	27	61

5 of 205 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	23	GLU
2	J	25	PHE
3	S	19	THR
2	H	72	VAL
2	I	25	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	24	ASN
3	O	75	GLN
3	T	228	HIS
2	M	24	ASN
2	M	73	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/227 (100%)	0.03	14 (6%) 24 17	43, 55, 79, 87	0
1	B	227/227 (100%)	0.05	12 (5%) 30 23	43, 55, 79, 87	0
1	C	227/227 (100%)	0.03	12 (5%) 30 23	43, 55, 79, 87	0
1	D	227/227 (100%)	0.07	14 (6%) 24 17	43, 55, 79, 87	0
1	E	227/227 (100%)	0.06	15 (6%) 22 16	43, 55, 79, 87	0
1	F	227/227 (100%)	0.01	13 (5%) 27 21	43, 55, 79, 87	0
1	G	227/227 (100%)	-0.03	11 (4%) 34 28	43, 55, 79, 87	0
2	H	203/203 (100%)	-0.17	3 (1%) 76 74	44, 50, 65, 76	0
2	I	203/203 (100%)	-0.13	2 (0%) 84 82	44, 50, 65, 76	0
2	J	203/203 (100%)	-0.13	4 (1%) 68 64	44, 50, 65, 76	0
2	K	203/203 (100%)	-0.14	4 (1%) 68 64	44, 50, 65, 76	0
2	L	203/203 (100%)	-0.13	3 (1%) 76 74	44, 50, 65, 76	0
2	M	203/203 (100%)	-0.12	4 (1%) 68 64	44, 50, 65, 76	0
2	N	203/203 (100%)	-0.15	7 (3%) 49 41	44, 50, 65, 76	0
3	O	218/228 (95%)	0.25	16 (7%) 18 12	46, 60, 80, 91	0
3	P	218/228 (95%)	0.15	20 (9%) 11 7	46, 60, 80, 91	0
3	Q	218/228 (95%)	0.23	18 (8%) 14 9	46, 60, 80, 91	0
3	R	218/228 (95%)	0.33	19 (8%) 13 8	46, 60, 80, 91	0
3	S	218/228 (95%)	0.16	16 (7%) 18 12	46, 60, 80, 91	0
3	T	218/228 (95%)	0.23	15 (6%) 20 14	46, 60, 80, 91	0
3	U	218/228 (95%)	0.29	20 (9%) 11 7	46, 60, 80, 91	0
All	All	4536/4606 (98%)	0.05	242 (5%) 30 23	43, 55, 78, 91	0

The worst 5 of 242 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	R	132	GLY	7.8
3	R	133	SER	7.4
3	Q	133	SER	6.6
2	I	202	ILE	6.3
3	T	131	SER	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.