



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:50 AM GMT

PDB ID : 3JVZ  
Title : E2 Ubiquitin-HECT  
Authors : Souphron, J.; Kamadurai, H.B.; Schulman, B.A.  
Deposited on : 2009-09-17  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	<div> <div>55%</div> <div>42%</div> <div>.</div> </div>
1	B	146	<div> <div>54%</div> <div>38%</div> <div>7%</div> <div>.</div> </div>
2	C	385	<div> <div>%</div> <div>50%</div> <div>44%</div> <div>.</div> <div>.</div> </div>
2	D	385	<div> <div>51%</div> <div>41%</div> <div>6%</div> <div>.</div> </div>
3	X	81	<div> <div>30%</div> <div>46%</div> <div>44%</div> <div>.</div> <div>6%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	Y	81	 A horizontal bar chart showing the quality of chain Y. The bar is divided into four segments: red (27%), green (44%), yellow (46%), and orange (6%). The segments are labeled with their respective percentages: 27%, 44%, 46%, and 6%.

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-conjugating enzyme E2 D2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1170	749	202	213	6			
1	B	146	Total	C	N	O	S	0	0	0
			1170	749	202	213	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	SER	LEU	ENGINEERED	UNP P62837
A	85	SER	CYS	ENGINEERED	UNP P62837
A	98	LYS	THR	ENGINEERED	UNP P62837
B	3	SER	LEU	ENGINEERED	UNP P62837
B	85	SER	CYS	ENGINEERED	UNP P62837
B	98	LYS	THR	ENGINEERED	UNP P62837

- Molecule 2 is a protein called E3 ubiquitin-protein ligase NEDD4-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	377	Total	C	N	O	S	0	0	0
			3148	2037	520	574	17			
2	D	374	Total	C	N	O	S	0	0	0
			3124	2021	517	569	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	571	GLY	-	EXPRESSION TAG	UNP Q96PU5
C	572	SER	-	EXPRESSION TAG	UNP Q96PU5
C	573	PRO	-	EXPRESSION TAG	UNP Q96PU5
C	574	GLU	-	EXPRESSION TAG	UNP Q96PU5
C	575	PHE	-	EXPRESSION TAG	UNP Q96PU5
C	922	ALA	CYS	ENGINEERED	UNP Q96PU5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	571	GLY	-	EXPRESSION TAG	UNP Q96PU5
D	572	SER	-	EXPRESSION TAG	UNP Q96PU5
D	573	PRO	-	EXPRESSION TAG	UNP Q96PU5
D	574	GLU	-	EXPRESSION TAG	UNP Q96PU5
D	575	PHE	-	EXPRESSION TAG	UNP Q96PU5
D	922	ALA	CYS	ENGINEERED	UNP Q96PU5

- Molecule 3 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	X	76	Total	C	N	O	S	0	0	0
			601	378	105	117	1			
3	Y	76	Total	C	N	O	S	0	0	0
			601	378	105	117	1			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-4	GLY	-	EXPRESSION TAG	UNP P62988
X	-3	SER	-	EXPRESSION TAG	UNP P62988
X	-2	GLY	-	EXPRESSION TAG	UNP P62988
X	-1	GLY	-	EXPRESSION TAG	UNP P62988
X	0	SER	-	EXPRESSION TAG	UNP P62988
Y	-4	GLY	-	EXPRESSION TAG	UNP P62988
Y	-3	SER	-	EXPRESSION TAG	UNP P62988
Y	-2	GLY	-	EXPRESSION TAG	UNP P62988
Y	-1	GLY	-	EXPRESSION TAG	UNP P62988
Y	0	SER	-	EXPRESSION TAG	UNP P62988


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin-conjugating enzyme E2 D2




- Molecule 1: Ubiquitin-conjugating enzyme E2 D2




- Molecule 2: E3 ubiquitin-protein ligase NEDD4-like




- Molecule 2: E3 ubiquitin-protein ligase NEDD4-like



- Molecule 3: Ubiquitin



- Molecule 3: Ubiquitin



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	173.98Å 199.89Å 109.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30 49.97 – 3.30	Depositor EDS
% Data completeness (in resolution range)	87.5 (50.00-3.30) 87.4 (49.97-3.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 3.33Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.217 , 0.267 0.220 , 0.230	Depositor DCC
$R_{free}$ test set	1288 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	74.2	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 25449 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	9814	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/1206	0.67	0/1642
1	B	0.41	0/1206	0.63	0/1642
2	C	0.49	0/3232	0.66	0/4366
2	D	0.48	1/3207 (0.0%)	0.72	1/4333 (0.0%)
3	X	0.40	0/607	0.70	0/816
3	Y	0.38	0/607	0.58	0/816
All	All	0.46	1/10065 (0.0%)	0.67	1/13615 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	837	ASP	CB-CG	5.05	1.62	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	764	GLN	N-CA-C	-5.95	94.94	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1170	0	1152	57	0
1	B	1170	0	1152	57	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	3148	0	3063	163	0
2	D	3124	0	3045	150	0
3	X	601	0	629	41	0
3	Y	601	0	629	40	0
All	All	9814	0	9670	480	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 480 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:597:ARG:HH11	2:D:597:ARG:HG3	1.25	1.00
2:C:817:ILE:O	2:C:820:ILE:HG22	1.61	1.00
2:C:675:ILE:H	2:C:675:ILE:HD12	1.26	0.97
2:C:848:ILE:HG13	2:C:906:THR:HG23	1.49	0.95
2:C:944:ALA:O	2:C:948:ALA:HB2	1.69	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/146 (99%)	133 (92%)	10 (7%)	1 (1%)	26	66
1	B	144/146 (99%)	124 (86%)	17 (12%)	3 (2%)	9	42
2	C	375/385 (97%)	321 (86%)	50 (13%)	4 (1%)	17	57
2	D	372/385 (97%)	328 (88%)	42 (11%)	2 (0%)	34	71
3	X	74/81 (91%)	64 (86%)	8 (11%)	2 (3%)	6	35
3	Y	74/81 (91%)	63 (85%)	9 (12%)	2 (3%)	6	35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1183/1224 (97%)	1033 (87%)	136 (12%)	14 (1%)	16	54

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	3	SER
2	D	825	GLU
1	B	129	THR
2	C	593	ASP
3	Y	2	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/130 (100%)	123 (95%)	7 (5%)	27	66
1	B	130/130 (100%)	118 (91%)	12 (9%)	11	40
2	C	341/347 (98%)	315 (92%)	26 (8%)	16	51
2	D	339/347 (98%)	306 (90%)	33 (10%)	10	38
3	X	68/70 (97%)	64 (94%)	4 (6%)	24	63
3	Y	68/70 (97%)	64 (94%)	4 (6%)	24	63
All	All	1076/1094 (98%)	990 (92%)	86 (8%)	15	49

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	841	ASN
2	D	604	ARG
3	X	64	GLU
2	C	887	ARG
2	C	932	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	856	ASN
2	D	596	ASN
3	Y	2	GLN
2	C	891	ASN
2	C	909	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/146 (100%)	0.11	0 <span>100</span> <span>100</span>	35, 80, 104, 114	0
1	B	146/146 (100%)	0.16	0 <span>100</span> <span>100</span>	52, 77, 99, 114	0
2	C	377/385 (97%)	0.04	4 (1%) <span>82</span> <span>78</span>	45, 72, 112, 122	0
2	D	374/385 (97%)	-0.03	0 <span>100</span> <span>100</span>	38, 67, 96, 126	0
3	X	76/81 (93%)	1.25	24 (31%) <span>1</span> <span>1</span>	93, 126, 127, 127	0
3	Y	76/81 (93%)	1.43	22 (28%) <span>1</span> <span>1</span>	95, 125, 127, 127	0
All	All	1195/1224 (97%)	0.21	50 (4%) <span>40</span> <span>33</span>	35, 75, 126, 127	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	X	1	MET	5.4
3	Y	66	THR	4.3
3	Y	15	LEU	4.2
3	Y	43	LEU	4.2
3	Y	34	GLU	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.