



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 05:54 PM GMT

PDB ID : 4JVI
Title : Crystal structure of PqsR co-inducer binding domain of *Pseudomonas aeruginosa* with inhibitor 3NH₂-7Cl-C9QZN
Authors : Ilangovan, A.; Williams, P.; Emsley, J.
Deposited on : 2013-03-25
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)


There are 3 unique types of molecules in this entry. The entry contains 1619 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator MvfR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	203	Total	C 1596	N 1011	O 277	S 302	0	0

- Molecule 2 is 3-AMINO-7-CHLORO-2-NONYLQUINAZOLIN-4(3H)-ONE (three-letter code: QZN) (formula: C₁₇H₂₄ClN₃O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C 22	Cl 17	N 1	O 3	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total 1 O 1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator MvfR



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.44Å 118.44Å 115.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.90 28.68 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-2.90) 99.3 (28.68-2.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ ¹	2.64 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R , R_{free}	0.204 , 0.263 0.217 , 0.238	Depositor DCC
R_{free} test set	531 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	101.1	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 76.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.47$, $< L^2 > = 0.31$	Xtriage
Outliers	0 of 10996 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1619	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: QZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.83	1/1626 (0.1%)	0.94	5/2206 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	174	PRO	N-CD	5.00	1.54	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	259	GLU	C-N-CD	5.97	140.94	128.40
1	A	104	PRO	C-N-CD	5.92	140.84	128.40
1	A	173	HIS	C-N-CD	5.83	140.63	128.40
1	A	209	ARG	C-N-CD	5.82	140.61	128.40
1	A	257	LEU	CA-CB-CG	-5.13	103.50	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1596	0	1590	140	2
2	A	22	0	24	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
All	All	1619	0	1614	140	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

All (140) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:GLN:HA	1:A:293:ARG:HD2	1.24	1.17
1:A:259:GLU:HG2	1:A:263:ILE:HA	1.39	1.01
1:A:162:VAL:O	1:A:289:ARG:NH1	1.95	0.98
1:A:98:LEU:HD11	1:A:127:THR:HG22	1.47	0.97
1:A:295:LEU:HD13	1:A:296:GLY:N	1.82	0.94
1:A:290:GLN:HA	1:A:293:ARG:CD	1.97	0.94
1:A:110:THR:O	1:A:114:VAL:HG23	1.67	0.94
1:A:290:GLN:CA	1:A:293:ARG:HD2	2.00	0.90
1:A:170:VAL:HG13	1:A:255:SER:CB	2.02	0.88
1:A:220:ASN:HD21	1:A:223:ASP:H	1.20	0.86
1:A:98:LEU:C	1:A:99:LEU:HD12	1.96	0.85
1:A:288:ALA:O	1:A:292:LEU:HD22	1.76	0.85
1:A:224:MET:O	1:A:228:VAL:HG12	1.77	0.84
1:A:98:LEU:HD12	1:A:99:LEU:N	1.93	0.83
1:A:160:GLN:HG3	1:A:270:TYR:CE1	2.14	0.83
1:A:294:GLU:OE1	1:A:295:LEU:N	2.12	0.82
1:A:182:SER:HA	1:A:185:SER:OG	1.78	0.81
1:A:170:VAL:HG13	1:A:255:SER:HB2	1.66	0.78
1:A:139:ASP:HA	1:A:157:ARG:NH2	1.99	0.78
1:A:98:LEU:HD11	1:A:127:THR:CG2	2.15	0.76
1:A:220:ASN:ND2	1:A:223:ASP:H	1.84	0.75
1:A:139:ASP:OD2	1:A:157:ARG:NH2	2.16	0.73
1:A:265:THR:HG21	2:A:401:QZN:H20	1.70	0.73
1:A:122:VAL:HG12	1:A:124:LEU:HD21	1.71	0.72
1:A:212:SER:O	1:A:215:VAL:CG2	2.39	0.70
1:A:170:VAL:HG22	1:A:254:LEU:HB2	1.74	0.69
1:A:253:VAL:HG12	1:A:255:SER:H	1.57	0.68
1:A:137:LYS:NZ	1:A:152:GLU:OE1	2.27	0.68
1:A:183:LEU:HD22	1:A:183:LEU:O	1.93	0.68
1:A:180:ASN:O	1:A:182:SER:OG	2.12	0.68
1:A:295:LEU:C	1:A:295:LEU:HD22	2.15	0.67
1:A:170:VAL:HG13	1:A:255:SER:HB3	1.75	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:VAL:H	1:A:255:SER:HB3	1.60	0.67
1:A:94:ASN:CB	1:A:121:MET:HG2	2.23	0.67
1:A:295:LEU:O	1:A:295:LEU:HD22	1.95	0.67
1:A:179:CYS:SG	1:A:180:ASN:N	2.68	0.66
1:A:183:LEU:C	1:A:183:LEU:HD13	2.16	0.66
1:A:225:LEU:HA	1:A:228:VAL:HG13	1.77	0.66
1:A:182:SER:O	1:A:185:SER:HB2	1.96	0.66
1:A:123:SER:C	1:A:124:LEU:HD23	2.16	0.66
1:A:122:VAL:CG1	1:A:124:LEU:HD21	2.26	0.66
1:A:113:SER:OG	1:A:291:ARG:NH2	2.28	0.66
1:A:105:PRO:HA	1:A:108:CYS:SG	2.36	0.65
1:A:167:LYS:NZ	1:A:264:ASP:OD2	2.30	0.65
1:A:182:SER:O	1:A:185:SER:N	2.30	0.65
1:A:213:ASP:N	1:A:213:ASP:OD1	2.30	0.64
1:A:209:ARG:HG2	1:A:209:ARG:HH11	1.62	0.64
1:A:132:SER:OG	1:A:148:THR:HG21	1.97	0.63
1:A:182:SER:O	1:A:185:SER:CB	2.48	0.62
1:A:139:ASP:HA	1:A:157:ARG:HH22	1.65	0.62
1:A:94:ASN:HB2	1:A:121:MET:HG2	1.82	0.62
1:A:200:ARG:HH11	1:A:200:ARG:CB	2.13	0.61
1:A:289:ARG:O	1:A:293:ARG:HG3	2.00	0.61
1:A:182:SER:HA	1:A:185:SER:CB	2.31	0.60
1:A:209:ARG:HH11	1:A:209:ARG:CG	2.15	0.60
1:A:122:VAL:HG12	1:A:124:LEU:CD2	2.33	0.59
1:A:294:GLU:OE1	1:A:295:LEU:CA	2.49	0.59
1:A:144:ASP:O	1:A:145:ILE:HG13	2.03	0.59
1:A:99:LEU:HD12	1:A:99:LEU:N	2.17	0.59
1:A:221:PHE:CE2	1:A:238:PRO:HD3	2.37	0.59
1:A:112:SER:O	1:A:116:LEU:HG	2.03	0.58
1:A:256:GLU:OE1	1:A:256:GLU:HA	2.04	0.57
1:A:98:LEU:HD12	1:A:99:LEU:H	1.69	0.57
1:A:228:VAL:O	1:A:231:GLY:HA3	2.05	0.57
1:A:212:SER:O	1:A:215:VAL:HG22	2.04	0.57
1:A:211:VAL:CG2	1:A:234:TRP:HH2	2.18	0.56
1:A:245:ARG:HG2	1:A:250:THR:HB	1.88	0.56
1:A:224:MET:O	1:A:228:VAL:CG1	2.50	0.56
1:A:194:GLN:HG3	1:A:234:TRP:CH2	2.40	0.56
1:A:199:SER:O	1:A:202:GLY:N	2.38	0.56
1:A:97:VAL:HG22	1:A:99:LEU:HD11	1.87	0.56
1:A:230:ALA:HB3	1:A:231:GLY:HA2	1.87	0.56
1:A:227:LEU:O	1:A:231:GLY:HA2	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:GLU:HG2	1:A:263:ILE:CA	2.27	0.55
1:A:211:VAL:HG21	1:A:234:TRP:HH2	1.72	0.54
1:A:242:VAL:HG12	1:A:242:VAL:O	2.08	0.54
1:A:115:LEU:O	1:A:119:PHE:HB2	2.08	0.54
1:A:97:VAL:CG2	1:A:99:LEU:HD11	2.39	0.53
1:A:188:SER:O	1:A:191:ASN:HB2	2.07	0.53
1:A:110:THR:HG21	1:A:292:LEU:HD13	1.90	0.53
1:A:211:VAL:HG21	1:A:234:TRP:CH2	2.44	0.52
1:A:97:VAL:HB	1:A:145:ILE:HB	1.92	0.52
1:A:220:ASN:C	1:A:220:ASN:ND2	2.63	0.52
1:A:113:SER:O	1:A:116:LEU:HB2	2.10	0.52
1:A:124:LEU:N	1:A:124:LEU:HD23	2.23	0.51
1:A:200:ARG:HH11	1:A:200:ARG:HB2	1.76	0.51
1:A:196:SER:HB3	1:A:215:VAL:HG13	1.92	0.50
1:A:184:HIS:O	1:A:187:ALA:N	2.43	0.50
1:A:211:VAL:CG2	1:A:234:TRP:CH2	2.94	0.50
1:A:220:ASN:HD22	1:A:220:ASN:C	2.15	0.49
1:A:211:VAL:HG23	1:A:212:SER:N	2.26	0.49
1:A:230:ALA:N	1:A:231:GLY:HA2	2.26	0.49
1:A:279:ARG:NH1	1:A:279:ARG:HB3	2.26	0.49
1:A:209:ARG:NH1	1:A:209:ARG:CG	2.73	0.49
1:A:159:ASN:HB2	1:A:271:TYR:CZ	2.47	0.49
1:A:144:ASP:C	1:A:145:ILE:HG13	2.31	0.48
1:A:182:SER:O	1:A:185:SER:CA	2.61	0.48
1:A:194:GLN:OE1	1:A:211:VAL:HG22	2.13	0.48
1:A:123:SER:O	1:A:124:LEU:HD23	2.12	0.48
1:A:140:ASN:ND2	1:A:141:ALA:N	2.61	0.48
1:A:182:SER:HA	1:A:185:SER:HG	1.76	0.48
1:A:182:SER:O	1:A:186:ILE:N	2.46	0.48
1:A:279:ARG:HH11	1:A:279:ARG:HB3	1.79	0.48
1:A:259:GLU:CG	1:A:263:ILE:HA	2.29	0.47
1:A:220:ASN:ND2	1:A:222:ASP:OD1	2.48	0.47
1:A:105:PRO:O	1:A:108:CYS:HB2	2.14	0.47
1:A:209:ARG:CZ	1:A:209:ARG:CB	2.91	0.47
1:A:212:SER:O	1:A:215:VAL:HG23	2.15	0.47
1:A:178:LEU:HD21	1:A:192:TYR:CE2	2.50	0.46
1:A:115:LEU:HA	1:A:115:LEU:HD23	1.73	0.46
1:A:295:LEU:HD13	1:A:296:GLY:H	1.69	0.46
1:A:155:ILE:HG22	1:A:156:SER:N	2.31	0.46
1:A:253:VAL:HG12	1:A:255:SER:N	2.29	0.45
1:A:100:ASP:OD2	1:A:100:ASP:C	2.55	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:LEU:HD13	1:A:184:HIS:N	2.31	0.44
1:A:109:ASP:OD2	1:A:109:ASP:N	2.50	0.44
1:A:285:LEU:O	1:A:289:ARG:HG3	2.18	0.43
1:A:194:GLN:HG3	1:A:234:TRP:CZ3	2.53	0.43
1:A:170:VAL:HG11	2:A:401:QZN:H5	2.01	0.43
1:A:292:LEU:O	1:A:295:LEU:HD12	2.19	0.43
1:A:196:SER:HB3	1:A:215:VAL:CG1	2.49	0.43
1:A:137:LYS:CE	1:A:152:GLU:OE1	2.67	0.42
1:A:165:TYR:CD2	1:A:165:TYR:N	2.87	0.42
1:A:183:LEU:C	1:A:183:LEU:HD22	2.37	0.42
1:A:140:ASN:C	1:A:140:ASN:ND2	2.73	0.42
1:A:227:LEU:HD12	1:A:227:LEU:HA	1.93	0.42
1:A:94:ASN:CA	1:A:121:MET:HG2	2.48	0.42
1:A:105:PRO:O	1:A:108:CYS:N	2.53	0.42
1:A:230:ALA:N	1:A:231:GLY:CA	2.83	0.42
1:A:295:LEU:C	1:A:295:LEU:HD13	2.37	0.42
1:A:259:GLU:HG3	1:A:259:GLU:O	2.20	0.42
1:A:133:LEU:HD13	1:A:152:GLU:HB2	2.02	0.41
1:A:173:HIS:CE1	1:A:175:GLN:HB2	2.55	0.41
1:A:163:LEU:HG	1:A:269:CYS:HB2	2.01	0.41
1:A:160:GLN:HG3	1:A:270:TYR:CZ	2.55	0.41
1:A:94:ASN:HA	1:A:121:MET:HG2	2.02	0.41
1:A:209:ARG:NH1	1:A:209:ARG:CB	2.84	0.40
1:A:127:THR:HG23	1:A:128:SER:O	2.22	0.40
1:A:98:LEU:HD12	1:A:98:LEU:C	2.41	0.40
1:A:182:SER:HA	1:A:185:SER:HB2	1.99	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:TYR:OH	1:A:259:GLU:OE2[9_555]	1.72	0.48
1:A:257:LEU:O	1:A:289:ARG:NH2[9_555]	1.80	0.40

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	201/216 (93%)	194 (96%)	7 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	178/189 (94%)	132 (74%)	46 (26%)	0 2

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	VAL
1	A	98	LEU
1	A	101	THR
1	A	121	MET
1	A	123	SER
1	A	140	ASN
1	A	142	GLU
1	A	148	THR
1	A	151	GLU
1	A	153	LEU
1	A	156	SER
1	A	170	VAL
1	A	175	GLN
1	A	180	ASN
1	A	182	SER
1	A	188	SER
1	A	189	LEU
1	A	197	LEU
1	A	200	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	SER
1	A	203	GLN
1	A	207	LEU
1	A	209	ARG
1	A	213	ASP
1	A	214	LYS
1	A	216	LEU
1	A	219	GLU
1	A	220	ASN
1	A	222	ASP
1	A	227	LEU
1	A	228	VAL
1	A	232	VAL
1	A	241	PHE
1	A	245	ARG
1	A	250	THR
1	A	256	GLU
1	A	264	ASP
1	A	280	SER
1	A	282	LEU
1	A	289	ARG
1	A	290	GLN
1	A	291	ARG
1	A	292	LEU
1	A	293	ARG
1	A	294	GLU
1	A	295	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	A	138	GLN
1	A	140	ASN
1	A	184	HIS
1	A	204	HIS
1	A	220	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	QZN	A	401	-	21,23,23	1.73	3 (14%)	18,30,30	1.24	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QZN	A	401	-	-	0/8/9/9	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	QZN	NAB-NAV	-6.06	1.24	1.41
2	A	401	QZN	CAG-CAT	-2.01	1.38	1.41
2	A	401	QZN	CAR-NAV	2.77	1.43	1.37

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	QZN	CAE-CAQ-CAG	-2.43	118.70	121.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	QZN	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/216 (93%)	-0.18	6 (2%) 54 47	68, 97, 140, 172	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	294	GLU	4.0
1	A	295	LEU	2.3
1	A	200	ARG	2.1
1	A	296	GLY	2.1
1	A	161	CYS	2.0
1	A	269	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no carbohydrates in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	QZN	A	401	22/22	0.96	0.20	0.73	80,88,105,110	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.