



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:54 AM GMT

PDB ID : 3K5K
Title : Discovery of a 2,4-Diamino-7-aminoalkoxy-quinazoline as a Potent Inhibitor of Histone Lysine Methyltransferase, G9a
Authors : Dong,A.; Wasney,G.A.; Liu,F.; Chen,X.; Allali-Hassani,A.; Senisterra,G.; Chau,I.; Bountra,C.; Weigelt,J.; Edwards,A.M.; Arrowsmith,C.H.; Frye,S.V.; Bochkarev,A.; Brown,P.J.; Jin,J.; Vedadi,M.
Deposited on : 2009-10-07
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	<div> <div>6%</div> <div>92%</div> <div>• •</div> </div>
1	B	283	<div> <div>10%</div> <div>93%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	A	3001	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase, H3 lysine-9 specific 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	11	3	0
			2126	1329	370	404	23			
1	B	269	Total	C	N	O	S	6	1	0
			2095	1307	362	403	23			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	911	GLY	-	EXPRESSION TAG	UNP Q96KQ7
A	912	SER	-	EXPRESSION TAG	UNP Q96KQ7
B	911	GLY	-	EXPRESSION TAG	UNP Q96KQ7
B	912	SER	-	EXPRESSION TAG	UNP Q96KQ7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	Zn	0	0
			4	4		
2	A	4	Total	Zn	0	0
			4	4		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).



-
- Chemical structure of DXQ, a complex molecule featuring a central benzimidazole core. The structure includes several fused and linked rings, with various substituents labeled with three-letter codes. Key features include a benzimidazole core with a benzene ring (C4, C5, C6) and an imidazole ring (N1, N3). Substituents include a large amide-like group (CAB, NBF, CAM, CAH, CAJ, OAX, OAY, OAW, CAA) and a complex side chain (CAG, CAF, C4, C5, C6, N1, N3, C2, NBI, CAS, CAR, NBG(S), CAN, CAI, CAO, CAK, CAP, CBE, CAL, CAQ, CBH, CAE). The structure is highly branched and contains multiple nitrogen atoms (N1, N3, NBI, NBH, NBF, NBG(S)) and oxygen atoms (OAX, OAY, OAW).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			35	26	7	2		



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			35	26	7	2		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Cl	0	0
			2	2		
5	A	2	Total	Cl	0	0
			2	2		


- Molecule 6 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	X	0	0
			1	1		
6	A	1	Total	X	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	236	Total	O	0	0
			236	236		
7	B	223	Total	O	0	0
			223	223		

- Molecule 1: Histone-lysine N-methyltransferase, H3 lysine-9 specific 3



- Chain B:  10% 93% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.67Å 78.07Å 72.51Å 90.00° 91.81° 90.00°	Depositor
Resolution (Å)	29.58 – 1.70 28.32 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.0 (29.58-1.70) 97.0 (28.32-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0102, COOT 0.5.2	Depositor
R, R_{free}	0.209 , 0.264 0.209 , 0.265	Depositor DCC
R_{free} test set	1367 reflections (2.07%)	DCC
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.9	EDS
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 67674 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4816	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.86 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.4178e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, ZN, SAH, DXQ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/2171	0.60	0/2946
1	B	0.54	0/2140	0.60	0/2906
All	All	0.54	0/4311	0.60	0/5852

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2126	0	1928	7	0
1	B	2095	0	1895	3	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
4	A	35	0	43	1	0
4	B	35	0	43	2	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	1	0	0	0	0
7	A	236	0	0	1	0
7	B	223	0	0	2	0
All	All	4816	0	3947	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:919:GLU:HG3	1:B:940:GLY:HA3	1.75	0.68
1:B:919:GLU:CD	1:B:1053:ARG:HH22	2.15	0.49
1:A:1060:GLN:HG2	7:A:208:HOH:O	2.13	0.48
4:B:2001:DXQ:HAP	7:B:344:HOH:O	2.13	0.47
1:A:962:MET:HE3	1:A:1133:PHE:CE1	2.53	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/283 (96%)	261 (96%)	10 (4%)	0	100	100
1	B	266/283 (94%)	253 (95%)	13 (5%)	0	100	100
All	All	537/566 (95%)	514 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/253 (87%)	219 (100%)	1 (0%)	92	88
1	B	221/253 (87%)	221 (100%)	0	100	100
All	All	441/506 (87%)	440 (100%)	1 (0%)	95	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	991	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	972	GLN
1	A	1060	GLN
1	B	1091	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 2 are unknown and 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SAH	A	1198	-	20,28,28	1.23	2 (10%)	19,40,40	2.33	4 (21%)
4	DXQ	A	2000	-	36,38,38	1.17	3 (8%)	46,52,52	2.54	12 (26%)
3	SAH	B	1198	-	20,28,28	1.26	4 (20%)	19,40,40	2.14	3 (15%)
4	DXQ	B	2001	-	36,38,38	1.26	4 (11%)	46,52,52	2.48	16 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	A	1198	-	-	0/7/31/31	0/3/3/3
4	DXQ	A	2000	-	-	0/13/38/38	0/3/4/4
3	SAH	B	1198	-	-	0/7/31/31	0/3/3/3
4	DXQ	B	2001	-	-	0/13/38/38	0/3/4/4

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	2001	DXQ	C6-C5	-4.21	1.39	1.44
4	A	2000	DXQ	C6-C5	-3.62	1.40	1.44
3	B	1198	SAH	C5'-SD	-2.33	1.76	1.81
4	B	2001	DXQ	CAG-CAY	2.01	1.40	1.36
3	B	1198	SAH	O4'-C1'	2.03	1.43	1.41

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2001	DXQ	CAG-C5-C6	-8.80	120.20	124.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2000	DXQ	CAG-C5-C6	-8.75	120.22	124.89
3	B	1198	SAH	N3-C2-N1	-7.96	122.80	128.89
3	A	1198	SAH	N3-C2-N1	-7.83	122.90	128.89
4	A	2000	DXQ	C5-C4-N3	-4.44	118.56	122.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2000	DXQ	1	0
4	B	2001	DXQ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/283 (96%)	0.42	18 (6%) 22 23	23, 33, 48, 68	2 (0%)
1	B	269/283 (95%)	0.46	27 (10%) 9 9	19, 33, 52, 67	1 (0%)
All	All	541/566 (95%)	0.44	45 (8%) 14 16	19, 33, 51, 68	3 (0%)

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1108[A]	SER	9.4
1	A	946	CYS	8.5
1	A	1189	LEU	7.4
1	A	977	VAL	6.7
1	A	1091	ASN	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	CL	A	3001	1/1	0.99	0.14	3.55	23,23,23,23	0
4	DXQ	B	2001	35/35	0.81	0.15	0.73	37,40,51,53	0
3	SAH	B	1198	26/26	0.93	0.11	0.42	29,31,33,34	0
4	DXQ	A	2000	35/35	0.88	0.11	-0.07	35,43,52,53	0
3	SAH	A	1198	26/26	0.94	0.08	-0.38	29,31,34,35	0
5	CL	B	3003	1/1	0.98	0.10	-0.45	28,28,28,28	0
2	ZN	B	1196	1/1	0.99	0.07	-0.70	29,29,29,29	0
2	ZN	B	1194	1/1	0.99	0.06	-0.91	27,27,27,27	0
2	ZN	A	1194	1/1	0.99	0.06	-1.51	33,33,33,33	0
2	ZN	B	1195	1/1	0.99	0.05	-1.52	27,27,27,27	0
2	ZN	B	1197	1/1	0.98	0.04	-2.14	37,37,37,37	0
2	ZN	A	1197	1/1	0.99	0.03	-2.43	37,37,37,37	0
2	ZN	A	1195	1/1	0.99	0.04	-2.80	32,32,32,32	0
2	ZN	A	1196	1/1	0.99	0.02	-4.66	35,35,35,35	0
5	CL	A	3004	1/1	0.88	0.10	-	59,59,59,59	0
6	UNX	B	1199	1/1	0.82	0.34	-	30,30,30,30	0
6	UNX	A	1199	1/1	0.88	0.34	-	30,30,30,30	0
5	CL	B	3005	1/1	0.90	0.08	-	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.