



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:55 AM GMT

PDB ID : 3K7D  
Title : C-terminal (adenylation) domain of E.coli Glutamine Synthetase Adenylyl-transferase  
Authors : Xu, Y.; Carr, P.D.; Vasudevan, S.G.; Ollis, D.L.  
Deposited on : 2009-10-12  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 85%, yellow 85%, yellow 98%, green 98%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>85%</span> <span>13%</span> <span>•</span> </div> </div>
1	B	498	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 6%, orange 6%, orange 81%, yellow 81%, yellow 96%, green 96%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>6%</span> <span>81%</span> <span>15%</span> <span>• •</span> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8407 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate-ammonia-ligase adenylyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	497	Total	C	N	O	S	0	0	0
			4005	2522	715	750	18			
1	B	495	Total	C	N	O	S	0	0	0
			3957	2497	695	747	18			

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	195	Total	O	0	0
			195	195		
3	B	185	Total	O	0	0
			185	185		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate-ammonia-ligase adenylyltransferase



- Molecule 1: Glutamate-ammonia-ligase adenylyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.54Å 122.86Å 143.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.13 – 2.40 47.13 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.4 (47.13-2.40) 96.4 (47.13-2.40)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.13 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.183 , 0.235 0.187 , 0.235	Depositor DCC
$R_{free}$ test set	2504 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 49134 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8407	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/4084	0.59	0/5537
1	B	0.46	0/4035	0.58	1/5476 (0.0%)
All	All	0.47	0/8119	0.59	1/11013 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	501	ARG	NE-CZ-NH1	6.49	123.54	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	836	GLY	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4005	0	3961	50	0
1	B	3957	0	3875	54	0
2	A	40	0	0	0	0
2	B	25	0	0	0	0
3	A	195	0	0	3	0
3	B	185	0	0	2	0
All	All	8407	0	7836	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:586:TYR:O	1:A:636:THR:HG21	1.39	1.15
1:B:462:GLU:HA	1:B:481:LEU:HD22	1.44	0.96
1:A:899:ALA:HB1	1:A:934:LEU:HD11	1.47	0.95
1:A:614:LEU:HD11	1:A:738:LEU:HD22	1.63	0.79
1:B:945:GLU:O	1:B:946:GLU:HB2	1.88	0.74
1:B:837:ASN:HB2	1:B:844:ASP:HB2	1.69	0.74
1:B:495:THR:HA	1:B:501:ARG:HH12	1.53	0.73
1:A:497:GLY:O	1:A:501:ARG:HG3	1.92	0.69
1:B:925:GLU:HA	1:B:925:GLU:OE1	1.92	0.68
1:B:462:GLU:CA	1:B:481:LEU:HD22	2.22	0.67
1:A:598:LEU:HD22	1:A:653:MET:HE1	1.79	0.65
1:A:586:TYR:O	1:A:636:THR:CG2	2.32	0.65
1:B:838:LYS:O	1:B:839:HIS:HB2	1.96	0.65
1:B:598:LEU:HD22	1:B:653:MET:HE2	1.79	0.63
1:A:631:ALA:HB1	1:A:636:THR:HG22	1.81	0.63
1:B:695:GLU:HG3	1:B:904:TYR:OH	1.99	0.63
1:A:496:ILE:HD11	1:A:501:ARG:HG2	1.81	0.63
1:B:598:LEU:HD22	1:B:653:MET:CE	2.29	0.62
1:A:573:PRO:O	1:A:576:LEU:HG	2.00	0.62
1:B:496:ILE:HG23	1:B:497:GLY:H	1.65	0.61
1:A:551:ALA:HB2	1:A:580:LEU:HD23	1.82	0.61
1:B:494:ARG:O	1:B:495:THR:O	2.19	0.59
1:B:836:GLY:HA2	1:B:847:ALA:CB	2.32	0.58
1:B:836:GLY:HA2	1:B:847:ALA:HB1	1.85	0.58
1:B:595:ARG:O	1:B:599:ARG:HG3	2.04	0.58
1:B:608:ASP:N	1:B:608:ASP:OD1	2.37	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:TYR:CE2	1:A:573:PRO:HB2	2.39	0.57
1:A:840:ARG:NH1	1:A:840:ARG:HB2	2.18	0.57
1:A:583:ASN:O	1:A:587:GLN:HG3	2.06	0.56
1:A:674:HIS:HE1	3:A:232:HOH:O	1.88	0.55
1:B:496:ILE:HG23	1:B:497:GLY:N	2.23	0.54
1:B:494:ARG:O	1:B:495:THR:C	2.46	0.54
1:B:933:GLU:OE1	1:B:936:ARG:NH1	2.39	0.54
1:A:838:LYS:NZ	1:B:750:GLU:OE1	2.24	0.54
1:B:945:GLU:O	1:B:946:GLU:CB	2.54	0.54
1:A:496:ILE:CD1	1:A:501:ARG:HG2	2.38	0.53
1:B:614:LEU:HB3	1:B:743:THR:HG21	1.90	0.52
1:B:829:GLU:OE2	1:B:832:ARG:NH1	2.41	0.52
1:A:471:HIS:HD2	3:A:156:HOH:O	1.93	0.52
1:A:676:ASN:O	1:A:677:GLU:HB2	2.09	0.52
1:A:766:THR:HG22	1:A:770:ALA:HB3	1.91	0.51
1:B:819:THR:O	1:B:823:GLU:HG3	2.11	0.50
1:A:826:GLU:HB2	1:B:780:TRP:CD2	2.46	0.50
1:A:664:GLN:HG3	3:A:369:HOH:O	2.11	0.50
1:B:708:HIS:O	1:B:767:SER:HA	2.13	0.49
1:A:873:LEU:HD22	1:A:886:LEU:HD13	1.94	0.49
1:B:606:PRO:O	1:B:613:GLN:NE2	2.45	0.49
1:A:615:GLU:OE2	1:A:745:SER:OG	2.31	0.49
1:A:840:ARG:HH11	1:A:840:ARG:HB2	1.77	0.48
1:A:491:LEU:CD2	1:A:536:VAL:HG21	2.43	0.48
1:B:740:SER:HB3	3:B:190:HOH:O	2.13	0.48
1:A:695:GLU:O	1:A:695:GLU:HG2	2.13	0.48
1:B:796:ASP:HB2	1:B:797:PRO:CD	2.44	0.48
1:B:834:HIS:O	1:B:835:LEU:HD12	2.13	0.48
1:B:547:GLU:HB2	1:B:548:PHE:CD2	2.49	0.47
1:A:614:LEU:CD1	1:A:738:LEU:HD22	2.40	0.47
1:A:598:LEU:HD22	1:A:653:MET:CE	2.43	0.47
1:A:614:LEU:HD11	1:A:738:LEU:CD2	2.41	0.47
1:B:806:VAL:O	1:B:810:ILE:HG12	2.15	0.46
1:A:859:THR:HB	1:A:882:ILE:HG21	1.98	0.46
1:B:455:LEU:HD12	1:B:468:VAL:HG21	1.96	0.46
1:A:608:ASP:N	1:A:608:ASP:OD1	2.42	0.46
1:B:574:LEU:O	1:B:574:LEU:HG	2.15	0.45
1:B:452:TRP:CZ3	1:B:513:SER:HA	2.52	0.45
1:B:455:LEU:HD22	1:B:484:ILE:HD13	1.97	0.45
1:A:591:THR:HB	1:A:648:TRP:CD1	2.52	0.45
1:A:514:ASP:C	1:A:514:ASP:OD1	2.56	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:496:ILE:HD13	1:B:496:ILE:C	2.38	0.44
1:A:541:TYR:CD2	1:A:573:PRO:HB2	2.53	0.43
1:A:796:ASP:HB2	1:A:797:PRO:CD	2.49	0.43
1:A:772:ALA:O	1:A:776:LYS:HG3	2.17	0.43
1:B:838:LYS:O	1:B:839:HIS:CB	2.64	0.43
1:B:832:ARG:NH2	1:B:854:ASP:OD1	2.51	0.43
1:A:541:TYR:HE2	1:A:573:PRO:HB2	1.79	0.43
1:A:471:HIS:H	1:A:471:HIS:CD2	2.37	0.43
1:A:872:LYS:O	1:A:875:ARG:HG3	2.19	0.43
1:B:669:TYR:OH	1:B:730:ARG:HD3	2.18	0.43
1:B:614:LEU:HD12	1:B:614:LEU:HA	1.86	0.42
1:B:664:GLN:OE1	1:B:734:ARG:NE	2.51	0.42
1:A:456:TRP:CE2	1:A:488:ARG:NH1	2.87	0.42
1:A:494:ARG:NH2	1:A:533:VAL:O	2.52	0.42
1:A:479:GLN:HA	1:A:479:GLN:HE21	1.84	0.42
1:B:707:LEU:HA	1:B:766:THR:O	2.20	0.42
1:A:668:ARG:HG2	1:A:669:TYR:CE1	2.54	0.42
1:A:489:LYS:HD3	1:B:493:LYS:HG3	2.02	0.42
1:A:934:LEU:C	1:A:934:LEU:HD23	2.41	0.42
1:B:756:ARG:HD2	1:B:763:MET:O	2.19	0.41
1:B:508:MET:N	1:B:509:PRO:CD	2.83	0.41
1:B:618:ARG:NH1	1:B:748:LEU:HB2	2.35	0.41
1:B:628:ILE:HD13	1:B:646:LEU:HD23	2.03	0.41
1:B:538:ARG:NH2	3:B:251:HOH:O	2.52	0.41
1:A:828:ARG:HG2	1:A:939:TRP:CH2	2.56	0.41
1:A:826:GLU:HB2	1:B:780:TRP:CE3	2.56	0.41
1:B:743:THR:C	1:B:745:SER:H	2.23	0.41
1:A:503:VAL:HG21	1:A:543:GLU:HG3	2.04	0.41
1:A:831:MET:O	1:A:835:LEU:HB2	2.21	0.41
1:A:687:GLY:O	1:A:790:ALA:HA	2.22	0.40
1:B:466:THR:HA	1:B:467:PRO:HD3	1.88	0.40
1:B:859:THR:HB	1:B:882:ILE:HG21	2.03	0.40
1:B:620:PHE:O	1:B:624:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/498 (99%)	489 (99%)	6 (1%)	0	100	100
1	B	491/498 (99%)	477 (97%)	10 (2%)	4 (1%)	24	35
All	All	986/996 (99%)	966 (98%)	16 (2%)	4 (0%)	39	56

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	495	THR
1	B	839	HIS
1	B	925	GLU
1	B	744	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/425 (99%)	400 (95%)	21 (5%)	30	48
1	B	412/425 (97%)	388 (94%)	24 (6%)	25	39
All	All	833/850 (98%)	788 (95%)	45 (5%)	27	43

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	479	GLN
1	A	496	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	536	VAL
1	A	545	LEU
1	A	555	LEU
1	A	587	GLN
1	A	607	GLU
1	A	608	ASP
1	A	611	GLU
1	A	614	LEU
1	A	636	THR
1	A	644	ASP
1	A	789	ARG
1	A	799	LEU
1	A	826	GLU
1	A	840	ARG
1	A	883	LEU
1	A	886	LEU
1	A	894	GLU
1	A	934	LEU
1	A	943	LEU
1	B	455	LEU
1	B	462	GLU
1	B	490	GLU
1	B	496	ILE
1	B	533	VAL
1	B	540	THR
1	B	580	LEU
1	B	607	GLU
1	B	608	ASP
1	B	611	GLU
1	B	614	LEU
1	B	633	ILE
1	B	640	MET
1	B	696	LEU
1	B	726	GLN
1	B	730	ARG
1	B	744	SER
1	B	745	SER
1	B	778	GLU
1	B	835	LEU
1	B	837	ASN
1	B	883	LEU
1	B	894	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	934	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	457	GLN
1	A	471	HIS
1	A	479	GLN
1	A	506	HIS
1	A	674	HIS
1	A	888	GLN
1	B	777	ASN
1	B	868	HIS
1	B	888	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	1	-	4,4,4	0.30	0	6,6,6	0.48	0
2	SO4	A	12	-	4,4,4	0.22	0	6,6,6	0.19	0
2	SO4	A	13	-	4,4,4	0.22	0	6,6,6	0.11	0
2	SO4	A	14	-	4,4,4	0.20	0	6,6,6	0.18	0
2	SO4	A	16	-	4,4,4	0.16	0	6,6,6	0.34	0
2	SO4	A	3	-	4,4,4	0.25	0	6,6,6	0.39	0
2	SO4	A	5	-	4,4,4	0.13	0	6,6,6	0.27	0
2	SO4	A	6	-	4,4,4	0.22	0	6,6,6	0.20	0
2	SO4	B	10	-	4,4,4	0.16	0	6,6,6	0.12	0
2	SO4	B	11	-	4,4,4	0.09	0	6,6,6	0.32	0
2	SO4	B	15	-	4,4,4	0.22	0	6,6,6	0.48	0
2	SO4	B	7	-	4,4,4	0.16	0	6,6,6	0.20	0
2	SO4	B	8	-	4,4,4	0.14	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1	-	-	0/0/0/0	0/0/0/0
2	SO4	A	12	-	-	0/0/0/0	0/0/0/0
2	SO4	A	13	-	-	0/0/0/0	0/0/0/0
2	SO4	A	14	-	-	0/0/0/0	0/0/0/0
2	SO4	A	16	-	-	0/0/0/0	0/0/0/0
2	SO4	A	3	-	-	0/0/0/0	0/0/0/0
2	SO4	A	5	-	-	0/0/0/0	0/0/0/0
2	SO4	A	6	-	-	0/0/0/0	0/0/0/0
2	SO4	B	10	-	-	0/0/0/0	0/0/0/0
2	SO4	B	11	-	-	0/0/0/0	0/0/0/0
2	SO4	B	15	-	-	0/0/0/0	0/0/0/0
2	SO4	B	7	-	-	0/0/0/0	0/0/0/0
2	SO4	B	8	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	497/498 (99%)	-0.06	6 (1%) 81 81	5, 15, 30, 50	0
1	B	495/498 (99%)	0.15	32 (6%) 22 22	6, 15, 31, 53	0
All	All	992/996 (99%)	0.05	38 (3%) 44 45	5, 15, 31, 53	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	492	ASP	7.1
1	B	496	ILE	4.9
1	B	498	PRO	4.2
1	B	493	LYS	3.8
1	B	491	LEU	3.8
1	B	485	ALA	3.5
1	A	492	ASP	3.3
1	B	470	ALA	3.3
1	B	475	ASP	3.0
1	B	455	LEU	3.0
1	B	479	GLN	3.0
1	B	472	LEU	2.9
1	B	946	GLU	2.8
1	B	494	ARG	2.7
1	B	507	LEU	2.6
1	B	550	ALA	2.6
1	B	719	GLU	2.5
1	B	836	GLY	2.5
1	A	459	ALA	2.5
1	B	833	ALA	2.5
1	B	465	THR	2.4
1	A	677	GLU	2.4
1	A	495	THR	2.3
1	B	608	ASP	2.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	482	THR	2.3
1	B	477	ARG	2.3
1	B	495	THR	2.2
1	A	719	GLU	2.2
1	B	506	HIS	2.2
1	B	504	LEU	2.2
1	B	451	GLN	2.2
1	B	483	LEU	2.1
1	B	837	ASN	2.1
1	B	464	ASP	2.1
1	B	454	GLU	2.1
1	A	493	LYS	2.0
1	B	480	VAL	2.0
1	B	835	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	13	5/5	0.96	0.20	1.47	45,46,46,47	4
2	SO4	A	16	5/5	0.83	0.21	0.47	67,70,71,71	0
2	SO4	B	15	5/5	0.94	0.17	-0.22	64,66,66,68	0
2	SO4	A	1	5/5	0.99	0.14	-0.69	29,30,32,34	0
2	SO4	B	8	5/5	0.85	0.22	-	81,82,83,84	0
2	SO4	B	11	5/5	0.86	0.21	-	66,68,71,72	0
2	SO4	A	12	5/5	0.91	0.36	-	73,73,74,76	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	14	5/5	0.96	0.11	-	58,59,61,63	0
2	SO4	A	6	5/5	0.91	0.23	-	73,76,77,77	0
2	SO4	A	3	5/5	0.99	0.11	-	23,25,27,29	0
2	SO4	B	7	5/5	0.97	0.15	-	45,51,51,54	0
2	SO4	A	5	5/5	0.94	0.20	-	60,60,62,64	0
2	SO4	B	10	5/5	0.88	0.20	-	87,88,88,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.