



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 28, 2016 – 06:42 PM EST

PDB ID : 5K7X  
Title : Fully ligated Adenylosuccinate Synthetase from *Pyrococcus horikoshii* OT3  
with GTP, IMP and Hadacidin  
Authors : Xie, Y.; Cheng, Z.  
Deposited on : 2016-05-27  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028320  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028320


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 74%, green 22%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>74%</span> <span>22%</span> <span>..</span> </div> </div>
1	B	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 77%, green 22%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>77%</span> <span>22%</span> <span>.</span> </div> </div>
1	C	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 70%, green 25%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>70%</span> <span>25%</span> <span>..</span> </div> </div>
1	D	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 5%, orange 1%, yellow 67%, green 29%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>5%</span> <span>67%</span> <span>29%</span> <span>..</span> </div> </div>
1	E	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 4%, orange 1%, yellow 76%, green 21%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>4%</span> <span>76%</span> <span>21%</span> <span>..</span> </div> </div>
1	F	340	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 78%, green 21%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>78%</span> <span>21%</span> <span>..</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMP	C	401	-	-	-	X
4	HDA	A	402	-	-	X	-
4	HDA	B	402	-	-	X	X
4	HDA	D	402	-	-	X	-
4	HDA	E	402	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 16066 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenylosuccinate synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			
1	B	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			
1	C	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			
1	D	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			
1	E	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			
1	F	337	Total	C	N	O	S	0	0	0
			2604	1647	450	498	9			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP O58187
B	0	HIS	-	expression tag	UNP O58187
C	0	HIS	-	expression tag	UNP O58187
D	0	HIS	-	expression tag	UNP O58187
E	0	HIS	-	expression tag	UNP O58187
F	0	HIS	-	expression tag	UNP O58187

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	E	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	F	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 3 is INOSINIC ACID (three-letter code: IMP) (formula: C<sub>10</sub>H<sub>13</sub>N<sub>4</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
3	B	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
3	C	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
3	D	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
3	E	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
3	F	1	Total	C	N	O	P	0	0
			23	10	4	8	1		

- Molecule 4 is HADACIDIN (three-letter code: HDA) (formula: C<sub>3</sub>H<sub>5</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	3	1	4		
4	B	1	Total	C	N	O	0	0
			8	3	1	4		
4	C	1	Total	C	N	O	0	0
			8	3	1	4		
4	D	1	Total	C	N	O	0	0
			8	3	1	4		
4	E	1	Total	C	N	O	0	0
			8	3	1	4		
4	F	1	Total	C	N	O	0	0
			8	3	1	4		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	Mg	0	0
			1	1		
5	E	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		
5	F	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	10	Total 10	O 10	0	0
6	B	13	Total 13	O 13	0	0
6	C	12	Total 12	O 12	0	0
6	D	5	Total 5	O 5	0	0
6	E	6	Total 6	O 6	0	0
6	F	12	Total 12	O 12	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adenylosuccinate synthetase



- Molecule 1: Adenylosuccinate synthetase




- Molecule 1: Adenylosuccinate synthetase









• Molecule 1: Adenylosuccinate synthetase



• Molecule 1: Adenylosuccinate synthetase



• Molecule 1: Adenylosuccinate synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.35Å 130.55Å 168.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.12 – 2.80 39.12 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.5 (39.12-2.80) 95.5 (39.12-2.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.206 , 0.263 0.206 , 0.263	Depositor DCC
$R_{free}$ test set	2691 reflections (4.88%)	DCC
Wilson B-factor (Å <sup>2</sup> )	48.6	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	16066	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, IMP, MG, HDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/2653	0.58	5/3586 (0.1%)
1	B	0.35	0/2653	0.52	3/3586 (0.1%)
1	C	0.36	1/2653 (0.0%)	0.59	7/3586 (0.2%)
1	D	0.38	1/2653 (0.0%)	0.63	4/3586 (0.1%)
1	E	0.37	0/2653	0.61	4/3586 (0.1%)
1	F	0.29	0/2653	0.50	1/3586 (0.0%)
All	All	0.36	2/15918 (0.0%)	0.57	24/21516 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	154	PRO	N-CD	5.72	1.55	1.47
1	D	238	PRO	N-CD	5.20	1.55	1.47

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	33	ALA	CB-CA-C	-15.01	87.59	110.10
1	A	258	ARG	CB-CA-C	12.33	135.06	110.40
1	C	150	LYS	CB-CA-C	-9.31	91.78	110.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	65	LYS	CB-CA-C	9.27	128.94	110.40
1	E	66	ALA	N-CA-CB	-8.82	97.75	110.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	336	ARG	Sidechain
1	E	299	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2600	81	0
1	B	2604	0	2599	76	0
1	C	2604	0	2600	106	2
1	D	2604	0	2600	108	0
1	E	2604	0	2599	79	0
1	F	2604	0	2600	71	2
2	A	32	0	12	5	0
2	B	32	0	12	5	0
2	C	32	0	12	5	0
2	D	32	0	12	5	0
2	E	32	0	12	5	0
2	F	32	0	12	5	0
3	A	23	0	11	4	0
3	B	23	0	11	3	0
3	C	23	0	11	2	0
3	D	23	0	11	4	0
3	E	23	0	11	4	0
3	F	23	0	11	3	0
4	A	8	0	4	7	0
4	B	8	0	4	4	0
4	C	8	0	4	3	0
4	D	8	0	4	4	0
4	E	8	0	4	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	8	0	4	2	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	A	10	0	0	1	0
6	B	13	0	0	3	0
6	C	12	0	0	1	0
6	D	5	0	0	5	0
6	E	6	0	0	1	0
6	F	12	0	0	3	0
All	All	16066	0	15760	502	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 502 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:27:ASP:OD2	1:E:336:ARG:NH1	1.58	1.34
1:E:298:TYR:HB2	6:E:503:HOH:O	1.44	1.15
1:C:166:ALA:O	1:C:171:SER:HB2	1.50	1.12
1:A:43:HIS:CD2	2:A:400:GTP:O2G	2.07	1.08
1:B:338:THR:HG21	1:E:338:THR:CG2	1.85	1.07

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:337:ASP:OD2	1:F:290:LYS:CD[2_454]	1.60	0.60
1:C:337:ASP:OD2	1:F:290:LYS:CE[2_454]	1.96	0.24

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/340 (98%)	318 (95%)	17 (5%)	0	100	100
1	B	335/340 (98%)	317 (95%)	18 (5%)	0	100	100
1	C	335/340 (98%)	310 (92%)	23 (7%)	2 (1%)	30	65
1	D	335/340 (98%)	315 (94%)	17 (5%)	3 (1%)	21	55
1	E	335/340 (98%)	317 (95%)	16 (5%)	2 (1%)	30	65
1	F	335/340 (98%)	318 (95%)	17 (5%)	0	100	100
All	All	2010/2040 (98%)	1895 (94%)	108 (5%)	7 (0%)	46	79

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	40	ASN
1	D	337	ASP
1	C	40	ASN
1	C	160	ALA
1	D	239	MET

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/278 (99%)	259 (94%)	16 (6%)	25	57
1	B	275/278 (99%)	271 (98%)	4 (2%)	72	93
1	C	275/278 (99%)	265 (96%)	10 (4%)	42	76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	275/278 (99%)	265 (96%)	10 (4%)	42	76
1	E	275/278 (99%)	263 (96%)	12 (4%)	35	69
1	F	275/278 (99%)	269 (98%)	6 (2%)	60	89
All	All	1650/1668 (99%)	1592 (96%)	58 (4%)	43	77

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	198	VAL
1	D	123	LYS
1	F	198	VAL
1	C	257	ARG
1	C	335	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	161	GLN
1	C	112	GLN
1	D	112	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 6 are monoatomic - leaving 18 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTP	A	400	5	26,34,34	0.94	1 (3%)	29,54,54	1.66	4 (13%)
3	IMP	A	401	-	21,25,25	1.23	3 (14%)	22,38,38	2.45	3 (13%)
4	HDA	A	402	5	2,7,7	1.10	0	1,8,8	2.97	1 (100%)
2	GTP	B	400	5	26,34,34	0.95	1 (3%)	29,54,54	1.64	3 (10%)
3	IMP	B	401	-	21,25,25	1.23	3 (14%)	22,38,38	2.44	3 (13%)
4	HDA	B	402	5	2,7,7	1.11	0	1,8,8	2.86	1 (100%)
2	GTP	C	400	5	26,34,34	0.95	1 (3%)	29,54,54	1.57	3 (10%)
3	IMP	C	401	-	21,25,25	1.24	3 (14%)	22,38,38	2.44	3 (13%)
4	HDA	C	402	5	2,7,7	0.96	0	1,8,8	1.86	0
2	GTP	D	400	5	26,34,34	0.93	1 (3%)	29,54,54	1.72	4 (13%)
3	IMP	D	401	-	21,25,25	1.23	3 (14%)	22,38,38	2.56	3 (13%)
4	HDA	D	402	5	2,7,7	1.32	0	1,8,8	1.38	0
2	GTP	E	400	5	26,34,34	0.93	1 (3%)	29,54,54	1.64	3 (10%)
3	IMP	E	401	-	21,25,25	1.23	3 (14%)	22,38,38	2.48	3 (13%)
4	HDA	E	402	5	2,7,7	1.20	0	1,8,8	2.41	1 (100%)
2	GTP	F	400	5	26,34,34	0.96	1 (3%)	29,54,54	1.65	3 (10%)
3	IMP	F	401	-	21,25,25	1.25	3 (14%)	22,38,38	2.54	3 (13%)
4	HDA	F	402	5	2,7,7	1.18	0	1,8,8	3.17	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	A	400	5	-	0/18/38/38	0/3/3/3
3	IMP	A	401	-	-	0/6/26/26	0/3/3/3
4	HDA	A	402	5	-	0/1/6/6	0/0/0/0
2	GTP	B	400	5	-	0/18/38/38	0/3/3/3
3	IMP	B	401	-	-	0/6/26/26	0/3/3/3
4	HDA	B	402	5	-	0/1/6/6	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	C	400	5	-	0/18/38/38	0/3/3/3
3	IMP	C	401	-	-	0/6/26/26	0/3/3/3
4	HDA	C	402	5	-	0/1/6/6	0/0/0/0
2	GTP	D	400	5	-	0/18/38/38	0/3/3/3
3	IMP	D	401	-	-	0/6/26/26	0/3/3/3
4	HDA	D	402	5	-	0/1/6/6	0/0/0/0
2	GTP	E	400	5	-	0/18/38/38	0/3/3/3
3	IMP	E	401	-	-	0/6/26/26	0/3/3/3
4	HDA	E	402	5	-	0/1/6/6	0/0/0/0
2	GTP	F	400	5	-	0/18/38/38	0/3/3/3
3	IMP	F	401	-	-	0/6/26/26	0/3/3/3
4	HDA	F	402	5	-	0/1/6/6	0/0/0/0

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	IMP	C2-N1	2.35	1.38	1.33
3	C	401	IMP	C2-N1	2.38	1.38	1.33
3	F	401	IMP	C2-N1	2.38	1.38	1.33
3	E	401	IMP	C2-N1	2.40	1.38	1.33
3	D	401	IMP	C2-N1	2.43	1.38	1.33

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	IMP	N3-C2-N1	-11.15	120.11	128.87
3	F	401	IMP	N3-C2-N1	-11.15	120.11	128.87
3	E	401	IMP	N3-C2-N1	-10.89	120.32	128.87
3	A	401	IMP	N3-C2-N1	-10.79	120.39	128.87
3	C	401	IMP	N3-C2-N1	-10.79	120.40	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

18 monomers are involved in 68 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	GTP	5	0
3	A	401	IMP	4	0
4	A	402	HDA	7	0
2	B	400	GTP	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	IMP	3	0
4	B	402	HDA	4	0
2	C	400	GTP	5	0
3	C	401	IMP	2	0
4	C	402	HDA	3	0
2	D	400	GTP	5	0
3	D	401	IMP	4	0
4	D	402	HDA	4	0
2	E	400	GTP	5	0
3	E	401	IMP	4	0
4	E	402	HDA	5	0
2	F	400	GTP	5	0
3	F	401	IMP	3	0
4	F	402	HDA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	337/340 (99%)	-0.18	3 (0%) 85 79	25, 41, 74, 98	0
1	B	337/340 (99%)	-0.17	2 (0%) 90 86	25, 41, 71, 97	0
1	C	337/340 (99%)	-0.14	3 (0%) 85 79	26, 48, 78, 101	0
1	D	337/340 (99%)	0.19	16 (4%) 35 24	30, 58, 105, 148	0
1	E	337/340 (99%)	0.19	13 (3%) 43 31	33, 59, 97, 152	0
1	F	337/340 (99%)	-0.05	1 (0%) 94 92	31, 47, 79, 117	0
All	All	2022/2040 (99%)	-0.03	38 (1%) 70 59	25, 49, 87, 152	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	122	GLY	4.5
1	E	305	ALA	4.1
1	D	305	ALA	3.6
1	A	338	THR	3.5
1	E	265	PHE	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	IMP	C	401	23/23	0.82	0.30	2.51	61,87,101,112	0
4	HDA	B	402	8/8	0.80	0.23	2.00	71,79,104,104	0
3	IMP	A	401	23/23	0.82	0.28	1.63	66,113,121,130	0
4	HDA	F	402	8/8	0.89	0.20	1.54	50,62,75,76	0
3	IMP	B	401	23/23	0.88	0.22	1.45	39,91,103,105	0
3	IMP	D	401	23/23	0.84	0.28	1.31	82,112,121,123	0
4	HDA	A	402	8/8	0.86	0.23	1.27	59,73,91,92	0
3	IMP	E	401	23/23	0.91	0.24	0.89	66,95,98,102	0
4	HDA	C	402	8/8	0.88	0.19	0.69	53,75,99,99	0
2	GTP	B	400	32/32	0.90	0.20	0.65	30,45,137,166	0
3	IMP	F	401	23/23	0.93	0.19	0.47	32,71,82,92	0
2	GTP	F	400	32/32	0.92	0.21	0.38	54,65,146,154	0
4	HDA	D	402	8/8	0.92	0.20	0.35	100,102,106,114	0
2	GTP	C	400	32/32	0.94	0.17	0.21	38,54,141,152	0
2	GTP	E	400	32/32	0.89	0.20	0.06	56,91,170,190	0
2	GTP	A	400	32/32	0.92	0.19	0.05	34,49,181,199	0
4	HDA	E	402	8/8	0.89	0.18	-0.00	70,83,102,108	0
2	GTP	D	400	32/32	0.90	0.21	-0.03	46,73,155,158	0
5	MG	F	403	1/1	0.87	0.17	-0.98	77,77,77,77	0
5	MG	C	403	1/1	0.92	0.07	-2.29	72,72,72,72	0
5	MG	B	403	1/1	0.83	0.12	-2.55	45,45,45,45	0
5	MG	A	403	1/1	0.89	0.07	-2.56	49,49,49,49	0
5	MG	E	403	1/1	0.86	0.11	-2.57	69,69,69,69	0
5	MG	D	403	1/1	0.90	0.08	-4.03	53,53,53,53	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.