



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2016 – 10:44 PM EDT

PDB ID : 5K9G
Title : Crystal Structure of GTP Cyclohydrolase-IB with Tris
Authors : Alvarez, J.; Stec, B.; Swairjo, M.A.
Deposited on : 2016-05-31
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>5%</div> </div> </div>
1	B	257	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>8%</div> <div>5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	302	-	-	-	X
3	EDO	B	304	-	-	-	X
3	EDO	B	306	-	-	-	X
3	EDO	B	308	-	-	-	X
5	FMT	B	309	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GTP cyclohydrolase FolE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	14	0
			2007	1276	335	384	12			
1	B	243	Total	C	N	O	S	0	9	0
			1971	1249	332	377	13			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	1
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	177	Total	O	0	4
			180	180		
6	B	206	Total	O	0	1
			207	207		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GTP cyclohydrolase FolE2



- Molecule 1: GTP cyclohydrolase FolE2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.71Å 100.56Å 114.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.16 – 1.90 43.73 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (68.16-1.90) 99.1 (43.73-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.51 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.173 , 0.221 0.174 , 0.221	Depositor DCC
R_{free} test set	2068 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4410	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, ZN, FMT, SNC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/2077	0.98	4/2809 (0.1%)
1	B	1.05	1/2018 (0.0%)	1.01	6/2727 (0.2%)
All	All	1.01	1/4095 (0.0%)	1.00	10/5536 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	193	TYR	CE1-CZ	5.36	1.45	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	LEU	CA-CB-CG	-9.28	93.95	115.30
1	B	204	VAL	CG1-CB-CG2	7.54	122.97	110.90
1	B	90	LEU	CA-CB-CG	-7.44	98.19	115.30
1	B	62	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	B	62	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	27	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	B	204	VAL	CA-CB-CG1	5.54	119.21	110.90
1	B	90	LEU	CB-CG-CD1	5.43	120.22	111.00
1	A	229	ASP	CB-CG-OD2	-5.41	113.44	118.30
1	A	81	ARG	NE-CZ-NH2	5.36	122.98	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2007	0	2027	21	0
1	B	1971	0	1981	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	12	1	0
3	B	24	0	36	2	0
4	B	8	0	11	0	0
5	B	3	0	1	0	0
6	A	180	0	0	5	0
6	B	207	0	0	6	0
All	All	4410	0	4068	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:SER:HB3	1:A:170[B]:THR:HG22	1.47	0.97
1:A:140[B]:MET:HE1	6:A:440:HOH:O	1.65	0.96
1:A:76[B]:PHE:HE2	1:A:135:HIS:CE1	1.88	0.91
1:A:76[B]:PHE:CE2	1:A:135:HIS:CE1	2.69	0.81
1:A:170[B]:THR:HG21	6:A:436:HOH:O	1.94	0.65
1:A:136:SER:CB	1:A:170[B]:THR:HG22	2.29	0.56
1:B:54[B]:GLU:CD	1:B:54[B]:GLU:H	2.09	0.56
1:B:197:LYS:HE2	6:B:546:HOH:O	2.04	0.56
3:B:304:EDO:H21	6:B:589:HOH:O	2.07	0.55
1:A:140[B]:MET:HE3	6:A:453:HOH:O	2.07	0.55
1:A:141:ILE:HD13	1:A:184:VAL:HG21	1.89	0.55
1:A:15:ASN:HA	6:A:431:HOH:O	2.08	0.53
1:A:163:SER:HB3	1:A:241:ASN:HA	1.92	0.52
1:B:33:LYS:HB2	1:B:73:VAL:HG11	1.92	0.51
1:A:19:ASN:H	1:A:19:ASN:HD22	1.60	0.50
1:B:198[B]:ARG:NH2	6:B:401:HOH:O	2.28	0.49
1:A:106:LYS:HG2	1:A:115:ARG:NH2	2.29	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:128[B]:MET:HE1	6:B:518:HOH:O	2.16	0.45
1:A:136:SER:HB3	1:A:170[B]:THR:CG2	2.33	0.44
1:A:163:SER:HB3	1:A:241:ASN:ND2	2.33	0.44
1:A:214:PHE:HB3	1:A:216:GLU:OE1	2.18	0.44
1:A:135:HIS:CD2	3:A:302:EDO:H21	2.54	0.43
1:A:233:LYS:HG2	6:A:546:HOH:O	2.20	0.41
1:B:191:GLN:HG2	6:B:553:HOH:O	2.20	0.41
1:A:19:ASN:ND2	1:B:255:ALA:H	2.19	0.41
1:B:170[B]:THR:OG1	1:B:234:SER:OG	2.37	0.41
1:A:168[B]:SER:OG	1:A:236:VAL:HB	2.21	0.41
1:A:163:SER:HB3	1:A:241:ASN:HD22	1.85	0.41
1:B:197:LYS:HD3	1:B:197:LYS:HA	1.82	0.41
1:B:70[B]:HIS:CB	6:B:477[B]:HOH:O	2.68	0.41
1:B:235:PHE:HA	3:B:304:EDO:H12	2.02	0.40
1:B:106:LYS:HD2	1:B:115:ARG:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/257 (100%)	255 (100%)	1 (0%)	0	100	100
1	B	249/257 (97%)	248 (100%)	1 (0%)	0	100	100
All	All	505/514 (98%)	503 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/224 (102%)	220 (96%)	8 (4%)	43	31
1	B	221/224 (99%)	214 (97%)	7 (3%)	46	35
All	All	449/448 (100%)	434 (97%)	15 (3%)	47	34

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LEU
1	A	90	LEU
1	A	99	VAL
1	A	106	LYS
1	A	156	TYR
1	A	163	SER
1	A	168[A]	SER
1	A	168[B]	SER
1	B	45	LEU
1	B	90	LEU
1	B	135	HIS
1	B	156	TYR
1	B	204	VAL
1	B	230[A]	LYS
1	B	230[B]	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN
1	A	55	GLN
1	A	135	HIS
1	A	241	ASN
1	B	211	ASN
1	B	241	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SNC	A	149	1	2,7,8	0.96	0	1,7,9	2.09	1 (100%)
1	SNC	B	149	1	2,7,8	0.72	0	1,7,9	2.22	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SNC	A	149	1	-	0/0/6/8	0/0/0/0
1	SNC	B	149	1	-	0/0/6/8	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	149	SNC	O-C-CA	-2.22	119.78	125.72
1	A	149	SNC	O-C-CA	-2.09	120.12	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	302	-	3,3,3	0.40	0	2,2,2	0.89	0
3	EDO	A	303[A]	-	3,3,3	0.57	0	2,2,2	0.55	0
4	TRS	B	302	2	7,7,7	0.83	0	9,9,9	1.30	1 (11%)
3	EDO	B	303	-	3,3,3	0.43	0	2,2,2	0.59	0
3	EDO	B	304	-	3,3,3	0.32	0	2,2,2	0.59	0
3	EDO	B	305	-	3,3,3	0.77	0	2,2,2	0.19	0
3	EDO	B	306	-	3,3,3	0.38	0	2,2,2	0.13	0
3	EDO	B	307	-	3,3,3	0.39	0	2,2,2	0.87	0
3	EDO	B	308	-	3,3,3	0.71	0	2,2,2	0.34	0
5	FMT	B	309	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	302	-	-	0/1/1/1	0/0/0/0
3	EDO	A	303[A]	-	-	0/1/1/1	0/0/0/0
4	TRS	B	302	2	-	0/9/9/9	0/0/0/0
3	EDO	B	303	-	-	0/1/1/1	0/0/0/0
3	EDO	B	304	-	-	0/1/1/1	0/0/0/0
3	EDO	B	305	-	-	0/1/1/1	0/0/0/0
3	EDO	B	306	-	-	0/1/1/1	0/0/0/0
3	EDO	B	307	-	-	0/1/1/1	0/0/0/0
3	EDO	B	308	-	-	0/1/1/1	0/0/0/0
5	FMT	B	309	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	302	TRS	C3-C-C2	-2.38	105.64	110.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	EDO	1	0
3	B	304	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/257 (94%)	0.21	13 (5%) 30 33	27, 41, 63, 85	0
1	B	242/257 (94%)	-0.17	2 (0%) 87 88	26, 35, 51, 75	0
All	All	486/514 (94%)	0.02	15 (3%) 52 56	26, 37, 61, 85	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	THR	4.7
1	A	156	TYR	4.4
1	A	155	GLN	4.4
1	A	13	LEU	4.2
1	A	153	ILE	3.9
1	A	198	ARG	3.9
1	A	158	ALA	2.9
1	A	114	ILE	2.5
1	A	151	LYS	2.4
1	B	54[A]	GLU	2.4
1	A	204	VAL	2.3
1	A	111	VAL	2.2
1	A	209	TYR	2.2
1	A	199	PRO	2.0
1	B	82[A]	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	SNC	A	149	8/9	0.88	0.26	-	58,61,79,84	0
1	SNC	B	149	8/9	0.97	0.06	-	32,37,45,56	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	B	306	4/4	0.94	0.21	10.14	51,54,57,58	0
3	EDO	B	308	4/4	0.93	0.26	9.32	39,47,48,48	0
5	FMT	B	309	3/3	0.82	0.21	3.45	58,58,62,74	0
3	EDO	A	302	4/4	0.95	0.17	2.85	40,45,47,54	0
3	EDO	B	304	4/4	0.95	0.14	2.37	47,47,50,55	0
4	TRS	B	302	8/8	0.94	0.12	1.27	36,46,50,61	0
3	EDO	A	303[A]	4/4	0.92	0.13	0.66	50,50,51,53	4
3	EDO	B	305	4/4	0.88	0.11	0.18	44,51,52,56	0
3	EDO	B	303	4/4	0.97	0.08	-0.26	38,41,43,44	0
2	ZN	B	301	1/1	1.00	0.05	-2.03	35,35,35,35	0
2	ZN	A	301	1/1	0.99	0.08	-3.35	56,56,56,56	0
3	EDO	B	307	4/4	0.92	0.19	-	50,57,62,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.