



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:02 PM GMT

PDB ID : 4KAY
Title : Structure of the soluble domain of Lipooligosaccharide phosphoethanolamine transferase A from *Neisseria meningitidis* - complex with Zn
Authors : Vrieling, A.; Wanty, C.; Anandan, A.
Deposited on : 2013-04-23
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6655 (1.80-1.76)
Clashscore	102246	7658 (1.80-1.76)
Ramachandran outliers	100387	7570 (1.80-1.76)
Sidechain outliers	100360	7569 (1.80-1.76)
RSRZ outliers	91569	6671 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 90% 9% . </div> </div>
1	B	335	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 93% 6% .. </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11984 atoms, of which 5344 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YhbX/YhjW/YijP/YjdB family protein.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
1	A	333	Total	C	H	N	O	P	S				
			5465	1735	2693	484	533	1	19		0	22	0
1	B	333	Total	C	H	N	O	P	S				
			5387	1709	2651	476	530	1	20		0	16	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Zn		
			3	3	0	0
2	A	3	Total	Zn		
			3	3	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	555	Total	O	0	0
			555	555		
4	B	561	Total	O	0	0
			561	561		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: YhbX/YhjW/YijP/YjdB family protein



- Molecule 1: YhbX/YhjW/YijP/YjdB family protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.07Å 90.52Å 91.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.80 – 1.78 19.80 – 1.78	Depositor EDS
% Data completeness (in resolution range)	97.8 (19.80-1.78) 97.9 (19.80-1.78)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 1.78Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.149 , 0.190 0.162 , 0.202	Depositor DCC
R_{free} test set	3369 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	1.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.9	EDS
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.56$, $\langle L^2 \rangle = 0.41$	Xtriage
Outliers	4 of 66464 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11984	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.06 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.8725e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	1/2882 (0.0%)	0.78	6/3905 (0.2%)
1	B	0.74	5/2813 (0.2%)	0.77	2/3812 (0.1%)
All	All	0.71	6/5695 (0.1%)	0.77	8/7717 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	490[A]	ARG	CA-C	-5.52	1.38	1.52
1	B	490[B]	ARG	CA-C	-5.52	1.38	1.52
1	B	439	ASN	CG-ND2	-5.50	1.19	1.32
1	A	216	GLN	CD-OE1	-5.34	1.12	1.24
1	B	439	ASN	CG-OD1	-5.24	1.12	1.24
1	B	463	TYR	C-O	-5.09	1.13	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	490[A]	ARG	NE-CZ-NH2	6.25	123.43	120.30
1	A	490[B]	ARG	NE-CZ-NH2	6.25	123.43	120.30
1	B	232	ARG	NE-CZ-NH1	5.97	123.29	120.30
1	B	489	PHE	C-N-CA	-5.90	106.94	121.70
1	A	412[B]	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	412[C]	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	412[D]	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	232	ARG	NE-CZ-NH1	5.39	122.99	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2772	2693	2716	22	6
1	B	2736	2651	2650	12	5
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	1	0
4	A	555	0	0	3	1
4	B	561	0	0	2	2
All	All	6640	5344	5366	35	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:435:ARG:NH2	4:A:1150:HOH:O	2.26	0.67
1:A:411:THR:OG1	1:A:414[A]:THR:HG23	1.94	0.67
1:A:297:ASP:OD2	1:A:300:LYS:NZ	2.32	0.57
1:A:212:ILE:HG13	1:A:213:PRO:HD2	1.86	0.55
1:B:212:ILE:HG13	1:B:213:PRO:HD2	1.89	0.55
1:A:272[B]:GLN:HB3	1:A:508[B]:GLU:HG2	1.91	0.53
1:B:263:ARG:O	1:B:266[A]:GLU:HB2	2.08	0.52
1:A:445[B]:SER:OG	1:A:486:SER:HB3	2.10	0.52
1:A:487:LYS:HE2	4:A:1246:HOH:O	2.10	0.49
1:A:412[C]:ARG:CG	1:A:457:LEU:O	2.61	0.49
1:A:412[C]:ARG:HG3	1:A:457:LEU:O	2.12	0.48
1:A:506:GLU:HB2	4:A:950:HOH:O	2.12	0.48
1:A:212:ILE:HG13	1:A:213:PRO:CD	2.44	0.48
1:B:445[B]:SER:OG	1:B:486:SER:HB3	2.13	0.48
1:B:411:THR:OG1	1:B:414[A]:THR:HG23	2.14	0.48
1:A:401:THR:C	1:A:414[B]:THR:CG2	2.84	0.46
1:A:383:HIS:O	1:A:390:ARG:NH2	2.49	0.46
1:A:445[B]:SER:HG	1:A:484:TRP:HZ3	1.64	0.45
1:B:401:THR:C	1:B:414[B]:THR:CG2	2.84	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:352:GLU:OE1	1:B:390:ARG:NH1	2.49	0.45
1:B:412:ARG:HG3	3:B:604:PO4:O2	2.16	0.45
1:B:212:ILE:HG13	1:B:213:PRO:CD	2.49	0.43
1:B:295:ASP:OD1	4:B:1102:HOH:O	2.21	0.43
1:A:494:GLY:HA2	1:A:542[A]:ARG:NH1	2.34	0.42
1:B:487:LYS:HE3	4:B:1204:HOH:O	2.19	0.42
1:A:271:PRO:O	1:A:272[B]:GLN:OE1	2.37	0.42
1:A:496:ASP:HB2	1:A:542[B]:ARG:CZ	2.50	0.42
1:A:317[B]:GLU:HG2	1:A:371:LYS:HD3	2.02	0.41
1:A:233:PHE:O	1:A:445[A]:SER:HA	2.20	0.41
1:A:233:PHE:O	1:A:445[B]:SER:HA	2.20	0.40
1:B:233:PHE:O	1:B:445[A]:SER:HA	2.21	0.40
1:A:352:GLU:OE1	1:A:390:ARG:NH1	2.53	0.40
1:B:408:ASN:OD1	1:B:408:ASN:N	2.47	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:ARG:HH12	1:B:490[B]:ARG:HH22[2_455]	1.18	0.42
1:A:266[B]:GLU:OE2	4:B:1018:HOH:O[2_555]	1.78	0.42
1:A:349:ARG:HH22	1:B:490[B]:ARG:HH21[2_455]	1.21	0.39
1:A:349:ARG:NH1	1:B:490[B]:ARG:NH2[2_455]	2.05	0.15
1:A:349:ARG:NH1	1:B:490[B]:ARG:HH22[2_455]	1.49	0.11
4:A:1128:HOH:O	4:B:1018:HOH:O[2_555]	2.12	0.08
1:A:349:ARG:NH2	1:B:490[B]:ARG:HH21[2_455]	1.58	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	352/335 (105%)	344 (98%)	8 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	345/335 (103%)	338 (98%)	7 (2%)	0	100	100
All	All	697/670 (104%)	682 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/293 (107%)	308 (98%)	5 (2%)	70	57
1	B	306/293 (104%)	303 (99%)	3 (1%)	82	75
All	All	619/586 (106%)	611 (99%)	8 (1%)	82	66

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	212	ILE
1	A	284	LEU
1	A	412[B]	ARG
1	A	412[C]	ARG
1	A	412[D]	ARG
1	B	212	ILE
1	B	284	LEU
1	B	408	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	280	1,2	8,10,11	1.28	1 (12%)	7,14,16	1.49	2 (28%)
1	TPO	B	280	1,2	8,10,11	1.22	1 (12%)	7,14,16	1.62	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	280	1,2	-	0/8/11/13	0/0/0/0
1	TPO	B	280	1,2	-	0/8/11/13	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	280	TPO	P-O3P	2.33	1.63	1.54
1	B	280	TPO	P-O3P	2.44	1.63	1.54

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	280	TPO	O-C-CA	-2.19	119.65	125.44
1	A	280	TPO	O-C-CA	-2.09	119.93	125.44
1	B	280	TPO	OG1-P-O1P	3.04	114.70	107.11
1	A	280	TPO	OG1-P-O1P	3.08	114.80	107.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	604	-	4,4,4	0.43	0	6,6,6	0.29	0
3	PO4	B	604	-	4,4,4	0.47	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	604	-	-	0/0/0/0	0/0/0/0
3	PO4	B	604	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	PO4	1	0
3	B	604	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/335 (99%)	-0.44	4 (1%) 81 81	6, 11, 26, 43	1 (0%)
1	B	332/335 (99%)	-0.44	5 (1%) 76 76	7, 12, 27, 45	3 (0%)
All	All	664/670 (99%)	-0.44	9 (1%) 78 77	6, 12, 26, 45	4 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	412[B]	ARG	3.7
1	B	212	ILE	3.0
1	B	543	PRO	2.7
1	B	211	ASN	2.6
1	A	543	PRO	2.5
1	B	213	PRO	2.3
1	A	212	ILE	2.2
1	B	412	ARG	2.2
1	A	211	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	280	11/12	0.95	0.14	-	10,14,47,48	0
1	TPO	B	280	11/12	0.94	0.13	-	9,14,50,54	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PO4	A	604	5/5	0.95	0.19	-0.65	41,45,55,65	0
2	ZN	B	602	1/1	0.96	0.05	-1.09	23,23,23,23	1
2	ZN	A	602	1/1	0.98	0.04	-1.28	21,21,21,21	1
2	ZN	B	601	1/1	0.99	0.04	-1.60	10,10,10,10	0
2	ZN	A	601	1/1	1.00	0.03	-2.66	11,11,11,11	0
2	ZN	B	603	1/1	0.94	0.06	-	51,51,51,51	0
3	PO4	B	604	5/5	0.94	0.17	-	40,43,60,65	0
2	ZN	A	603	1/1	0.91	0.05	-	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.