



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 11:45 PM BST

PDB ID : 2KIL
Title : NMR structure of the H103G mutant SO2144 H-NOX domain from *Shewanella oneidensis* in the Fe(II)CO ligation state
Authors : Erbil, W.K.
Deposited on : 2009-05-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 76%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	181	<div> <div>49%</div> <div>31%</div> <div>•</div> <div>19%</div> </div>

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:4-A:29, A:45-A:108, A:117-A:132, A:138-A:178 (147)	0.24	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 7, 8, 9, 11, 13, 14, 15, 17, 18, 19, 20
2	3, 6, 10
3	12, 16


- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms						Trace
1	A	181	Total	C	H	N	O	S	0
			2856	910	1423	247	264	12	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	103	GLY	HIS	ENGINEERED	UNP Q8EF49

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					
2	A	1	Total	C	Fe	H	N	O
			73	34	1	30	4	4

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms		
3	A	1	Total	C	O
			2	1	1

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).




Mol	Chain	Residues	Atoms			
4	A	1	Total	C	H	N
			9	3	4	2

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Putative uncharacterized protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

- Molecule 1: Putative uncharacterized protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16278
Number of chemical shift lists	1
Total number of shifts	1818
Number of shifts mapped to atoms	1804
Number of unparsed shifts	0
Number of shifts with mapping errors	14
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	76%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM, IMD

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1160	1154	1146	54±7
2	A	43	30	30	2±2
4	A	5	4	4	0±0
All	All	24200	23760	23600	1086

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

5 of 342 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:25:LEU:HD13	1:A:26:LEU:N	0.89	1.82	18	20
1:A:122:LEU:N	1:A:122:LEU:HD22	0.76	1.96	9	6
1:A:25:LEU:O	1:A:25:LEU:HD22	0.69	1.87	2	13
1:A:25:LEU:C	1:A:25:LEU:HD22	0.69	2.08	12	9
1:A:157:ILE:O	1:A:157:ILE:HD13	0.68	1.88	17	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/181 (81%)	130±2 (89±2%)	13±3 (9±2%)	4±1 (2±1%)	11	48
All	All	2940/3620 (81%)	2605 (89%)	262 (9%)	73 (2%)	11	48

5 of 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	98	ILE	20
1	A	100	ASP	20
1	A	165	MET	11
1	A	86	ASP	4
1	A	4	ILE	4

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/159 (81%)	121±2 (93±1%)	8±2 (7±1%)	25	71
All	All	2580/3180 (81%)	2411 (93%)	169 (7%)	25	71

5 of 38 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	47	LEU	20
1	A	25	LEU	20
1	A	51	VAL	17
1	A	21	VAL	12
1	A	163	THR	12

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	HEM	A	182	3,4	24,50,50	1.96±0.03	0±0 (0±0%)
3	CMO	A	183	2	0,1,1	0.00±0.00	-
4	IMD	A	184	2	4,5,5	0.48±0.01	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	HEM	A	182	3,4	16,82,82	1.31±0.02	0±0 (0±0%)
3	CMO	A	183	2	0,0,0	0.00±0.00	-
4	IMD	A	184	2	4,5,5	0.76±0.02	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	182	3,4	-	0±0,6,54,54	0±0,0,8,8
3	CMO	A	183	2	-	0±0,0,0,0	0±0,0,0,0
4	IMD	A	184	2	-	0±0,0,0,0	0±0,1,1,1

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	A	182	HEM	C2D-C3D-CAD-CBD	1

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 76% for the well-defined parts and 71% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 16278

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1818
Number of shifts mapped to atoms	1804
Number of unparsed shifts	0
Number of shifts with mapping errors	14
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	20

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 14) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	HEM	HMC1	3.261	0.03	1
UNMAPPED	1	HEM	HMD1	2.659	0.03	1
UNMAPPED	1	IMD	HD1	10.596	0.03	1
UNMAPPED	1	HEM	HHB	9.493	0.03	1
UNMAPPED	1	HEM	HHD	9.569	0.03	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	171	-0.32 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	159	0.55 ± 0.16	Should be applied
$^{13}\text{C}'$	0	—	—

Continued on next page...

Continued from previous page...

Nucleus	# values	Correction \pm precision, ppm	Suggested action
¹⁵ N	166	0.10 \pm 0.26	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 76%, i.e. 1376 atoms were assigned a chemical shift out of a possible 1811. 0 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	578/731 (79%)	289/292 (99%)	147/294 (50%)	142/145 (98%)
Sidechain	758/906 (84%)	462/524 (88%)	296/349 (85%)	0/33 (0%)
Aromatic	40/174 (23%)	39/92 (42%)	0/65 (0%)	1/17 (6%)
Overall	1376/1811 (76%)	790/908 (87%)	443/708 (63%)	143/195 (73%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	77	LEU	HD11	-2.25	2.16 – -0.64	-10.8
1	A	77	LEU	HD13	-2.25	2.16 – -0.64	-10.8
1	A	77	LEU	HD12	-2.25	2.16 – -0.64	-10.8
1	A	115	LEU	HD23	-2.04	2.14 – -0.66	-9.9
1	A	115	LEU	HD21	-2.04	2.14 – -0.66	-9.9
1	A	115	LEU	HD22	-2.04	2.14 – -0.66	-9.9
1	A	145	LEU	HD13	-2.02	2.16 – -0.64	-9.9
1	A	145	LEU	HD12	-2.02	2.16 – -0.64	-9.9
1	A	145	LEU	HD11	-2.02	2.16 – -0.64	-9.9
1	A	145	LEU	HD21	-1.37	2.14 – -0.66	-7.5
1	A	145	LEU	HD23	-1.37	2.14 – -0.66	-7.5
1	A	145	LEU	HD22	-1.37	2.14 – -0.66	-7.5
1	A	73	LEU	HD21	-1.24	2.14 – -0.66	-7.1
1	A	73	LEU	HD23	-1.24	2.14 – -0.66	-7.1
1	A	73	LEU	HD22	-1.24	2.14 – -0.66	-7.1
1	A	98	ILE	HG21	-0.97	2.13 – -0.57	-6.5
1	A	98	ILE	HG22	-0.97	2.13 – -0.57	-6.5
1	A	98	ILE	HG23	-0.97	2.13 – -0.57	-6.5
1	A	73	LEU	HG	-0.55	3.16 – -0.14	-6.2
1	A	10	GLU	HA	1.99	6.30 – 2.20	-5.5

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

