



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:06 PM GMT

PDB ID : 4KI0
Title : Crystal structure of the maltose-binding protein/maltose transporter complex
in an outward-facing conformation bound to maltohexaose
Authors : Oldham, M.L.; Chen, S.; Chen, J.
Deposited on : 2013-05-01
Resolution : 2.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	<div> <div>6%</div> <div>87% 10% •</div> </div>
1	B	381	<div> <div>11%</div> <div>86% 11% •</div> </div>
2	E	380	<div> <div>7%</div> <div>94% • •</div> </div>
3	F	514	<div> <div>9%</div> <div>85% 9% • 5%</div> </div>
4	G	296	<div> <div>6%</div> <div>88% 9% • •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	UMQ	F	607	-	-	-	X
7	UMQ	F	608	-	-	-	X
7	UMQ	F	609	-	-	-	X
7	UMQ	G	303	-	-	-	X
7	UMQ	G	306	-	-	-	X
7	UMQ	G	307	-	-	-	X
8	GLC	F	602	X	-	-	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 15618 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	371	Total	C	N	O	S	0	1	0
			2884	1824	518	529	13			
1	B	371	Total	C	N	O	S	0	0	0
			2876	1819	515	529	13			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	372	ALA	-	EXPRESSION TAG	UNP C9QV42
A	373	SER	-	EXPRESSION TAG	UNP C9QV42
A	374	ALA	-	EXPRESSION TAG	UNP C9QV42
A	375	SER	-	EXPRESSION TAG	UNP C9QV42
A	376	HIS	-	EXPRESSION TAG	UNP C9QV42
A	377	HIS	-	EXPRESSION TAG	UNP C9QV42
A	378	HIS	-	EXPRESSION TAG	UNP C9QV42
A	379	HIS	-	EXPRESSION TAG	UNP C9QV42
A	380	HIS	-	EXPRESSION TAG	UNP C9QV42
A	381	HIS	-	EXPRESSION TAG	UNP C9QV42
B	372	ALA	-	EXPRESSION TAG	UNP C9QV42
B	373	SER	-	EXPRESSION TAG	UNP C9QV42
B	374	ALA	-	EXPRESSION TAG	UNP C9QV42
B	375	SER	-	EXPRESSION TAG	UNP C9QV42
B	376	HIS	-	EXPRESSION TAG	UNP C9QV42
B	377	HIS	-	EXPRESSION TAG	UNP C9QV42
B	378	HIS	-	EXPRESSION TAG	UNP C9QV42
B	379	HIS	-	EXPRESSION TAG	UNP C9QV42
B	380	HIS	-	EXPRESSION TAG	UNP C9QV42
B	381	HIS	-	EXPRESSION TAG	UNP C9QV42

- Molecule 2 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	370	Total	C	N	O	S	0	0	0
			2877	1853	469	549	6			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	371	ALA	-	EXPRESSION TAG	UNP P0AEX9
E	372	SER	-	EXPRESSION TAG	UNP P0AEX9
E	373	ALA	-	EXPRESSION TAG	UNP P0AEX9
E	374	SER	-	EXPRESSION TAG	UNP P0AEX9
E	375	HIS	-	EXPRESSION TAG	UNP P0AEX9
E	376	HIS	-	EXPRESSION TAG	UNP P0AEX9
E	377	HIS	-	EXPRESSION TAG	UNP P0AEX9
E	378	HIS	-	EXPRESSION TAG	UNP P0AEX9
E	379	HIS	-	EXPRESSION TAG	UNP P0AEX9
E	380	HIS	-	EXPRESSION TAG	UNP P0AEX9

- Molecule 3 is a protein called Maltose transport system permease protein MalF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	490	Total	C	N	O	S	0	1	0
			3819	2512	607	683	17			


- Molecule 4 is a protein called Binding-protein-dependent transport systems inner membrane component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	289	Total	C	N	O	S	0	1	0
			2238	1500	357	373	8			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
6	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 7 is UNDECYL-MALTOSIDE (three-letter code: UMQ) (formula: $C_{23}H_{44}O_{11}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	E	1	Total	C	O	0	0
			34	23	11		
7	F	1	Total	C	O	0	0
			13	12	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			13	12	1		
7	F	1	Total	C	O	0	0
			13	12	1		
7	G	1	Total	C	O	0	0
			34	23	11		
7	G	1	Total	C	O	0	0
			13	12	1		
7	G	1	Total	C	O	0	0
			13	12	1		
7	G	1	Total	C	O	0	0
			13	12	1		
7	G	1	Total	C	O	0	0
			23	17	6		
7	G	1	Total	C	O	0	0
			13	12	1		
7	G	1	Total	C	O	0	0
			34	23	11		

- Molecule 8 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	F	4	Total	C	O	0	0
			34	18	16		

- Molecule 9 is (1R)-2-{{[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PALMITOYLOXY)METHYL]ETHYL (11E)-OCTADEC-11-ENOATE (three-letter code: PGV) (formula: C₄₀H₇₇O₁₀P).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	F	1	Total	C	O	P	0	0
			51	40	10	1		
9	F	1	Total	C	O	P	0	0
			51	40	10	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	112	Total	O	0	0
			112	112		
10	B	123	Total	O	0	0
			123	123		
10	E	105	Total	O	0	0
			105	105		
10	F	87	Total	O	0	0
			87	87		
10	G	81	Total	O	0	0
			81	81		

- Molecule 1: ABC transporter related protein





- Molecule 4: Binding-protein-dependent transport systems inner membrane component



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	81.59 Å 96.63 Å 112.32 Å 85.30° 79.11° 73.00°	Depositor
Resolution (Å)	19.82 – 2.38 19.81 – 2.38	Depositor EDS
% Data completeness (in resolution range)	87.9 (19.82-2.38) 87.6 (19.81-2.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.38 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.192 , 0.228 0.196 , 0.228	Depositor DCC
R_{free} test set	6156 reflections (5.70%)	DCC
Wilson B-factor (Å ²)	58.8	Xtriage
Anisotropy	0.812	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 64.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 113613 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15618	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGV, ANP, MG, GLC, UMQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.31	0/2937	0.55	0/3982
1	B	0.31	0/2926	0.54	0/3968
2	E	0.32	0/2946	0.48	0/3998
3	F	0.32	0/3917	0.51	0/5333
4	G	0.33	0/2304	0.52	0/3150
All	All	0.32	0/15030	0.52	0/20431

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
8	F	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
8	F	602	GLC	C1

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2884	0	2954	22	0
1	B	2876	0	2942	19	0
2	E	2877	0	2859	6	0
3	F	3819	0	3853	29	0
4	G	2238	0	2323	20	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	31	0	13	0	0
6	B	31	0	13	1	0
7	E	34	0	44	0	0
7	F	39	0	69	0	0
7	G	143	0	213	1	0
8	F	34	0	28	0	0
9	F	102	0	152	0	0
10	A	112	0	0	6	0
10	B	123	0	0	3	0
10	E	105	0	0	1	0
10	F	87	0	0	1	0
10	G	81	0	0	2	0
All	All	15618	0	15463	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:429:LEU:HD23	4:G:172:LEU:HD12	1.53	0.88
1:A:173:ARG:CZ	10:A:1690:HOH:O	2.36	0.73
4:G:295:LYS:NZ	10:G:410:HOH:O	2.21	0.73
1:A:223:HIS:CE1	1:A:368:GLU:HG2	2.27	0.68
4:G:212:VAL:HG22	4:G:284[A]:TRP:CD2	2.30	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/381 (97%)	363 (98%)	7 (2%)	0	100	100
1	B	369/381 (97%)	349 (95%)	18 (5%)	2 (0%)	34	46
2	E	368/380 (97%)	360 (98%)	7 (2%)	1 (0%)	46	61
3	F	487/514 (95%)	469 (96%)	15 (3%)	3 (1%)	30	40
4	G	288/296 (97%)	283 (98%)	4 (1%)	1 (0%)	46	61
All	All	1882/1952 (96%)	1824 (97%)	51 (3%)	7 (0%)	39	53

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	37	GLN
2	E	173	ASN
3	F	38	GLY
1	B	38	SER
3	F	36	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/323 (98%)	305 (97%)	10 (3%)	46	66
1	B	314/323 (97%)	303 (96%)	11 (4%)	43	62
2	E	297/305 (97%)	296 (100%)	1 (0%)	94	98
3	F	402/424 (95%)	385 (96%)	17 (4%)	36	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	G	232/237 (98%)	222 (96%)	10 (4%)	35 52
All	All	1560/1612 (97%)	1511 (97%)	49 (3%)	47 67

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	20	LEU
3	F	59	ASN
4	G	137	LEU
3	F	37	GLN
3	F	112	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	37	GLN
3	F	106	GLN
3	F	437	ASN
2	E	218	ASN
3	F	376	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

Of 4 carbohydrates modelled in this entry, 1 is modelled with single atom - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	GLC	F	602	8	11,11,12	0.64	0	14,15,17	1.67	5 (35%)
8	GLC	F	603	8	11,11,12	0.64	0	14,15,17	1.09	1 (7%)
8	GLC	F	604	8	11,11,12	0.61	0	14,15,17	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GLC	F	602	8	1/1/4/5	0/2/19/22	0/1/1/1
8	GLC	F	603	8	-	0/2/19/22	0/1/1/1
8	GLC	F	604	8	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	F	602	GLC	O4-C4-C3	-2.77	104.11	110.34
8	F	602	GLC	C2-C3-C4	2.06	114.54	111.04
8	F	602	GLC	C1-C2-C3	2.25	112.21	109.54
8	F	602	GLC	C3-C4-C5	2.49	114.54	110.20
8	F	603	GLC	C1-O5-C5	2.98	116.03	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
8	F	602	GLC	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	ANP	A	1502	5	27,33,33	1.99	6 (22%)	30,52,52	2.15	7 (23%)
6	ANP	B	402	5	27,33,33	2.00	7 (25%)	30,52,52	2.27	6 (20%)
7	UMQ	E	401	-	35,35,35	0.41	0	46,46,46	0.71	0
9	PGV	F	605	-	50,50,50	1.07	3 (6%)	51,56,56	1.05	4 (7%)
9	PGV	F	606	-	50,50,50	1.07	3 (6%)	51,56,56	0.97	3 (5%)
7	UMQ	F	607	-	12,12,35	0.36	0	11,11,46	0.40	0
7	UMQ	F	608	-	12,12,35	0.38	0	11,11,46	0.37	0
7	UMQ	F	609	-	12,12,35	0.35	0	11,11,46	0.42	0
7	UMQ	G	301	-	35,35,35	0.52	1 (2%)	46,46,46	1.05	2 (4%)
7	UMQ	G	302	-	12,12,35	0.38	0	11,11,46	0.43	0
7	UMQ	G	303	-	12,12,35	0.35	0	11,11,46	0.42	0
7	UMQ	G	304	-	12,12,35	0.38	0	11,11,46	0.38	0
7	UMQ	G	305	-	23,23,35	0.48	0	28,28,46	0.68	0
7	UMQ	G	306	-	12,12,35	0.36	0	11,11,46	0.42	0
7	UMQ	G	307	-	35,35,35	0.52	1 (2%)	46,46,46	1.05	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ANP	A	1502	5	-	1/12/38/38	0/3/3/3
6	ANP	B	402	5	-	1/12/38/38	0/3/3/3
7	UMQ	E	401	-	-	0/20/60/60	0/2/2/2
9	PGV	F	605	-	-	0/55/55/55	0/0/0/0
9	PGV	F	606	-	-	0/55/55/55	0/0/0/0
7	UMQ	F	607	-	-	0/10/10/60	0/0/0/2
7	UMQ	F	608	-	-	0/10/10/60	0/0/0/2
7	UMQ	F	609	-	-	0/10/10/60	0/0/0/2
7	UMQ	G	301	-	-	0/20/60/60	0/2/2/2
7	UMQ	G	302	-	-	0/10/10/60	0/0/0/2
7	UMQ	G	303	-	-	0/10/10/60	0/0/0/2
7	UMQ	G	304	-	-	0/10/10/60	0/0/0/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	UMQ	G	305	-	-	0/14/34/60	0/1/1/2
7	UMQ	G	306	-	-	0/10/10/60	0/0/0/2
7	UMQ	G	307	-	-	1/20/60/60	0/2/2/2

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	402	ANP	PG-O3G	-2.08	1.50	1.56
6	B	402	ANP	PG-O2G	-2.06	1.50	1.56
6	A	1502	ANP	PG-O3G	-2.01	1.51	1.56
7	G	307	UMQ	O1'-C1'	2.24	1.44	1.40
7	G	301	UMQ	O1'-C1'	2.36	1.44	1.40

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	402	ANP	N3-C2-N1	-8.25	122.58	128.89
6	A	1502	ANP	N3-C2-N1	-7.86	122.88	128.89
6	B	402	ANP	O1G-PG-N3B	-5.18	103.95	111.90
6	A	1502	ANP	O1G-PG-N3B	-3.97	105.81	111.90
6	B	402	ANP	O1B-PB-N3B	-2.78	107.64	111.90

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	402	ANP	O1B-PB-N3B-PG
6	A	1502	ANP	O1B-PB-N3B-PG
7	G	307	UMQ	CA-O1'-C1'-O5'

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	402	ANP	1	0
7	G	307	UMQ	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/381 (97%)	0.30	21 (5%) 27 31	54, 79, 108, 126	0
1	B	371/381 (97%)	0.57	43 (11%) 6 7	54, 92, 147, 170	0
2	E	370/380 (97%)	0.26	28 (7%) 17 19	59, 83, 111, 132	0
3	F	490/514 (95%)	0.50	45 (9%) 11 13	53, 85, 134, 153	0
4	G	289/296 (97%)	0.30	17 (5%) 26 30	51, 71, 115, 148	0
All	All	1891/1952 (96%)	0.40	154 (8%) 15 17	51, 82, 130, 170	0

The worst 5 of 154 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	372	ALA	9.6
1	B	251	ILE	8.2
1	B	295	ILE	7.1
3	F	37	GLN	6.9
4	G	70	ALA	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
8	GLC	F	602	11/12	0.95	0.11	-0.88	73,80,87,95	0
8	GLC	F	604	11/12	0.99	0.07	-1.63	61,62,66,67	0
8	GLC	F	603	11/12	0.98	0.09	-1.65	61,64,68,68	0
8	GLC	F	601	1/12	0.82	0.26	-	91,91,91,91	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	UMQ	F	609	13/34	0.81	0.34	6.48	95,101,111,111	0
7	UMQ	F	607	13/34	0.81	0.26	5.49	87,90,92,94	0
7	UMQ	G	306	13/34	0.67	0.29	5.00	90,99,118,119	0
7	UMQ	F	608	13/34	0.70	0.29	4.89	97,100,115,115	0
7	UMQ	G	307	34/34	0.74	0.35	3.45	99,135,156,157	0
7	UMQ	G	303	13/34	0.86	0.19	2.52	78,86,99,101	0
7	UMQ	G	301	34/34	0.68	0.33	1.70	91,132,157,158	0
9	PGV	F	605	51/51	0.88	0.23	1.51	89,98,109,111	0
9	PGV	F	606	51/51	0.78	0.25	1.38	86,106,129,134	0
7	UMQ	G	305	23/34	0.78	0.22	1.20	94,106,127,136	0
7	UMQ	E	401	34/34	0.93	0.19	0.98	80,87,96,97	0
6	ANP	A	1502	31/31	0.98	0.11	-0.20	55,62,75,81	0
6	ANP	B	402	31/31	0.96	0.11	-0.46	58,70,84,89	0
5	MG	B	401	1/1	0.96	0.10	-0.79	59,59,59,59	0
5	MG	A	1501	1/1	0.99	0.04	-2.36	57,57,57,57	0
7	UMQ	G	304	13/34	0.81	0.28	-	88,100,110,111	0
7	UMQ	G	302	13/34	0.80	0.21	-	86,93,96,99	0

6.5 Other polymers [i](#)

There are no such residues in this entry.