



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 11:27 PM BST

PDB ID : 2KL5  
Title : Solution NMR Structure of protein yutD from B.subtilis, Northeast Structural Genomics Consortium Target SR232  
Authors : Liu, G.; Hamilton, K.; Xiao, R.; Ciccosanti, C.; Ho, C.J.; Everett, J.; Nair, R.; Acton, T.; Rost, B.; Montelione, G.T.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2009-06-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 81%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	110	<div> <div style="width: 76%; background-color: green;"></div> <div style="width: 21%; background-color: cyan;"></div> <div style="width: 3%; background-color: yellow;"></div> <div style="width: 2%; background-color: orange;"></div> <div style="width: 2%; background-color: red;"></div> </div> <div>76% .. 21%</div>

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:9-A:66, A:75-A:103 (87)	0.53	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 6, 8, 10, 11, 12, 14, 15, 16, 18
2	1, 5, 9, 19
3	7, 13
Single-model clusters	17; 20

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1828 atoms, of which 893 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Uncharacterized protein yutD.

Mol	Chain	Residues	Atoms						Trace
1	A	110	Total	C	H	N	O	S	0
			1828	602	893	159	169	5	

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP O32127
A	2	ILE	-	EXPRESSION TAG	UNP O32127
A	3	GLY	-	EXPRESSION TAG	UNP O32127
A	4	MET	-	EXPRESSION TAG	UNP O32127
A	5	SER	-	EXPRESSION TAG	UNP O32127
A	6	GLU	-	EXPRESSION TAG	UNP O32127
A	7	LYS	-	EXPRESSION TAG	UNP O32127
A	8	ARG	-	EXPRESSION TAG	UNP O32127
A	9	GLY	-	EXPRESSION TAG	UNP O32127
A	10	GLU	-	EXPRESSION TAG	UNP O32127
A	11	ILE	-	EXPRESSION TAG	UNP O32127
A	103	LEU	-	EXPRESSION TAG	UNP O32127
A	104	GLU	-	EXPRESSION TAG	UNP O32127
A	105	HIS	-	EXPRESSION TAG	UNP O32127
A	106	HIS	-	EXPRESSION TAG	UNP O32127
A	107	HIS	-	EXPRESSION TAG	UNP O32127
A	108	HIS	-	EXPRESSION TAG	UNP O32127
A	109	HIS	-	EXPRESSION TAG	UNP O32127
A	110	HIS	-	EXPRESSION TAG	UNP O32127



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, simulated annealing, molecular dynamics, null*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CNS	geometry optimization	
CNS	structure solution	
CYANA	refinement	3.0
CYANA	geometry optimization	3.0
CYANA	structure solution	3.0
AutoStructure	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 16384
Number of chemical shift lists	1
Total number of shifts	1077
Number of shifts mapped to atoms	1077
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	81%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	735	706	705	5±2
All	All	14700	14120	14100	95

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 55 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:88:CYS:SG	1:A:92:CYS:N	0.75	2.59	13	5
1:A:88:CYS:HA	1:A:92:CYS:HB3	0.69	1.64	14	1
1:A:12:MET:SD	1:A:12:MET:N	0.64	2.70	5	8
1:A:12:MET:N	1:A:12:MET:SD	0.60	2.74	4	10
1:A:15:ILE:HG12	1:A:16:GLN:H	0.58	1.58	6	1

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/110 (79%)	78±2 (90±3%)	7±2 (8±2%)	2±1 (2±1%)	15	58
All	All	1740/2200 (79%)	1565 (90%)	144 (8%)	31 (2%)	15	58

5 of 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	54	GLY	7
1	A	91	GLY	4
1	A	103	LEU	3
1	A	10	GLU	3
1	A	56	GLY	2

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/97 (78%)	73±1 (96±2%)	3±1 (4±2%)	46	87
All	All	1520/1940 (78%)	1466 (96%)	54 (4%)	46	87

5 of 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	12	MET	19
1	A	45	LYS	6
1	A	27	LYS	6
1	A	102	LYS	3
1	A	38	ARG	3

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 81% for the well-defined parts and 67% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 16384

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1077
Number of shifts mapped to atoms	1077
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	87	$0.20 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	82	$0.39 \pm 0.19$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	71	$0.64 \pm 0.22$	Should be applied
$^{15}\text{N}$	68	$-0.13 \pm 0.30$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 81%, i.e. 927 atoms were assigned a chemical shift out of a possible 1138. 11 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	368/435 (85%)	149/174 (86%)	152/174 (87%)	67/87 (77%)
Sidechain	443/547 (81%)	274/320 (86%)	166/201 (83%)	3/26 (12%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	116/156 (74%)	77/82 (94%)	38/71 (54%)	1/3 (33%)
Overall	927/1138 (81%)	500/576 (87%)	356/446 (80%)	71/116 (61%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	61	LYS	HE2	1.84	3.87 – 1.97	-5.7
1	A	31	ASN	H	5.09	11.45 – 5.25	-5.3

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

