



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:08 PM GMT

PDB ID : 4KON  
Title : The structure of hemagglutinin from avian-origin H7N9 influenza virus in complex with human receptor analog 6'SLNLN (NeuAc<#945;2-6Gal<#946;1-4GlcNAc<#946;1-3Gal<#946;1-4Glc)  
Authors : Shi, Y.; Zhang, W.; Wang, F.; Qi, J.; Song, H.; Wu, Y.; Gao, F.; Zhang, Y.; Fan, Z.; Gong, W.; Wang, D.; Shu, Y.; Wang, Y.; Yan, J.; Gao, G.F.  
Deposited on : 2013-05-12  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	<div> <div>8%</div> <div>71%</div> <div>25%</div> <div>• •</div> </div>
2	B	169	<div> <div>8%</div> <div>72%</div> <div>20%</div> <div>7%</div> <div>•</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2395	1488	433	459	15			

- Molecule 2 is a protein called Hemagglutinin HA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	169	Total	C	N	O	S	0	0	0
			1368	845	237	279	7			

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			32	17	1	14		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	65	Total	O	0	0
			65	65		
5	B	23	Total	O	0	0
			23	23		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Hemagglutinin HA1



#### • Molecule 2: Hemagglutinin HA2



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.24Å 116.24Å 296.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.65 – 2.60 47.65 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.65-2.60) 99.8 (47.65-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.18 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.218 , 0.256 0.220 , 0.262	Depositor DCC
$R_{free}$ test set	1229 reflections (5.11%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.4	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.9	EDS
Estimated twinning fraction	0.006 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/3*k+1/3*l 0.021 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+1/3*l,-4/3*h+4/3*k+1/3*l 0.007 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k+1/3*l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 24065 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3925	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2441	0.58	1/3300 (0.0%)
2	B	0.37	0/1392	0.53	0/1876
All	All	0.38	0/3833	0.56	1/5176 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	150	ALA	CB-CA-C	10.72	126.18	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2395	0	2352	50	0
2	B	1368	0	1264	35	0
3	A	28	0	26	0	0
3	B	14	0	13	2	0
4	A	32	0	28	0	0
5	A	65	0	0	4	0
5	B	23	0	0	4	0
All	All	3925	0	3683	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:328:ALA:H	2:B:329:GLY:HA3	0.96	1.11
2:B:328:ALA:N	2:B:329:GLY:HA3	1.60	1.06
2:B:327:ILE:O	2:B:327:ILE:HG22	1.60	0.99
2:B:397:GLN:HG3	5:B:620:HOH:O	1.72	0.88
1:A:165:SER:OG	5:A:710:HOH:O	1.91	0.87
2:B:326:ALA:O	2:B:327:ILE:HB	1.75	0.85
2:B:328:ALA:N	2:B:329:GLY:CA	2.44	0.80
2:B:327:ILE:O	2:B:327:ILE:CG2	2.34	0.74
2:B:327:ILE:O	2:B:328:ALA:HB3	1.88	0.73
1:A:121:ARG:NH1	1:A:145:SER:O	2.22	0.73
1:A:96:GLU:OE2	5:A:702:HOH:O	2.09	0.71
1:A:121:ARG:HB3	1:A:144:LEU:HB2	1.74	0.69
2:B:327:ILE:O	2:B:328:ALA:CB	2.41	0.67
2:B:396:LYS:HE3	3:B:501:NAG:H81	1.76	0.66
1:A:87:CYS:O	1:A:215:ASN:ND2	2.29	0.65
1:A:148:ASP:OD2	1:A:148:ASP:N	2.29	0.65
2:B:326:ALA:O	2:B:327:ILE:CB	2.46	0.64
1:A:293:VAL:HG11	2:B:386:LEU:HD13	1.80	0.63
1:A:207:SER:O	1:A:211:ARG:NH2	2.24	0.63
2:B:463:HIS:HB3	2:B:487:ALA:HB2	1.80	0.62
1:A:6:GLY:HA3	5:B:623:HOH:O	1.97	0.62
1:A:192:LEU:N	1:A:239:ASN:OD1	2.33	0.62
1:A:44:LYS:HE2	1:A:269:GLU:HB2	1.83	0.60
1:A:183:THR:HG22	1:A:188:SER:HA	1.84	0.60
1:A:282:ASN:HB3	2:B:377:ILE:HG23	1.84	0.59
1:A:195:VAL:HG23	1:A:200:TYR:HE2	1.67	0.59
1:A:291:ARG:HH21	2:B:388:ASP:HB3	1.67	0.58
1:A:191:LYS:HA	1:A:239:ASN:HD21	1.69	0.58
2:B:380:THR:HG22	2:B:382:GLN:H	1.68	0.57
2:B:475:ASN:ND2	2:B:477:THR:OG1	2.36	0.57
1:A:32:THR:HG23	1:A:285:PHE:HD1	1.69	0.57
3:B:501:NAG:H83	5:B:621:HOH:O	2.03	0.57
1:A:158:SER:HA	5:A:720:HOH:O	2.05	0.56
1:A:154:GLN:OE1	1:A:239:ASN:HB3	2.06	0.55
2:B:326:ALA:HB2	2:B:330:PHE:CE2	2.41	0.55
2:B:375:ARG:NH2	2:B:424:GLU:OE2	2.39	0.54
2:B:485:GLU:HA	2:B:488:MET:HB2	1.90	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:450:ASN:OD1	2:B:450:ASN:N	2.39	0.54
2:B:490:ASN:OD1	2:B:490:ASN:N	2.43	0.52
1:A:194:THR:HG23	1:A:237:SER:HB2	1.92	0.52
1:A:70:LEU:O	1:A:110:LYS:NZ	2.39	0.52
2:B:485:GLU:O	2:B:489:GLN:HG2	2.10	0.52
1:A:49:VAL:HG23	1:A:74:ALA:HB2	1.91	0.52
1:A:163:ARG:HD3	1:A:250:PHE:CZ	2.44	0.51
1:A:139:GLU:OE1	1:A:247:ARG:HD3	2.10	0.51
2:B:387:ILE:HD12	2:B:387:ILE:H	1.75	0.51
1:A:98:LEU:HD11	1:A:168:LEU:HD21	1.91	0.51
1:A:117:TYR:O	5:A:752:HOH:O	2.18	0.50
2:B:440:TYR:CE1	2:B:457:GLY:HA2	2.48	0.49
1:A:147:THR:O	1:A:149:ASN:N	2.45	0.49
2:B:481:SER:HA	2:B:484:ARG:HB2	1.94	0.49
2:B:376:LEU:HD22	2:B:420:LEU:HD21	1.94	0.49
2:B:484:ARG:NE	2:B:488:MET:SD	2.67	0.49
1:A:186:TYR:CZ	1:A:241:ALA:HA	2.48	0.48
1:A:35:ARG:HD3	1:A:303:ARG:HG2	1.95	0.47
1:A:32:THR:HG23	1:A:285:PHE:CD1	2.50	0.47
1:A:179:THR:O	1:A:183:THR:HG23	2.15	0.46
1:A:4:CYS:HA	2:B:458:CYS:HA	1.97	0.46
1:A:32:THR:HG22	1:A:305:LEU:HB2	1.97	0.45
1:A:3:ILE:O	2:B:459:PHE:N	2.37	0.45
1:A:173:ILE:HB	1:A:222:ASP:HB2	1.99	0.44
2:B:440:TYR:HE1	2:B:457:GLY:HA2	1.83	0.44
1:A:191:LYS:HA	1:A:239:ASN:ND2	2.32	0.44
1:A:124:GLY:HA3	1:A:142:TRP:HB3	2.01	0.43
2:B:469:CYS:O	2:B:472:SER:OG	2.31	0.43
1:A:51:LEU:HB3	1:A:54:CYS:O	2.19	0.42
1:A:192:LEU:O	1:A:238:PHE:HA	2.19	0.42
1:A:117:TYR:CE2	1:A:157:LYS:HE3	2.55	0.42
1:A:170:VAL:O	1:A:245:PRO:HB3	2.20	0.41
1:A:149:ASN:HB3	1:A:150:ALA:H	1.66	0.41
1:A:93:VAL:HG21	1:A:224:HIS:CE1	2.55	0.41
1:A:6:GLY:CA	5:B:623:HOH:O	2.63	0.41
2:B:339:ILE:H	2:B:339:ILE:HG12	1.69	0.41
1:A:228:LEU:HA	1:A:228:LEU:HD23	1.89	0.41
1:A:89:PRO:HB2	1:A:220:ARG:HD3	2.03	0.41
1:A:191:LYS:O	1:A:206:PRO:HD2	2.21	0.41
1:A:111:GLU:OE2	1:A:163:ARG:NH1	2.55	0.40
1:A:171:TRP:CH2	1:A:226:LEU:HD23	2.57	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:444:LYS:HB2	2:B:459:PHE:CZ	2.56	0.40
2:B:472:SER:O	2:B:478:TYR:N	2.54	0.40
1:A:106:GLY:HA2	1:A:255:SER:HB3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/314 (99%)	290 (93%)	18 (6%)	4 (1%)	15	30
2	B	167/169 (99%)	144 (86%)	16 (10%)	7 (4%)	3	4
All	All	479/483 (99%)	434 (91%)	34 (7%)	11 (2%)	8	14

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	327	ILE
2	B	328	ALA
2	B	448	ARG
1	A	148	ASP
2	B	484	ARG
1	A	132	SER
1	A	150	ALA
1	A	154	GLN
2	B	464	LYS
2	B	481	SER
2	B	377	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/263 (100%)	237 (90%)	26 (10%)	10	18
2	B	144/144 (100%)	127 (88%)	17 (12%)	6	12
All	All	407/407 (100%)	364 (89%)	43 (11%)	8	15

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	8	HIS
1	A	19	LEU
1	A	20	THR
1	A	24	VAL
1	A	32	THR
1	A	38	ILE
1	A	73	SER
1	A	94	ASN
1	A	111	GLU
1	A	121	ARG
1	A	126	THR
1	A	130	ARG
1	A	144	LEU
1	A	145	SER
1	A	148	ASP
1	A	149	ASN
1	A	162	THR
1	A	176	SER
1	A	181	GLU
1	A	194	THR
1	A	217	LEU
1	A	227	MET
1	A	228	LEU
1	A	256	MET
1	A	316	GLU
2	B	332	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	373	LEU
2	B	385	GLU
2	B	386	LEU
2	B	387	ILE
2	B	388	ASP
2	B	442	ARG
2	B	448	ARG
2	B	450	ASN
2	B	454	ASP
2	B	465	CYS
2	B	468	ASP
2	B	475	ASN
2	B	480	HIS
2	B	484	ARG
2	B	485	GLU
2	B	490	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	213	GLN
2	B	475	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SIA	A	603	4	16,20,21	3.79	6 (37%)	18,28,31	2.47	5 (27%)
4	GAL	A	604	4	12,12,12	0.63	0	17,17,17	1.02	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SIA	A	603	4	-	0/14/34/38	0/1/1/1
4	GAL	A	604	4	-	0/2/22/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603	SIA	C7-C6	-9.29	1.41	1.52
4	A	603	SIA	C3-C2	-6.17	1.42	1.52
4	A	603	SIA	C3-C4	-5.62	1.43	1.52
4	A	603	SIA	C6-C5	-2.36	1.49	1.53
4	A	603	SIA	C10-N5	3.88	1.49	1.34
4	A	603	SIA	O6-C6	6.47	1.54	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	603	SIA	O6-C2-C3	-8.15	94.18	109.86
4	A	603	SIA	O6-C6-C5	-3.37	102.96	108.48
4	A	604	GAL	O5-C1-C2	-2.69	105.50	109.80
4	A	603	SIA	C8-C7-C6	-2.56	107.87	113.01
4	A	604	GAL	C1-O5-C5	2.38	117.86	113.47
4	A	603	SIA	C3-C4-C5	2.47	114.23	111.47
4	A	603	SIA	O9-C9-C8	2.82	117.23	111.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	A	601	1	14,14,15	0.43	0	15,19,21	1.67	1 (6%)
3	NAG	A	602	1	14,14,15	0.48	0	15,19,21	0.95	1 (6%)
3	NAG	B	501	2	14,14,15	0.48	0	15,19,21	1.03	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
3	NAG	B	501	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	NAG	C1-O5-C5	2.28	115.14	112.25
3	A	602	NAG	C1-O5-C5	2.86	115.87	112.25
3	A	601	NAG	C1-O5-C5	5.43	119.14	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	314/314 (100%)	-0.19	3 (0%) 84 81	29, 57, 101, 134	0
2	B	169/169 (100%)	0.35	13 (7%) 16 11	24, 83, 130, 158	0
All	All	483/483 (100%)	-0.00	16 (3%) 50 43	24, 64, 119, 158	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	480	HIS	4.6
2	B	352	GLY	3.9
2	B	353	GLU	3.6
2	B	354	GLY	3.3
2	B	482	LYS	2.7
2	B	441	GLU	2.7
2	B	464	LYS	2.7
2	B	348	GLN	2.4
1	A	190	ASN	2.4
2	B	437	ASP	2.3
2	B	468	ASP	2.3
1	A	148	ASP	2.3
2	B	322	GLY	2.2
1	A	12	ASN	2.2
2	B	346	ARG	2.1
2	B	355	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	SIA	A	603	20/21	0.91	0.28	2.00	84,90,101,108	0
4	GAL	A	604	12/12	0.76	0.38	-	103,139,162,171	0

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	B	501	14/15	0.94	0.15	-0.10	67,84,93,94	0
3	NAG	A	601	14/15	0.82	0.24	-	103,112,120,122	0
3	NAG	A	602	14/15	0.81	0.28	-	110,118,121,121	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.