



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:09 PM GMT

PDB ID : 4KR3
Title : Glycyl-tRNA synthetase mutant E71G in complex with tRNA-Gly
Authors : Qin, X.; Hao, Z.; Tian, Q.; Zhang, Z.; Zhou, C.; Xie, W.
Deposited on : 2013-05-16
Resolution : 3.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1092 (3.28-3.20)
Clashscore	102246	1227 (3.28-3.20)
Ramachandran outliers	100387	1204 (3.28-3.20)
Sidechain outliers	100360	1203 (3.28-3.20)
RSRZ outliers	91569	1097 (3.28-3.20)
RNA backbone	2183	1001 (3.74-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	<div> <div>50%</div> <div>22%</div> <div>•</div> <div>27%</div> </div>
2	C	74	<div> <div>5%</div> <div>45%</div> <div>42%</div> <div>8%</div> <div>5%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	466	3632	2313	632	669	18	0	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	MET	-	EXPRESSION TAG	UNP P41250
A	58	ALA	-	EXPRESSION TAG	UNP P41250
A	59	SER	-	EXPRESSION TAG	UNP P41250
A	71	GLY	GLU	ENGINEERED MUTATION	UNP P41250
A	686	LEU	-	EXPRESSION TAG	UNP P41250
A	687	GLU	-	EXPRESSION TAG	UNP P41250
A	688	HIS	-	EXPRESSION TAG	UNP P41250
A	689	HIS	-	EXPRESSION TAG	UNP P41250
A	690	HIS	-	EXPRESSION TAG	UNP P41250
A	691	HIS	-	EXPRESSION TAG	UNP P41250
A	692	HIS	-	EXPRESSION TAG	UNP P41250
A	693	HIS	-	EXPRESSION TAG	UNP P41250

- Molecule 2 is a RNA chain called Gly-tRNA-CCC.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	70	1496	662	257	505	72	0	0	0

- Molecule 3 is GLYCINE (three-letter code: GLY) (formula: C₂H₅NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total 12	O 12	0	0
5	C	2	Total 2	O 2	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.83Å 87.41Å 80.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.42 – 3.23 38.42 – 3.24	Depositor EDS
% Data completeness (in resolution range)	89.1 (38.42-3.23) 88.7 (38.42-3.24)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.04 (at 3.25Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.239 , 0.292 0.239 , 0.293	Depositor DCC
R_{free} test set	700 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	82.6	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 47.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 14068 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5178	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3716	0.53	1/5033 (0.0%)
2	C	0.30	0/1632	0.84	0/2541
All	All	0.40	0/5348	0.65	1/7574 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	GLY	N-CA-C	5.22	126.15	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3632	0	3495	78	0
2	C	1496	0	754	23	0
3	A	5	0	2	0	0
4	A	31	0	13	2	0
5	A	12	0	0	0	0
5	C	2	0	0	0	0
All	All	5178	0	4264	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:54:U:H3	2:C:58:A:H2	1.24	0.81
4:A:702:ANP:O3A	4:A:702:ANP:O3G	1.99	0.79
1:A:412:CYS:HG	1:A:416:SER:HG	1.31	0.78
2:C:30:G:H1	2:C:40:C:H42	1.45	0.65
1:A:372:LYS:HB3	1:A:396:LYS:HB3	1.80	0.63
1:A:410:ARG:NH2	1:A:522:GLU:OE2	2.32	0.63
1:A:622:THR:O	1:A:628:HIS:ND1	2.28	0.63
1:A:648:LEU:HA	1:A:651:ILE:HD12	1.82	0.62
1:A:316:ASP:OD1	1:A:337:ARG:NH1	2.34	0.61
1:A:629:THR:O	1:A:642:ARG:NH2	2.34	0.61
4:A:702:ANP:O2A	4:A:702:ANP:O1B	2.19	0.59
2:C:68:G:H2'	2:C:69:G:H8	1.68	0.58
1:A:144:PHE:HB2	1:A:225:ASN:HB2	1.86	0.58
1:A:297:ILE:HB	1:A:523:PRO:HG2	1.86	0.57
1:A:576:PHE:O	1:A:580:LEU:HB2	2.05	0.56
1:A:392:ASP:HB3	1:A:406:GLY:HA2	1.88	0.56
2:C:30:G:H2'	2:C:31:A:O4'	2.07	0.55
1:A:412:CYS:SG	1:A:416:SER:OG	2.55	0.54
1:A:438:VAL:HG12	1:A:440:VAL:HG13	1.90	0.54
1:A:279:GLU:OE1	1:A:288:ARG:NH2	2.41	0.53
1:A:582:GLU:O	1:A:586:ARG:HG3	2.08	0.53
1:A:595:ASP:OD1	1:A:595:ASP:N	2.41	0.53
1:A:653:GLN:O	1:A:657:ASN:ND2	2.30	0.53
2:C:18:G:N2	2:C:57:G:H2'	2.25	0.52
1:A:568:SER:O	1:A:570:ASN:N	2.37	0.52
1:A:378:HIS:CG	1:A:390:CYS:HB3	2.45	0.52
1:A:388:CYS:O	1:A:389:ASP:HB2	2.09	0.51
1:A:619:ASP:HB2	1:A:675:GLN:NE2	2.25	0.51
1:A:102:ALA:O	1:A:106:ASN:ND2	2.44	0.51
1:A:415:LEU:HD22	1:A:426:LEU:HD13	1.93	0.51
1:A:258:LEU:HG	1:A:263:GLY:HA2	1.92	0.51
1:A:261:ASN:ND2	1:A:267:PHE:HB3	2.26	0.50
2:C:15:G:H2'	2:C:16:U:C6	2.46	0.50
1:A:314:VAL:HG21	1:A:354:TYR:HA	1.94	0.50
1:A:233:GLY:O	1:A:235:GLY:N	2.41	0.50
2:C:2:C:H2'	2:C:3:G:C8	2.46	0.50
1:A:79:PHE:HA	1:A:99:VAL:HG13	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1:GTP:H2'	2:C:2:C:C6	2.47	0.49
2:C:2:C:H2'	2:C:3:G:H8	1.77	0.49
1:A:139:GLY:O	1:A:143:LYS:HB2	2.13	0.49
1:A:556:VAL:HG23	1:A:557:VAL:HG13	1.93	0.48
1:A:634:ASP:HB2	1:A:662:TRP:CE2	2.49	0.48
1:A:562:CYS:HB2	1:A:652:VAL:HG11	1.95	0.48
1:A:281:SER:HB2	2:C:1:GTP:N7	2.29	0.47
1:A:247:ALA:N	1:A:296:GLU:OE2	2.47	0.47
1:A:372:LYS:HB2	1:A:537:HIS:CE1	2.50	0.47
1:A:298:GLU:OE2	1:A:522:GLU:HG3	2.15	0.47
1:A:326:ALA:HA	1:A:331:GLN:HG3	1.97	0.47
1:A:437:THR:HA	1:A:510:THR:HA	1.97	0.47
1:A:359:ILE:O	1:A:363:LEU:HG	2.15	0.46
2:C:7:C:O2'	2:C:49:C:OP2	2.27	0.46
2:C:26:G:H2'	2:C:27:C:C6	2.50	0.46
1:A:529:ARG:O	1:A:533:THR:HG23	2.15	0.46
2:C:8:U:H5'	2:C:49:C:OP2	2.15	0.46
1:A:410:ARG:NH1	1:A:414:ASP:OD2	2.49	0.45
1:A:563:SER:HB3	1:A:612:VAL:HG11	1.97	0.45
1:A:565:LEU:HB2	1:A:617:THR:HG22	1.98	0.45
1:A:599:SER:OG	2:C:38:U:OP2	2.16	0.45
1:A:618:ILE:HG23	1:A:622:THR:HG21	1.98	0.45
1:A:606:ARG:NE	2:C:27:C:OP1	2.48	0.45
1:A:565:LEU:O	1:A:617:THR:HA	2.16	0.45
1:A:410:ARG:HG3	1:A:520:VAL:HG13	1.98	0.44
2:C:37:A:H4'	2:C:38:U:OP2	2.18	0.44
2:C:24:A:H2'	2:C:25:U:O4'	2.17	0.44
1:A:587:HIS:CD2	1:A:649:PRO:HB2	2.52	0.44
1:A:131:PRO:HG2	1:A:134:VAL:HG23	1.99	0.44
1:A:610:ILE:HG13	1:A:612:VAL:HG23	2.00	0.44
2:C:68:G:H2'	2:C:69:G:C8	2.50	0.43
1:A:323:SER:O	1:A:327:GLN:HG3	2.18	0.43
1:A:553:PHE:O	1:A:635:ARG:NH2	2.42	0.43
1:A:620:PHE:H	1:A:675:GLN:HE21	1.66	0.43
1:A:649:PRO:O	1:A:653:GLN:HG3	2.18	0.43
1:A:138:SER:OG	1:A:140:HIS:ND1	2.32	0.43
1:A:640:GLN:NE2	2:C:36:C:H41	2.17	0.43
2:C:28:A:H2'	2:C:29:A:C8	2.54	0.43
1:A:541:VAL:HG22	1:A:549:THR:HG22	2.00	0.42
1:A:319:LEU:O	1:A:335:LYS:HA	2.19	0.42
1:A:616:VAL:HA	1:A:631:THR:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLN:HA	1:A:117:GLN:HB2	2.01	0.42
1:A:299:HIS:O	1:A:520:VAL:HA	2.18	0.42
1:A:407:CYS:HA	1:A:523:PRO:HA	2.01	0.42
1:A:530:ILE:O	1:A:534:VAL:HG23	2.19	0.42
1:A:242:LEU:HB3	1:A:276:PHE:CD1	2.54	0.42
1:A:283:ARG:NH2	2:C:1:GTP:O6	2.52	0.42
1:A:308:HIS:HA	1:A:309:PRO:HD3	1.85	0.42
1:A:368:ILE:HG23	1:A:537:HIS:CE1	2.55	0.42
1:A:532:TYR:O	1:A:535:PHE:HB2	2.20	0.42
1:A:137:THR:HG21	1:A:418:HIS:HA	2.02	0.42
1:A:255:LYS:HB3	1:A:255:LYS:HE2	1.79	0.42
1:A:625:LYS:HD3	1:A:625:LYS:HA	1.76	0.41
2:C:51:G:H2'	2:C:52:G:C8	2.56	0.41
2:C:19:G:N2	2:C:57:G:H1'	2.35	0.41
1:A:295:ALA:HB3	1:A:525:PHE:HB2	2.01	0.41
1:A:629:THR:HA	1:A:644:GLU:HA	2.03	0.40
1:A:246:THR:O	1:A:271:GLN:NE2	2.54	0.40
1:A:644:GLU:HG3	1:A:647:GLU:HG2	2.03	0.40
1:A:287:ILE:O	1:A:529:ARG:NE	2.54	0.40
1:A:560:PHE:HB2	1:A:612:VAL:HG22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	458/637 (72%)	420 (92%)	31 (7%)	7 (2%)	13 53

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	389	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	379	MET
1	A	569	GLN
1	A	671	LEU
1	A	234	PRO
1	A	411	SER
1	A	628	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	380/555 (68%)	358 (94%)	22 (6%)	25 65

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	SER
1	A	117	GLN
1	A	132	GLU
1	A	259	GLU
1	A	317	LEU
1	A	322	TYR
1	A	331	GLN
1	A	343	GLU
1	A	350	THR
1	A	352	LEU
1	A	374	ARG
1	A	389	ASP
1	A	394	GLU
1	A	410	ARG
1	A	427	VAL
1	A	507	PHE
1	A	580	LEU
1	A	599	SER
1	A	606	ARG
1	A	646	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	654	ASP
1	A	659	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	ASN
1	A	331	GLN
1	A	640	GLN
1	A	675	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	69/74 (93%)	13 (18%)	1 (1%)

All (13) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	9	G
2	C	18	G
2	C	19	G
2	C	21	A
2	C	33	U
2	C	37	A
2	C	38	U
2	C	48	C
2	C	51	G
2	C	60	U
2	C	65	G
2	C	70	C
2	C	71	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	C	1	GTP

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GTP	C	1	2	25,34,34	2.72	8 (32%)	34,54,54	2.06	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	C	1	2	-	0/18/38/38	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	GTP	C8-N7	-5.69	1.23	1.34
2	C	1	GTP	C2'-C3'	-4.98	1.39	1.53
2	C	1	GTP	O4'-C4'	-2.88	1.38	1.45
2	C	1	GTP	PB-O2B	-2.17	1.45	1.54
2	C	1	GTP	O3'-C3'	-2.15	1.37	1.43
2	C	1	GTP	C5-C4	2.83	1.46	1.40
2	C	1	GTP	C2-N2	5.47	1.45	1.34
2	C	1	GTP	O6-C6	7.10	1.41	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GTP	C4-C5-N7	-5.08	104.81	109.48
2	C	1	GTP	N3-C2-N1	-3.94	121.45	127.44
2	C	1	GTP	C1'-N9-C4	-3.76	121.27	126.94
2	C	1	GTP	PA-O3A-PB	-3.66	122.45	132.73
2	C	1	GTP	C5-C6-N1	-3.47	118.85	123.59
2	C	1	GTP	PB-O3B-PG	-3.00	122.62	132.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GTP	C6-C5-C4	-2.45	117.97	120.90
2	C	1	GTP	O5'-C5'-C4'	2.56	118.57	109.12
2	C	1	GTP	C6-N1-C2	4.21	121.78	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GTP	3	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GLY	A	701	-	1,4,4	0.32	0	0,4,4	0.00	-
4	ANP	A	702	-	27,33,33	2.78	8 (29%)	30,52,52	3.33	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLY	A	701	-	-	0/0/2/2	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	702	-	-	0/12/38/38	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	ANP	O4'-C4'	-2.39	1.39	1.45
4	A	702	ANP	O3'-C3'	-2.04	1.38	1.43
4	A	702	ANP	C2'-C3'	-2.04	1.47	1.53
4	A	702	ANP	PB-N3B	2.43	1.69	1.63
4	A	702	ANP	C6-N6	2.74	1.43	1.34
4	A	702	ANP	C8-N7	2.84	1.40	1.34
4	A	702	ANP	PG-O1G	4.19	1.50	1.46
4	A	702	ANP	PB-O1B	11.59	1.59	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	702	ANP	O1B-PB-N3B	-12.00	93.49	111.90
4	A	702	ANP	N3-C2-N1	-10.44	120.90	128.89
4	A	702	ANP	O3A-PB-N3B	-3.29	97.38	106.44
4	A	702	ANP	PA-O3A-PB	-2.83	123.18	132.67
4	A	702	ANP	O2B-PB-O3A	3.09	119.13	105.09
4	A	702	ANP	O2B-PB-O1B	5.13	120.70	110.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	702	ANP	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	466/637 (73%)	-0.01	4 (0%)	85 79	51, 76, 106, 140	0
2	C	69/74 (93%)	0.23	4 (5%)	26 17	65, 100, 142, 159	0
All	All	535/711 (75%)	0.02	8 (1%)	76 65	51, 78, 117, 159	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	381	ASN	4.0
2	C	20	U	3.6
1	A	224	PHE	2.8
2	C	72	C	2.8
1	A	441	VAL	2.7
2	C	19	G	2.6
1	A	274	ASN	2.3
2	C	71	G	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GTP	C	1	32/32	0.82	0.27	-0.09	113,120,169,173	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GLY	A	701	5/5	0.73	0.26	0.44	110,111,111,111	0
4	ANP	A	702	31/31	0.92	0.23	0.30	64,69,239,331	0

6.5 Other polymers [i](#)

There are no such residues in this entry.