



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:06 AM GMT

PDB ID : 3K VX  
Title : JNK3 bound to aminopyrimidine inhibitor, SR-3562  
Authors : Habel, J.E.; Laughlin, J.D.; LoGrasso, P.  
Deposited on : 2009-11-30  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>8%</div> <div>74%</div> <div>13%</div> <div>•</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FMY	A	403	-	-	X	-




In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	1	0
			2537	1630	429	459	19			

- Molecule 2 is N-[(2Z)-4-(3-FLUORO-5-MORPHOLIN-4-YLPHENYL)PYRIMIDIN-2(1H)-YLIDENE]-4-(3-MORPHOLIN-4-YL-1H-1,2,4-TRIAZOL-1-YL)ANILINE (three-letter code: FMY) (formula:  $C_{26}H_{27}FN_8O_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			37	26	1	8	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	27	Total O 27 27	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 10



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.36 Å 125.64 Å 68.74 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.38 – 2.40 23.38 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.6 (23.38-2.40) 98.8 (23.38-2.40)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.41 (at 2.38 Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.220 , 0.269 0.227 , 0.285	Depositor DCC
$R_{free}$ test set	707 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	55.6	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 48.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 14037 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2601	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OCY, FMY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	2/2566 (0.1%)	0.92	2/3488 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	CYS	CB-SG	-6.84	1.70	1.82
1	A	111	GLU	CG-CD	5.13	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	MET	CG-SD-CE	-5.55	91.31	100.20
1	A	112	LEU	CA-CB-CG	5.10	127.02	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2537	0	2421	55	0
2	A	37	0	27	22	0
3	A	27	0	0	0	1
All	All	2601	0	2448	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:403:FMY:H03	2:A:403:FMY:C09	1.32	1.55
2:A:403:FMY:H09A	2:A:403:FMY:C03	1.36	1.50
1:A:287:MET:HG2	1:A:299:GLU:HG2	1.27	1.12
1:A:74:ALA:H	2:A:403:FMY:H15	1.15	1.11
1:A:212:ARG:HH11	1:A:212:ARG:HG3	0.96	1.10
2:A:403:FMY:H05	2:A:403:FMY:C09	1.85	1.07
2:A:403:FMY:H08	2:A:403:FMY:H01	1.39	1.05
1:A:212:ARG:NH1	1:A:212:ARG:HG3	1.72	0.89
1:A:77:ILE:HD13	1:A:77:ILE:H	1.35	0.89
2:A:403:FMY:O20	2:A:403:FMY:H05	1.72	0.89
1:A:212:ARG:HH11	1:A:212:ARG:CG	1.83	0.88
2:A:403:FMY:H01	2:A:403:FMY:N32	1.88	0.85
1:A:287:MET:CG	1:A:299:GLU:HG2	2.07	0.82
1:A:211:ALA:O	1:A:212:ARG:HB2	1.78	0.81
1:A:287:MET:HG2	1:A:299:GLU:CG	2.10	0.80
1:A:211:ALA:O	1:A:212:ARG:CB	2.30	0.80
1:A:193:SER:O	2:A:403:FMY:H10	1.82	0.80
1:A:74:ALA:N	2:A:403:FMY:H15	1.97	0.79
1:A:77:ILE:CD1	1:A:77:ILE:H	2.00	0.74
1:A:364:ASP:O	1:A:366:ALA:N	2.23	0.72
2:A:403:FMY:C01	2:A:403:FMY:N32	2.52	0.72
1:A:161:LEU:HD12	1:A:257:VAL:HA	1.71	0.71
1:A:212:ARG:NH1	1:A:212:ARG:CG	2.45	0.69
1:A:365:PRO:O	1:A:366:ALA:HB2	1.95	0.66
1:A:187:HIS:O	1:A:188:ARG:HB2	1.96	0.65
1:A:173:MET:HE2	1:A:249:VAL:HG22	1.79	0.65
1:A:211:ALA:O	1:A:212:ARG:CG	2.51	0.59
1:A:70:ILE:HD11	1:A:148:LEU:HD11	1.84	0.58
2:A:403:FMY:C05	2:A:403:FMY:C09	2.73	0.57
1:A:149:MET:H	2:A:403:FMY:H35	1.70	0.56
1:A:161:LEU:HD22	1:A:319:PRO:HD2	1.88	0.55
1:A:120:HIS:CE1	1:A:176:GLY:HA2	2.43	0.54
1:A:75:GLN:OE1	2:A:403:FMY:H16A	2.09	0.53
2:A:403:FMY:H09A	2:A:403:FMY:H05	1.76	0.52
1:A:120:HIS:HB3	1:A:123:ILE:CG1	2.40	0.51
1:A:120:HIS:HB3	1:A:123:ILE:HG12	1.92	0.51
1:A:301:ARG:HG3	1:A:302:PRO:HD2	1.93	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:ARG:NH2	1:A:86:LEU:HD22	2.26	0.51
1:A:318:PHE:C	1:A:319:PRO:O	2.49	0.51
2:A:403:FMY:H08	2:A:403:FMY:C01	2.27	0.50
1:A:70:ILE:HG22	1:A:78:VAL:O	2.12	0.49
1:A:97:ARG:O	1:A:100:GLN:NE2	2.20	0.49
1:A:149:MET:H	2:A:403:FMY:C35	2.26	0.48
1:A:318:PHE:O	1:A:319:PRO:O	2.30	0.48
1:A:287:MET:O	1:A:289:LYS:N	2.47	0.48
1:A:328:LYS:HD2	1:A:328:LYS:HA	1.71	0.48
2:A:403:FMY:C03	2:A:403:FMY:C09	2.28	0.47
2:A:403:FMY:H03	2:A:403:FMY:C13	2.31	0.47
1:A:365:PRO:O	1:A:366:ALA:CB	2.61	0.47
1:A:197:VAL:HG23	1:A:197:VAL:O	2.15	0.46
1:A:53:VAL:HG11	1:A:79:CYS:SG	2.56	0.46
1:A:364:ASP:O	1:A:365:PRO:C	2.54	0.45
1:A:287:MET:C	1:A:289:LYS:H	2.19	0.45
1:A:75:GLN:OE1	1:A:93:LYS:HE2	2.17	0.45
2:A:403:FMY:H09A	2:A:403:FMY:C05	2.41	0.44
1:A:149:MET:N	2:A:403:FMY:H35	2.33	0.44
1:A:210:LEU:HD12	1:A:210:LEU:HA	1.71	0.44
1:A:149:MET:H	2:A:403:FMY:C34	2.31	0.43
1:A:242:GLU:HG2	1:A:242:GLU:H	1.33	0.43
1:A:94:LYS:HE3	1:A:143:TYR:OH	2.19	0.42
1:A:152:ASN:OD1	1:A:154:OCY:HB3	2.19	0.42
1:A:338:LYS:HD3	1:A:348:ILE:HD12	2.02	0.41
1:A:171:TYR:O	1:A:175:OCY:HB3	2.20	0.41
1:A:114:LEU:HD13	1:A:185:ILE:HD11	2.01	0.41
1:A:287:MET:CG	1:A:299:GLU:CG	2.84	0.41
1:A:283:CYS:HA	1:A:284:PRO:HD2	1.86	0.40
1:A:149:MET:H	2:A:403:FMY:H34	1.86	0.40
1:A:319:PRO:O	1:A:320:ALA:HB2	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:417:HOH:O	3:A:418:HOH:O[3_555]	2.04	0.16



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	322/364 (88%)	307 (95%)	10 (3%)	5 (2%)	12 16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	319	PRO
1	A	366	ALA
1	A	288	LYS
1	A	365	PRO
1	A	320	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	255/326 (78%)	244 (96%)	11 (4%)	35 55

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ILE
1	A	86	LEU
1	A	95	LEU
1	A	112	LEU
1	A	199	SER
1	A	210	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	212	ARG
1	A	242	GLU
1	A	287	MET
1	A	316	SER
1	A	383	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	OCY	A	154	1	7,8,9	1.10	1 (14%)	5,8,10	2.53	1 (20%)
1	OCY	A	175	1	7,8,9	1.71	1 (14%)	5,8,10	1.80	1 (20%)
1	OCY	A	201	1	7,8,9	0.86	0	5,8,10	2.86	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCY	A	154	1	-	0/5/7/9	0/0/0/0
1	OCY	A	175	1	-	0/5/7/9	0/0/0/0
1	OCY	A	201	1	-	0/5/7/9	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	175	OCY	CB-SG	-3.99	1.73	1.81
1	A	154	OCY	CB-SG	-2.46	1.76	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	OCY	CE-CD-SG	-2.24	106.05	113.77
1	A	201	OCY	O-C-CA	-2.21	119.73	125.49
1	A	154	OCY	CB-SG-CD	5.13	117.79	102.41
1	A	201	OCY	CB-SG-CD	5.70	119.51	102.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	154	OCY	1	0
1	A	175	OCY	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMY	A	403	-	36,42,42	1.85	7 (19%)	46,58,58	2.99	20 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMY	A	403	-	-	0/14/36/36	0/6/6/6

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	403	FMY	C26-N30	-5.88	1.32	1.44
2	A	403	FMY	C23-N19	-5.77	1.32	1.42
2	A	403	FMY	C03-C01	2.12	1.42	1.38
2	A	403	FMY	C27-N29	2.22	1.39	1.35
2	A	403	FMY	C13-N28	2.33	1.50	1.46
2	A	403	FMY	C06-C22	2.41	1.41	1.37
2	A	403	FMY	C03-C26	2.60	1.42	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	403	FMY	C08-C24-C33	-9.89	107.60	120.54
2	A	403	FMY	C10-C14-N28	-5.15	100.95	110.02
2	A	403	FMY	N17-C05-N30	-4.41	105.17	112.90
2	A	403	FMY	C03-C01-C23	-4.38	114.71	120.40
2	A	403	FMY	C13-N28-C25	-3.67	108.08	117.92
2	A	403	FMY	F37-C22-C07	-3.39	113.69	118.22
2	A	403	FMY	C04-C26-C03	-3.33	116.04	121.21
2	A	403	FMY	C24-C33-N32	-2.89	112.28	116.13
2	A	403	FMY	C14-N28-C25	-2.63	110.85	117.92
2	A	403	FMY	C08-C25-N28	-2.51	118.67	121.36
2	A	403	FMY	O20-C09-C13	-2.10	107.02	111.84
2	A	403	FMY	C15-N29-C27	-2.09	118.53	121.81
2	A	403	FMY	C02-C23-C01	2.05	122.60	119.17
2	A	403	FMY	C34-C33-C24	2.32	126.82	121.84
2	A	403	FMY	C31-N32-C33	2.60	119.35	115.77
2	A	403	FMY	C10-O20-C09	2.72	119.06	109.89
2	A	403	FMY	C16-N29-C15	3.86	119.70	111.59
2	A	403	FMY	F37-C22-C06	4.25	123.88	118.22
2	A	403	FMY	C01-C03-C26	5.93	125.27	119.23
2	A	403	FMY	C06-C24-C33	7.91	130.88	120.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	FMY	22	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	327/364 (89%)	0.27	29 (8%) 12 12	8, 32, 48, 94	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	ALA	5.4
1	A	401	SER	4.4
1	A	249	VAL	4.2
1	A	250	GLY	3.8
1	A	225	VAL	3.6
1	A	118	VAL	3.5
1	A	321	ASP	3.4
1	A	251	CYS	3.2
1	A	340	LEU	3.1
1	A	264	PRO	3.1
1	A	70	ILE	3.1
1	A	226	THR	2.9
1	A	248	SER	2.8
1	A	320	ALA	2.8
1	A	319	PRO	2.7
1	A	170	LEU	2.5
1	A	95	LEU	2.4
1	A	252	ILE	2.4
1	A	247	TRP	2.2
1	A	270	ASP	2.2
1	A	46	ASN	2.2
1	A	342	ILE	2.2
1	A	367	GLU	2.2
1	A	142	VAL	2.2
1	A	173	MET	2.1
1	A	228	TYR	2.1
1	A	144	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	253	MET	2.0
1	A	239	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	OCY	A	201	9/10	0.90	0.12	-	29,31,37,39	0
1	OCY	A	175	9/10	0.94	0.13	-	25,29,48,50	0
1	OCY	A	154	9/10	0.95	0.09	-	42,44,53,54	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FMY	A	403	37/37	0.79	0.24	0.73	53,74,111,111	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.