



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:35 PM GMT

PDB ID : 1KZJ
Title : Crystal Structure of EcTS W80G/dUMP/CB3717 Complex
Authors : Fritz, T.A.; Liu, L.; Finer-Moore, J.S.; Stroud, R.M.
Deposited on : 2002-02-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	 61% 36% •
1	B	264	 63% 33% •
1	C	264	 64% 30% 6%
1	D	264	 61% 38% •
1	E	264	 60% 35% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	264	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CXM	E	1	-	-	X	-
3	CB3	E	309	-	-	-	X
3	CB3	F	311	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2143	1366	370	395	12	0	0	0
1	B	264	2143	1366	370	395	12	0	0	0
1	C	264	2143	1366	370	395	12	0	0	0
1	D	264	2143	1366	370	395	12	0	0	0
1	E	261	2122	1352	367	391	12	0	0	0
1	F	264	2143	1366	370	395	12	0	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
A	80	GLY	TRP	ENGINEERED	UNP P0A884
B	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
B	80	GLY	TRP	ENGINEERED	UNP P0A884
C	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
C	80	GLY	TRP	ENGINEERED	UNP P0A884
D	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
D	80	GLY	TRP	ENGINEERED	UNP P0A884
E	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
E	80	GLY	TRP	ENGINEERED	UNP P0A884
F	1	CXM	MET	MODIFIED RESIDUE	UNP P0A884
F	80	GLY	TRP	ENGINEERED	UNP P0A884

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C₉H₁₃N₂O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
2	A	1	Total	20	9	2	8	1	0	0
2	B	1	Total	20	9	2	8	1	0	0
2	C	1	Total	20	9	2	8	1	0	0
2	D	1	Total	20	9	2	8	1	0	0
2	E	1	Total	20	9	2	8	1	0	0
2	F	1	Total	20	9	2	8	1	0	0

- Molecule 3 is 10-PROPARGYL-5,8-DIDEAZAFOLIC ACID (three-letter code: CB3) (formula: $C_{24}H_{23}N_5O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	35	24	5	6	0	0
3	B	1	35	24	5	6	0	0
3	C	1	35	24	5	6	0	0
3	D	1	35	24	5	6	0	0
3	E	1	35	24	5	6	0	0
3	F	1	35	24	5	6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	54	54	54	0	0
4	B	54	54	54	0	0
4	C	62	62	62	0	0
4	D	34	34	34	0	0
4	E	36	36	36	0	0
4	F	52	52	52	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Thymidylate synthase




- Molecule 1: Thymidylate synthase




- Molecule 1: Thymidylate synthase




- Molecule 1: Thymidylate synthase



- Molecule 1: Thymidylate synthase



- Molecule 1: Thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.55Å 84.57Å 111.52Å 90.00° 111.37° 90.00°	Depositor
Resolution (Å)	25.35 – 2.60 25.35 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.2 (25.35-2.60) 98.3 (25.35-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 2.60Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.215 , 0.258 0.213 , 0.256	Depositor DCC
R_{free} test set	3545 reflections (7.06%)	DCC
Wilson B-factor (Å ²)	21.8	Xtrriage
Anisotropy	0.230	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.9	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	1 of 50201 reflections (0.002%)	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13459	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.90 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.4247e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CB3, UMP, CXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2190	0.66	1/2972 (0.0%)
1	B	0.39	0/2190	0.65	0/2972
1	C	0.37	0/2190	0.64	0/2972
1	D	0.37	0/2190	0.63	0/2972
1	E	0.39	0/2169	0.66	1/2944 (0.0%)
1	F	0.39	0/2190	0.65	0/2972
All	All	0.38	0/13119	0.65	2/17804 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	147	HIS	N-CA-C	-5.38	96.47	111.00
1	A	27	LEU	N-CA-C	-5.09	97.26	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2143	0	2072	89	0
1	B	2143	0	2072	70	0
1	C	2143	0	2072	80	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2143	0	2073	85	0
1	E	2122	0	2048	89	0
1	F	2143	0	2072	73	0
2	A	20	0	10	0	0
2	B	20	0	10	1	0
2	C	20	0	10	1	0
2	D	20	0	11	1	0
2	E	20	0	11	1	0
2	F	20	0	10	2	0
3	A	35	0	21	2	0
3	B	35	0	21	2	0
3	C	35	0	21	3	0
3	D	35	0	21	6	0
3	E	35	0	21	5	0
3	F	35	0	21	1	0
4	A	54	0	0	2	0
4	B	54	0	0	4	0
4	C	62	0	0	6	0
4	D	34	0	0	1	0
4	E	36	0	0	1	0
4	F	52	0	0	3	0
All	All	13459	0	12597	476	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 476 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:127:ARG:HH21	1:C:127:ARG:HG3	1.27	0.99
1:D:83:TRP:HH2	3:D:307:CB3:H8	1.25	0.98
1:B:233:LYS:HD2	1:B:248:GLU:HG2	1.44	0.97
1:A:59:LEU:HD23	1:A:183:LEU:HD23	1.45	0.96
1:C:127:ARG:CG	1:C:127:ARG:HH21	1.83	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/264 (99%)	246 (94%)	15 (6%)	1 (0%)	39	65
1	B	262/264 (99%)	240 (92%)	18 (7%)	4 (2%)	13	26
1	C	262/264 (99%)	244 (93%)	15 (6%)	3 (1%)	17	36
1	D	262/264 (99%)	240 (92%)	20 (8%)	2 (1%)	24	46
1	E	259/264 (98%)	235 (91%)	20 (8%)	4 (2%)	13	26
1	F	262/264 (99%)	244 (93%)	15 (6%)	3 (1%)	17	36
All	All	1569/1584 (99%)	1449 (92%)	103 (7%)	17 (1%)	17	36

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	94	TYR
1	A	94	TYR
1	B	23	GLY
1	B	79	ILE
1	E	94	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	218 (94%)	13 (6%)	26	50
1	B	231/231 (100%)	215 (93%)	16 (7%)	19	38
1	C	231/231 (100%)	212 (92%)	19 (8%)	14	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	231/231 (100%)	219 (95%)	12 (5%)	29	54
1	E	229/231 (99%)	214 (93%)	15 (7%)	21	40
1	F	231/231 (100%)	219 (95%)	12 (5%)	29	54
All	All	1384/1386 (100%)	1297 (94%)	87 (6%)	22	44

5 of 87 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	138	LEU
1	D	39	GLN
1	F	107	ARG
1	C	176	PHE
1	C	222	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	87	ASN
1	C	215	GLN
1	F	117	ASN
1	C	117	ASN
1	C	134	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CXM	A	1	1	6,10,11	0.95	1 (16%)	4,11,13	0.76	0
1	CXM	B	1	1	6,10,11	1.35	1 (16%)	4,11,13	0.84	0
1	CXM	C	1	1	6,10,11	1.03	1 (16%)	4,11,13	0.75	0
1	CXM	D	1	1	6,10,11	1.19	1 (16%)	4,11,13	0.90	0
1	CXM	E	1	1	6,10,11	1.01	1 (16%)	4,11,13	0.85	0
1	CXM	F	1	1	6,10,11	1.44	1 (16%)	4,11,13	1.05	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
1	CXM	B	1	1	-	0/6/10/12	0/0/0/0
1	CXM	C	1	1	-	0/6/10/12	0/0/0/0
1	CXM	D	1	1	-	0/6/10/12	0/0/0/0
1	CXM	E	1	1	-	0/6/10/12	0/0/0/0
1	CXM	F	1	1	-	0/6/10/12	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	CXM	CA-N	2.09	1.49	1.46
1	E	1	CXM	CA-N	2.22	1.49	1.46
1	C	1	CXM	CA-N	2.35	1.49	1.46
1	D	1	CXM	CA-N	2.63	1.49	1.46
1	B	1	CXM	CA-N	3.11	1.50	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	1	CXM	O-C-CA	-2.03	120.09	125.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1	CXM	2	0
1	B	1	CXM	1	0
1	C	1	CXM	1	0
1	D	1	CXM	3	0
1	E	1	CXM	6	0
1	F	1	CXM	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	UMP	A	300	1	16,21,21	3.94	4 (25%)	23,31,31	2.43	6 (26%)
3	CB3	A	301	-	31,37,37	1.87	7 (22%)	35,51,51	1.99	7 (20%)
2	UMP	B	302	1	16,21,21	4.16	4 (25%)	23,31,31	2.53	5 (21%)
3	CB3	B	303	-	31,37,37	1.81	9 (29%)	35,51,51	1.77	6 (17%)
2	UMP	C	304	1	16,21,21	4.07	4 (25%)	23,31,31	2.05	5 (21%)
3	CB3	C	305	-	31,37,37	1.96	9 (29%)	35,51,51	2.34	12 (34%)
2	UMP	D	306	-	16,21,21	3.30	4 (25%)	23,31,31	1.91	3 (13%)
3	CB3	D	307	-	31,37,37	2.07	10 (32%)	35,51,51	2.05	11 (31%)
2	UMP	E	308	-	16,21,21	3.75	5 (31%)	23,31,31	1.94	3 (13%)
3	CB3	E	309	-	31,37,37	1.90	6 (19%)	35,51,51	1.89	7 (20%)
2	UMP	F	310	1	16,21,21	4.82	5 (31%)	23,31,31	2.43	6 (26%)
3	CB3	F	311	-	31,37,37	2.29	12 (38%)	35,51,51	2.71	13 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	300	1	-	0/6/22/22	0/2/2/2
3	CB3	A	301	-	-	0/21/28/28	0/3/3/3
2	UMP	B	302	1	-	0/6/22/22	0/2/2/2
3	CB3	B	303	-	-	0/21/28/28	0/3/3/3
2	UMP	C	304	1	-	0/6/22/22	0/2/2/2
3	CB3	C	305	-	-	0/21/28/28	0/3/3/3
2	UMP	D	306	-	-	0/6/22/22	0/2/2/2
3	CB3	D	307	-	-	0/21/28/28	0/3/3/3
2	UMP	E	308	-	-	0/6/22/22	0/2/2/2
3	CB3	E	309	-	-	0/21/28/28	0/3/3/3
2	UMP	F	310	1	-	0/6/22/22	0/2/2/2
3	CB3	F	311	-	-	0/21/28/28	0/3/3/3

The worst 5 of 79 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	311	CB3	CP1-CP2	-4.09	1.41	1.47
2	E	308	UMP	P-OP2	-2.17	1.46	1.54
2	F	310	UMP	P-OP2	-2.05	1.47	1.54
3	B	303	CB3	C16-C11	2.01	1.42	1.39
3	D	307	CB3	CA-N	2.01	1.49	1.46

The worst 5 of 84 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	311	CB3	CP1-N10-C9	-6.16	102.12	117.28
3	A	301	CB3	N1-C2-N3	-6.04	118.24	127.44
3	F	311	CB3	N1-C2-N3	-6.02	118.27	127.44
2	B	302	UMP	C5-C6-N1	-5.68	106.67	120.58
2	F	310	UMP	C5-C6-N1	-5.31	107.56	120.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	CB3	2	0
2	B	302	UMP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	303	CB3	2	0
2	C	304	UMP	1	0
3	C	305	CB3	3	0
2	D	306	UMP	1	0
3	D	307	CB3	6	0
2	E	308	UMP	1	0
3	E	309	CB3	5	0
2	F	310	UMP	2	0
3	F	311	CB3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	263/264 (99%)	-0.34	1 (0%) 93 91	3, 16, 29, 39	0
1	B	263/264 (99%)	-0.20	2 (0%) 87 85	4, 17, 36, 45	0
1	C	263/264 (99%)	-0.24	1 (0%) 93 91	3, 19, 33, 50	0
1	D	263/264 (99%)	-0.05	7 (2%) 58 51	5, 22, 42, 62	0
1	E	260/264 (98%)	-0.12	5 (1%) 70 64	6, 22, 43, 60	0
1	F	263/264 (99%)	-0.22	2 (0%) 87 85	7, 20, 31, 44	0
All	All	1575/1584 (99%)	-0.20	18 (1%) 82 79	3, 19, 37, 62	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	262	VAL	5.6
1	D	22	THR	4.4
1	B	22	THR	3.3
1	E	22	THR	3.3
1	D	263	ALA	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
1	CXM	F	1	11/12	0.94	0.16	-	21,24,26,27	0
1	CXM	B	1	11/12	0.96	0.12	-	12,15,24,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CXM	C	1	11/12	0.95	0.15	-	22,26,28,28	0
1	CXM	E	1	11/12	0.92	0.15	-	27,30,34,34	0
1	CXM	A	1	11/12	0.96	0.14	-	13,17,24,27	0
1	CXM	D	1	11/12	0.95	0.14	-	21,22,24,25	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CB3	E	309	35/35	0.61	0.39	2.60	67,78,92,92	0
3	CB3	F	311	35/35	0.84	0.22	2.13	18,26,47,48	0
3	CB3	C	305	35/35	0.89	0.20	1.99	10,24,51,53	0
3	CB3	B	303	35/35	0.84	0.23	1.69	24,31,50,52	0
3	CB3	A	301	35/35	0.90	0.19	0.98	9,24,36,37	0
3	CB3	D	307	35/35	0.77	0.28	0.55	43,49,63,64	0
2	UMP	B	302	20/20	0.95	0.16	-0.15	16,17,21,21	0
2	UMP	E	308	20/20	0.95	0.15	-0.47	12,15,21,23	0
2	UMP	D	306	20/20	0.96	0.14	-0.52	18,21,23,24	0
2	UMP	F	310	20/20	0.95	0.14	-0.87	12,15,20,20	0
2	UMP	C	304	20/20	0.96	0.12	-0.98	12,14,17,18	0
2	UMP	A	300	20/20	0.98	0.10	-1.45	4,7,11,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.