



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:36 PM GMT

PDB ID : 1L0V
Title : Quinol-Fumarate Reductase with Menaquinol Molecules
Authors : Iverson, T.M.; Luna-Chavez, C.; Croal, L.R.; Cecchini, G.; Rees, D.C.
Deposited on : 2002-02-13
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	602	
1	M	602	
2	B	243	
2	N	243	
3	C	130	
3	O	130	
4	D	119	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	P	119	 <div>45%49%</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	MQ7	D	700	-	-	X	-
9	FAD	M	803	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 17046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate reductase flavoprotein subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			
1	M	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			

- Molecule 2 is a protein called Fumarate reductase iron-sulfur protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			
2	N	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			


- Molecule 3 is a protein called Fumarate reductase 15 kDa hydrophobic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			
3	O	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			

- Molecule 4 is a protein called Fumarate reductase 13 kDa hydrophobic protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			
4	P	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			

- Molecule 5 is OXALOACETATE ION (three-letter code: OAA) (formula: C₄H₃O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			9	4	5		
5	M	1	Total	C	O	0	0
			9	4	5		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			4	2	2		
6	N	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			7	3	4		
7	N	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 8 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	N	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 9 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
9	M	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 10 is MENAQUINONE-7 (three-letter code: MQ7) (formula: $C_{46}H_{64}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	D	1	Total	C	O	0	0
			24	22	2		
10	B	1	Total	C	O	0	0
			24	22	2		
10	P	1	Total	C	O	0	0
			24	22	2		
10	N	1	Total	C	O	0	0
			24	22	2		

- Molecule 11 is O-DODECANYL OCTAETHYLENE GLYCOL (three-letter code: CE1) (formula: C₂₈H₅₈O₉).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	D	1	Total	C	O	0	0
			37	28	9		
11	D	1	Total	C	O	0	0
			37	28	9		
11	O	1	Total	C	O	0	0
			37	28	9		
11	O	1	Total	C	O	0	0
			37	28	9		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

• Molecule 1: Fumarate reductase flavoprotein subunit




• Molecule 1: Fumarate reductase flavoprotein subunit







• Molecule 2: Fumarate reductase iron-sulfur protein




• Molecule 2: Fumarate reductase iron-sulfur protein




• Molecule 3: Fumarate reductase 15 kDa hydrophobic protein




• Molecule 3: Fumarate reductase 15 kDa hydrophobic protein



- Molecule 4: Fumarate reductase 13 kDa hydrophobic protein



- Molecule 4: Fumarate reductase 13 kDa hydrophobic protein



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.59 Å 138.09 Å 275.25 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-3.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.245 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	17046	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MQ7, OAA, SF4, F3S, FES, CE1, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/4540	0.80	2/6139 (0.0%)
1	M	0.48	0/4540	0.75	1/6139 (0.0%)
2	B	0.98	6/1931 (0.3%)	1.04	11/2617 (0.4%)
2	N	0.80	9/1931 (0.5%)	0.89	8/2617 (0.3%)
3	C	0.50	0/1094	0.69	1/1496 (0.1%)
3	O	0.52	0/1094	0.68	0/1496
4	D	0.53	0/956	0.74	0/1303
4	P	0.99	2/956 (0.2%)	0.78	1/1303 (0.1%)
All	All	0.66	17/17042 (0.1%)	0.81	24/23110 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	N	0	3
4	P	0	1
All	All	0	5

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	9	ASP	C-N	-21.89	0.83	1.34
2	B	243	ARG	CA-C	19.61	2.04	1.52
2	B	243	ARG	C-O	17.19	1.56	1.23
4	P	95	ALA	C-N	13.40	1.57	1.33
2	N	69	VAL	C-N	-13.14	1.03	1.34

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	243	ARG	CB-CG-CD	16.45	154.36	111.60
2	B	243	ARG	NE-CZ-NH1	-11.92	114.34	120.30
2	N	54	ARG	O-C-N	-11.67	104.03	122.70
2	B	242	PRO	CA-N-CD	-11.40	95.53	111.50
2	B	243	ARG	NE-CZ-NH2	11.02	125.81	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	243	ARG	Sidechain
2	N	183	SER	Mainchain
2	N	54	ARG	Mainchain
2	N	64	SER	Mainchain
4	P	9	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4448	0	4335	143	0
1	M	4448	0	4333	662	0
2	B	1888	0	1837	72	0
2	N	1888	0	1833	172	0
3	C	1058	0	1108	52	0
3	O	1058	0	1108	82	0
4	D	926	0	971	90	0
4	P	926	0	970	106	0
5	A	9	0	2	1	0
5	M	9	0	2	1	0
6	B	4	0	0	1	0
6	N	4	0	0	0	0
7	B	7	0	0	0	0
7	N	7	0	0	1	0
8	B	8	0	0	0	0
8	N	8	0	0	0	0
9	A	53	0	31	8	0
9	M	53	0	31	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	B	24	0	23	4	0
10	D	24	0	23	26	0
10	N	24	0	23	14	0
10	P	24	0	23	20	0
11	D	74	0	116	8	0
11	O	74	0	116	1	0
All	All	17046	0	16885	1285	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1285 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:HIS:NE2	9:A:703:FAD:HM82	1.16	1.45
1:M:98:LEU:CD2	2:N:132:ASN:HD21	1.28	1.42
1:M:44:HIS:NE2	9:M:803:FAD:HM82	1.06	1.32
1:M:493:ASP:OD2	2:N:50:ASP:HA	1.27	1.31
1:M:98:LEU:HD23	2:N:132:ASN:ND2	1.42	1.31

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/602 (96%)	529 (92%)	37 (6%)	9 (2%)	12	48
1	M	575/602 (96%)	355 (62%)	153 (27%)	67 (12%)	0	3
2	B	241/243 (99%)	215 (89%)	22 (9%)	4 (2%)	11	47
2	N	241/243 (99%)	205 (85%)	34 (14%)	2 (1%)	24	62
3	C	128/130 (98%)	121 (94%)	5 (4%)	2 (2%)	12	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	O	128/130 (98%)	113 (88%)	12 (9%)	3 (2%)	8	39
4	D	117/119 (98%)	101 (86%)	13 (11%)	3 (3%)	7	36
4	P	117/119 (98%)	99 (85%)	12 (10%)	6 (5%)	2	19
All	All	2122/2188 (97%)	1738 (82%)	288 (14%)	96 (4%)	3	21

5 of 96 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	242	PRO
4	D	99	VAL
1	M	57	GLN
1	M	76	LEU
1	M	85	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/475 (97%)	448 (97%)	12 (3%)	54	81
1	M	460/475 (97%)	400 (87%)	60 (13%)	5	22
2	B	205/205 (100%)	197 (96%)	8 (4%)	39	75
2	N	205/205 (100%)	191 (93%)	14 (7%)	20	57
3	C	111/111 (100%)	104 (94%)	7 (6%)	22	60
3	O	111/111 (100%)	103 (93%)	8 (7%)	18	54
4	D	97/97 (100%)	94 (97%)	3 (3%)	47	79
4	P	97/97 (100%)	93 (96%)	4 (4%)	37	74
All	All	1746/1776 (98%)	1630 (93%)	116 (7%)	21	59

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	161	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	M	311	VAL
3	O	37	TRP
1	M	174	ASN
1	M	242	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	148	GLN
1	M	256	ASN
2	N	225	GLN
1	M	174	ASN
1	M	204	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	OAA	A	702	-	2,8,8	2.11	1 (50%)	2,10,10	0.90	0
9	FAD	A	703	1	48,58,58	1.70	12 (25%)	54,89,89	2.47	12 (22%)
6	FES	B	244	2	0,4,4	0.00	-	0,4,4	0.00	-
7	F3S	B	245	2	0,9,9	0.00	-	0,15,15	0.00	-
8	SF4	B	246	2	0,12,12	0.00	-	0,24,24	0.00	-
10	MQ7	B	701	-	25,25,49	3.48	6 (24%)	33,34,63	2.21	8 (24%)
10	MQ7	D	700	-	25,25,49	3.47	5 (20%)	33,34,63	2.20	11 (33%)
11	CE1	D	710	-	36,36,36	1.00	0	35,35,35	1.71	15 (42%)
11	CE1	D	810	-	36,36,36	1.10	0	35,35,35	1.74	16 (45%)
5	OAA	M	802	-	2,8,8	1.79	1 (50%)	2,10,10	1.93	1 (50%)
9	FAD	M	803	1	48,58,58	1.80	12 (25%)	54,89,89	2.26	9 (16%)
6	FES	N	244	2	0,4,4	0.00	-	0,4,4	0.00	-
7	F3S	N	245	2	0,9,9	0.00	-	0,15,15	0.00	-
8	SF4	N	246	2	0,12,12	0.00	-	0,24,24	0.00	-
10	MQ7	N	801	-	25,25,49	3.66	5 (20%)	33,34,63	2.34	9 (27%)
11	CE1	O	811	-	36,36,36	1.25	4 (11%)	35,35,35	1.91	13 (37%)
11	CE1	O	812	-	36,36,36	1.12	0	35,35,35	1.59	9 (25%)
10	MQ7	P	800	-	25,25,49	3.40	5 (20%)	33,34,63	2.18	10 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OAA	A	702	-	-	0/2/8/8	0/0/0/0
9	FAD	A	703	1	-	0/30/50/50	0/6/6/6
6	FES	B	244	2	-	0/0/4/4	0/1/1/1
7	F3S	B	245	2	-	0/0/24/24	0/0/3/3
8	SF4	B	246	2	-	0/0/48/48	0/6/5/5
10	MQ7	B	701	-	-	0/13/33/61	0/2/2/2
10	MQ7	D	700	-	-	0/13/33/61	0/2/2/2
11	CE1	D	710	-	-	0/34/34/34	0/0/0/0
11	CE1	D	810	-	-	0/34/34/34	0/0/0/0
5	OAA	M	802	-	-	0/2/8/8	0/0/0/0
9	FAD	M	803	1	-	0/30/50/50	0/6/6/6
6	FES	N	244	2	-	0/0/4/4	0/1/1/1
7	F3S	N	245	2	-	0/0/24/24	0/0/3/3
8	SF4	N	246	2	-	0/0/48/48	0/6/5/5
10	MQ7	N	801	-	-	0/13/33/61	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	CE1	O	811	-	-	0/34/34/34	0/0/0/0
11	CE1	O	812	-	-	0/34/34/34	0/0/0/0
10	MQ7	P	800	-	-	0/13/33/61	0/2/2/2

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	N	801	MQ7	C11-C12	-11.17	1.33	1.50
10	D	700	MQ7	C11-C12	-10.33	1.34	1.50
10	P	800	MQ7	C11-C12	-10.16	1.34	1.50
10	B	701	MQ7	C11-C12	-9.58	1.35	1.50
10	P	800	MQ7	C14-C13	-5.76	1.36	1.50

The worst 5 of 113 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	703	FAD	N3A-C2A-N1A	-12.55	119.29	128.89
9	M	803	FAD	N3A-C2A-N1A	-10.52	120.84	128.89
10	N	801	MQ7	C11-C3-C4	-7.13	110.33	118.47
10	D	700	MQ7	C11-C3-C4	-6.90	110.59	118.47
10	B	701	MQ7	C11-C3-C4	-6.88	110.61	118.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 109 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	702	OAA	1	0
9	A	703	FAD	8	0
6	B	244	FES	1	0
10	B	701	MQ7	4	0
10	D	700	MQ7	26	0
11	D	710	CE1	3	0
11	D	810	CE1	5	0
5	M	802	OAA	1	0
9	M	803	FAD	24	0
7	N	245	F3S	1	0
10	N	801	MQ7	14	0
11	O	812	CE1	1	0
10	P	800	MQ7	20	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.