



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:11 AM GMT

PDB ID : 3L3S
Title : Crystal structure of an enoyl-CoA hydrotase/isomerase family protein from *Silicibacter pomeroyi*
Authors : Eswaramoorthy, S.; Silberstein, M.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2009-12-17
Resolution : 2.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	 2% 76% 11% • 12%
1	B	263	 % 76% 11% • 11%
1	C	263	 74% 13% • 11%
1	D	263	 % 76% 11% • 12%
1	E	263	 % 74% 12% • 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%79%8%•12%</div></div>
1	G	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>3%74%12%•11%</div></div>
1	H	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%77%10%•12%</div></div>
1	I	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>3%77%10%•11%</div></div>
1	J	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>3%75%13%•11%</div></div>
1	K	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>9%74%12%•12%</div></div>
1	L	263	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>11%70%14%•12%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 20649 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-CoA hydratase/isomerase family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	Se	0	0	0
			1711	1077	311	310	6	7			
1	B	234	Total	C	N	O	S	Se	0	0	0
			1721	1086	312	310	6	7			
1	C	234	Total	C	N	O	S	Se	0	0	0
			1722	1085	310	314	6	7			
1	D	232	Total	C	N	O	S	Se	0	0	0
			1710	1079	308	310	6	7			
1	E	231	Total	C	N	O	S	Se	0	0	0
			1697	1072	307	305	6	7			
1	F	231	Total	C	N	O	S	Se	0	0	0
			1701	1074	307	307	6	7			
1	G	233	Total	C	N	O	S	Se	0	0	0
			1710	1079	309	309	6	7			
1	H	232	Total	C	N	O	S	Se	0	0	0
			1706	1077	308	308	6	7			
1	I	234	Total	C	N	O	S	Se	0	0	0
			1718	1083	310	312	6	7			
1	J	233	Total	C	N	O	S	Se	0	0	0
			1710	1079	309	309	6	7			
1	K	232	Total	C	N	O	S	Se	0	0	0
			1707	1076	308	310	6	7			
1	L	231	Total	C	N	O	S	Se	0	0	0
			1702	1073	307	309	6	7			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MSE	-	expression tag	UNP Q5LQZ3
A	3	SER	-	expression tag	UNP Q5LQZ3
A	4	LEU	-	expression tag	UNP Q5LQZ3
A	257	GLU	-	expression tag	UNP Q5LQZ3
A	258	GLY	-	expression tag	UNP Q5LQZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	259	HIS	-	expression tag	UNP Q5LQZ3
A	260	HIS	-	expression tag	UNP Q5LQZ3
A	261	HIS	-	expression tag	UNP Q5LQZ3
A	262	HIS	-	expression tag	UNP Q5LQZ3
A	263	HIS	-	expression tag	UNP Q5LQZ3
A	264	HIS	-	expression tag	UNP Q5LQZ3
B	2	MSE	-	expression tag	UNP Q5LQZ3
B	3	SER	-	expression tag	UNP Q5LQZ3
B	4	LEU	-	expression tag	UNP Q5LQZ3
B	257	GLU	-	expression tag	UNP Q5LQZ3
B	258	GLY	-	expression tag	UNP Q5LQZ3
B	259	HIS	-	expression tag	UNP Q5LQZ3
B	260	HIS	-	expression tag	UNP Q5LQZ3
B	261	HIS	-	expression tag	UNP Q5LQZ3
B	262	HIS	-	expression tag	UNP Q5LQZ3
B	263	HIS	-	expression tag	UNP Q5LQZ3
B	264	HIS	-	expression tag	UNP Q5LQZ3
C	2	MSE	-	expression tag	UNP Q5LQZ3
C	3	SER	-	expression tag	UNP Q5LQZ3
C	4	LEU	-	expression tag	UNP Q5LQZ3
C	257	GLU	-	expression tag	UNP Q5LQZ3
C	258	GLY	-	expression tag	UNP Q5LQZ3
C	259	HIS	-	expression tag	UNP Q5LQZ3
C	260	HIS	-	expression tag	UNP Q5LQZ3
C	261	HIS	-	expression tag	UNP Q5LQZ3
C	262	HIS	-	expression tag	UNP Q5LQZ3
C	263	HIS	-	expression tag	UNP Q5LQZ3
C	264	HIS	-	expression tag	UNP Q5LQZ3
D	2	MSE	-	expression tag	UNP Q5LQZ3
D	3	SER	-	expression tag	UNP Q5LQZ3
D	4	LEU	-	expression tag	UNP Q5LQZ3
D	257	GLU	-	expression tag	UNP Q5LQZ3
D	258	GLY	-	expression tag	UNP Q5LQZ3
D	259	HIS	-	expression tag	UNP Q5LQZ3
D	260	HIS	-	expression tag	UNP Q5LQZ3
D	261	HIS	-	expression tag	UNP Q5LQZ3
D	262	HIS	-	expression tag	UNP Q5LQZ3
D	263	HIS	-	expression tag	UNP Q5LQZ3
D	264	HIS	-	expression tag	UNP Q5LQZ3
E	2	MSE	-	expression tag	UNP Q5LQZ3
E	3	SER	-	expression tag	UNP Q5LQZ3
E	4	LEU	-	expression tag	UNP Q5LQZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	257	GLU	-	expression tag	UNP Q5LQZ3
E	258	GLY	-	expression tag	UNP Q5LQZ3
E	259	HIS	-	expression tag	UNP Q5LQZ3
E	260	HIS	-	expression tag	UNP Q5LQZ3
E	261	HIS	-	expression tag	UNP Q5LQZ3
E	262	HIS	-	expression tag	UNP Q5LQZ3
E	263	HIS	-	expression tag	UNP Q5LQZ3
E	264	HIS	-	expression tag	UNP Q5LQZ3
F	2	MSE	-	expression tag	UNP Q5LQZ3
F	3	SER	-	expression tag	UNP Q5LQZ3
F	4	LEU	-	expression tag	UNP Q5LQZ3
F	257	GLU	-	expression tag	UNP Q5LQZ3
F	258	GLY	-	expression tag	UNP Q5LQZ3
F	259	HIS	-	expression tag	UNP Q5LQZ3
F	260	HIS	-	expression tag	UNP Q5LQZ3
F	261	HIS	-	expression tag	UNP Q5LQZ3
F	262	HIS	-	expression tag	UNP Q5LQZ3
F	263	HIS	-	expression tag	UNP Q5LQZ3
F	264	HIS	-	expression tag	UNP Q5LQZ3
G	2	MSE	-	expression tag	UNP Q5LQZ3
G	3	SER	-	expression tag	UNP Q5LQZ3
G	4	LEU	-	expression tag	UNP Q5LQZ3
G	257	GLU	-	expression tag	UNP Q5LQZ3
G	258	GLY	-	expression tag	UNP Q5LQZ3
G	259	HIS	-	expression tag	UNP Q5LQZ3
G	260	HIS	-	expression tag	UNP Q5LQZ3
G	261	HIS	-	expression tag	UNP Q5LQZ3
G	262	HIS	-	expression tag	UNP Q5LQZ3
G	263	HIS	-	expression tag	UNP Q5LQZ3
G	264	HIS	-	expression tag	UNP Q5LQZ3
H	2	MSE	-	expression tag	UNP Q5LQZ3
H	3	SER	-	expression tag	UNP Q5LQZ3
H	4	LEU	-	expression tag	UNP Q5LQZ3
H	257	GLU	-	expression tag	UNP Q5LQZ3
H	258	GLY	-	expression tag	UNP Q5LQZ3
H	259	HIS	-	expression tag	UNP Q5LQZ3
H	260	HIS	-	expression tag	UNP Q5LQZ3
H	261	HIS	-	expression tag	UNP Q5LQZ3
H	262	HIS	-	expression tag	UNP Q5LQZ3
H	263	HIS	-	expression tag	UNP Q5LQZ3
H	264	HIS	-	expression tag	UNP Q5LQZ3
I	2	MSE	-	expression tag	UNP Q5LQZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	3	SER	-	expression tag	UNP Q5LQZ3
I	4	LEU	-	expression tag	UNP Q5LQZ3
I	257	GLU	-	expression tag	UNP Q5LQZ3
I	258	GLY	-	expression tag	UNP Q5LQZ3
I	259	HIS	-	expression tag	UNP Q5LQZ3
I	260	HIS	-	expression tag	UNP Q5LQZ3
I	261	HIS	-	expression tag	UNP Q5LQZ3
I	262	HIS	-	expression tag	UNP Q5LQZ3
I	263	HIS	-	expression tag	UNP Q5LQZ3
I	264	HIS	-	expression tag	UNP Q5LQZ3
J	2	MSE	-	expression tag	UNP Q5LQZ3
J	3	SER	-	expression tag	UNP Q5LQZ3
J	4	LEU	-	expression tag	UNP Q5LQZ3
J	257	GLU	-	expression tag	UNP Q5LQZ3
J	258	GLY	-	expression tag	UNP Q5LQZ3
J	259	HIS	-	expression tag	UNP Q5LQZ3
J	260	HIS	-	expression tag	UNP Q5LQZ3
J	261	HIS	-	expression tag	UNP Q5LQZ3
J	262	HIS	-	expression tag	UNP Q5LQZ3
J	263	HIS	-	expression tag	UNP Q5LQZ3
J	264	HIS	-	expression tag	UNP Q5LQZ3
K	2	MSE	-	expression tag	UNP Q5LQZ3
K	3	SER	-	expression tag	UNP Q5LQZ3
K	4	LEU	-	expression tag	UNP Q5LQZ3
K	257	GLU	-	expression tag	UNP Q5LQZ3
K	258	GLY	-	expression tag	UNP Q5LQZ3
K	259	HIS	-	expression tag	UNP Q5LQZ3
K	260	HIS	-	expression tag	UNP Q5LQZ3
K	261	HIS	-	expression tag	UNP Q5LQZ3
K	262	HIS	-	expression tag	UNP Q5LQZ3
K	263	HIS	-	expression tag	UNP Q5LQZ3
K	264	HIS	-	expression tag	UNP Q5LQZ3
L	2	MSE	-	expression tag	UNP Q5LQZ3
L	3	SER	-	expression tag	UNP Q5LQZ3
L	4	LEU	-	expression tag	UNP Q5LQZ3
L	257	GLU	-	expression tag	UNP Q5LQZ3
L	258	GLY	-	expression tag	UNP Q5LQZ3
L	259	HIS	-	expression tag	UNP Q5LQZ3
L	260	HIS	-	expression tag	UNP Q5LQZ3
L	261	HIS	-	expression tag	UNP Q5LQZ3
L	262	HIS	-	expression tag	UNP Q5LQZ3
L	263	HIS	-	expression tag	UNP Q5LQZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	264	HIS	-	expression tag	UNP Q5LQZ3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total O 5 5	0	0
2	B	20	Total O 20 20	0	0
2	C	19	Total O 19 19	0	0
2	D	16	Total O 16 16	0	0
2	E	14	Total O 14 14	0	0
2	F	20	Total O 20 20	0	0
2	G	8	Total O 8 8	0	0
2	H	9	Total O 9 9	0	0
2	I	6	Total O 6 6	0	0
2	J	3	Total O 3 3	0	0
2	K	10	Total O 10 10	0	0
2	L	4	Total O 4 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enoyl-CoA hydratase/isomerase family protein




- Molecule 1: Enoyl-CoA hydratase/isomerase family protein



- Molecule 1: Enoyl-CoA hydratase/isomerase family protein




- Molecule 1: Enoyl-CoA hydratase/isomerase family protein










- Molecule 1: Enoyl-CoA hydratase/isomerase family protein





- Molecule 1: Enoyl-CoA hydratase/isomerase family protein




- Molecule 1: Enoyl-CoA hydratase/isomerase family protein




- Molecule 1: Enoyl-CoA hydratase/isomerase family protein





- Molecule 1: Enoyl-CoA hydratase/isomerase family protein








- Molecule 1: Enoyl-CoA hydratase/isomerase family protein



- Molecule 1: Enoyl-CoA hydratase/isomerase family protein



- Molecule 1: Enoyl-CoA hydratase/isomerase family protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.40Å 124.28Å 147.00Å 90.00° 102.44° 90.00°	Depositor
Resolution (Å)	50.00 – 2.32 49.65 – 2.32	Depositor EDS
% Data completeness (in resolution range)	92.3 (50.00-2.32) 75.1 (49.65-2.32)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.32Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.263 , 0.294 0.262 , 0.291	Depositor DCC
R_{free} test set	5004 reflections (4.03%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 4.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 130326 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	20649	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: OCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/1725	0.79	0/2335
1	B	0.53	0/1737	0.80	3/2354 (0.1%)
1	C	0.55	0/1737	0.81	1/2353 (0.0%)
1	D	0.55	0/1725	0.81	1/2337 (0.0%)
1	E	0.55	0/1712	0.80	0/2320
1	F	0.54	0/1716	0.78	1/2325 (0.0%)
1	G	0.56	0/1725	0.85	2/2337 (0.1%)
1	H	0.54	0/1721	0.80	1/2332 (0.0%)
1	I	0.57	0/1733	0.80	4/2347 (0.2%)
1	J	0.54	0/1725	0.78	0/2337
1	K	0.55	0/1722	0.82	3/2333 (0.1%)
1	L	0.56	0/1717	0.84	3/2326 (0.1%)
All	All	0.55	0/20695	0.81	19/28036 (0.1%)

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	111	LEU	CA-CB-CG	7.27	132.01	115.30
1	I	235	LEU	CA-CB-CG	5.91	128.90	115.30
1	F	111	LEU	CA-CB-CG	5.80	128.65	115.30
1	L	55	LEU	CA-CB-CG	5.75	128.52	115.30
1	G	55	LEU	CA-CB-CG	5.71	128.42	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1711	0	1729	11	0
1	B	1721	0	1737	15	0
1	C	1722	0	1738	24	0
1	D	1710	0	1731	12	0
1	E	1697	0	1720	13	0
1	F	1701	0	1725	11	0
1	G	1710	0	1730	17	0
1	H	1706	0	1727	22	0
1	I	1718	0	1736	11	0
1	J	1710	0	1731	12	0
1	K	1707	0	1722	23	0
1	L	1702	0	1720	20	0
2	A	5	0	0	0	0
2	B	20	0	0	1	0
2	C	19	0	0	2	0
2	D	16	0	0	0	0
2	E	14	0	0	0	0
2	F	20	0	0	1	0
2	G	8	0	0	0	0
2	H	9	0	0	0	0
2	I	6	0	0	0	0
2	J	3	0	0	0	0
2	K	10	0	0	0	0
2	L	4	0	0	0	0
All	All	20649	0	20746	164	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 164 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:130:ALA:H	1:F:187:ASN:HD22	1.37	0.72
1:B:130:ALA:H	1:B:187:ASN:HD22	1.35	0.71
1:H:44:ARG:HH22	1:K:44:ARG:HH22	1.40	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:130:ALA:H	1:D:187:ASN:HD22	1.39	0.68
1:I:130:ALA:H	1:I:187:ASN:HD22	1.41	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/263 (86%)	214 (94%)	9 (4%)	4 (2%)	11	8
1	B	229/263 (87%)	221 (96%)	8 (4%)	0	100	100
1	C	229/263 (87%)	216 (94%)	11 (5%)	2 (1%)	21	23
1	D	227/263 (86%)	216 (95%)	7 (3%)	4 (2%)	11	8
1	E	226/263 (86%)	216 (96%)	5 (2%)	5 (2%)	8	6
1	F	226/263 (86%)	217 (96%)	6 (3%)	3 (1%)	15	14
1	G	228/263 (87%)	218 (96%)	6 (3%)	4 (2%)	11	8
1	H	227/263 (86%)	218 (96%)	5 (2%)	4 (2%)	11	8
1	I	229/263 (87%)	218 (95%)	9 (4%)	2 (1%)	21	23
1	J	228/263 (87%)	213 (93%)	10 (4%)	5 (2%)	8	6
1	K	227/263 (86%)	214 (94%)	9 (4%)	4 (2%)	11	8
1	L	226/263 (86%)	208 (92%)	11 (5%)	7 (3%)	5	3
All	All	2729/3156 (86%)	2589 (95%)	96 (4%)	44 (2%)	12	10

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	GLU
1	D	86	PHE
1	G	46	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	86	PHE
1	K	86	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/188 (89%)	153 (91%)	15 (9%)	12	14
1	B	169/188 (90%)	155 (92%)	14 (8%)	14	16
1	C	170/188 (90%)	161 (95%)	9 (5%)	28	37
1	D	169/188 (90%)	158 (94%)	11 (6%)	21	27
1	E	167/188 (89%)	152 (91%)	15 (9%)	12	13
1	F	168/188 (89%)	159 (95%)	9 (5%)	27	36
1	G	168/188 (89%)	151 (90%)	17 (10%)	9	10
1	H	168/188 (89%)	160 (95%)	8 (5%)	31	42
1	I	169/188 (90%)	155 (92%)	14 (8%)	14	16
1	J	168/188 (89%)	153 (91%)	15 (9%)	12	14
1	K	168/188 (89%)	154 (92%)	14 (8%)	14	16
1	L	168/188 (89%)	152 (90%)	16 (10%)	11	12
All	All	2020/2256 (90%)	1863 (92%)	157 (8%)	16	19

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	224	HIS
1	G	229	LEU
1	L	100	ASP
1	G	10	LEU
1	G	117	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	187	ASN
1	H	187	ASN
1	L	122	GLN
1	G	187	ASN
1	H	144	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	A	149	1	7,8,9	2.22	2 (28%)	7,11,13	3.67	2 (28%)
1	OCS	B	149	1	7,8,9	2.14	2 (28%)	7,11,13	3.99	3 (42%)
1	OCS	C	149	1	7,8,9	2.04	2 (28%)	7,11,13	3.49	3 (42%)
1	OCS	D	149	1	7,8,9	2.03	3 (42%)	7,11,13	3.65	2 (28%)
1	OCS	E	149	1	7,8,9	1.97	2 (28%)	7,11,13	3.21	3 (42%)
1	OCS	F	149	1	7,8,9	1.99	2 (28%)	7,11,13	3.12	2 (28%)
1	OCS	G	149	1	7,8,9	1.88	2 (28%)	7,11,13	4.28	2 (28%)
1	OCS	H	149	1	7,8,9	2.27	2 (28%)	7,11,13	4.04	3 (42%)
1	OCS	I	149	1	7,8,9	2.30	3 (42%)	7,11,13	3.51	3 (42%)
1	OCS	J	149	1	7,8,9	2.52	3 (42%)	7,11,13	4.34	3 (42%)
1	OCS	K	149	1	7,8,9	2.14	3 (42%)	7,11,13	4.20	2 (28%)
1	OCS	L	149	1	7,8,9	1.56	2 (28%)	7,11,13	4.37	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	149	1	-	0/4/7/9	0/0/0/0
1	OCS	B	149	1	-	0/4/7/9	0/0/0/0
1	OCS	C	149	1	-	0/4/7/9	0/0/0/0
1	OCS	D	149	1	-	0/4/7/9	0/0/0/0
1	OCS	E	149	1	-	0/4/7/9	0/0/0/0
1	OCS	F	149	1	-	0/4/7/9	0/0/0/0
1	OCS	G	149	1	-	0/4/7/9	0/0/0/0
1	OCS	H	149	1	-	0/4/7/9	0/0/0/0
1	OCS	I	149	1	-	0/4/7/9	0/0/0/0
1	OCS	J	149	1	-	0/4/7/9	0/0/0/0
1	OCS	K	149	1	-	0/4/7/9	0/0/0/0
1	OCS	L	149	1	-	0/4/7/9	0/0/0/0

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	149	OCS	OD3-SG	-4.69	1.30	1.45
1	A	149	OCS	OD3-SG	-4.36	1.31	1.45
1	J	149	OCS	OD3-SG	-4.26	1.32	1.45
1	B	149	OCS	OD3-SG	-3.93	1.33	1.45
1	G	149	OCS	OD3-SG	-3.89	1.33	1.45

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	149	OCS	OD1-SG-CB	-10.77	97.86	106.94
1	J	149	OCS	OD1-SG-CB	-10.74	97.89	106.94
1	H	149	OCS	OD1-SG-CB	-9.58	98.87	106.94
1	K	149	OCS	OD1-SG-CB	-9.52	98.91	106.94
1	G	149	OCS	OD1-SG-CB	-9.50	98.93	106.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	E	149	OCS	1	0
1	G	149	OCS	1	0
1	J	149	OCS	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	L	149	OCS	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/263 (85%)	0.16	4 (1%) 71 78	14, 36, 52, 62	0
1	B	226/263 (85%)	-0.13	2 (0%) 85 89	15, 28, 46, 55	0
1	C	226/263 (85%)	-0.01	0 100 100	15, 30, 47, 56	0
1	D	224/263 (85%)	0.10	3 (1%) 79 84	15, 33, 49, 59	0
1	E	223/263 (84%)	0.02	2 (0%) 85 89	18, 34, 49, 58	0
1	F	223/263 (84%)	0.06	3 (1%) 79 84	17, 31, 45, 58	0
1	G	225/263 (85%)	0.18	8 (3%) 46 55	20, 35, 51, 66	0
1	H	224/263 (85%)	0.33	3 (1%) 79 84	20, 40, 54, 65	0
1	I	226/263 (85%)	0.24	8 (3%) 48 56	16, 38, 52, 59	0
1	J	225/263 (85%)	0.32	9 (4%) 42 50	20, 38, 53, 59	0
1	K	224/263 (85%)	0.64	24 (10%) 8 12	21, 46, 60, 66	0
1	L	223/263 (84%)	0.86	28 (12%) 5 8	21, 50, 61, 63	0
All	All	2693/3156 (85%)	0.23	94 (3%) 48 56	14, 36, 55, 66	0

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	27	PRO	6.2
1	H	85	ALA	5.8
1	G	248	GLY	5.5
1	L	27	PRO	4.6
1	L	197	THR	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	OCS	H	149	9/10	0.97	0.09	-	34,34,38,40	0
1	OCS	K	149	9/10	0.91	0.14	-	35,36,40,40	0
1	OCS	L	149	9/10	0.91	0.09	-	46,48,51,51	0
1	OCS	G	149	9/10	0.95	0.12	-	25,33,42,43	0
1	OCS	A	149	9/10	0.97	0.11	-	25,28,32,33	0
1	OCS	B	149	9/10	0.94	0.13	-	32,35,41,42	0
1	OCS	E	149	9/10	0.95	0.14	-	30,30,39,40	0
1	OCS	F	149	9/10	0.96	0.11	-	30,32,38,38	0
1	OCS	C	149	9/10	0.98	0.11	-	22,23,35,36	0
1	OCS	D	149	9/10	0.95	0.14	-	29,32,36,36	0
1	OCS	I	149	9/10	0.92	0.12	-	34,36,45,46	0
1	OCS	J	149	9/10	0.96	0.11	-	36,37,46,46	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.