



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2017 – 03:58 PM EST

PDB ID : 5L38
Title : The structure of the hexagonal shell protein MSM0272 from the RMM micro-compartment
Authors : Mallette, E.; Kimber, M.S.
Deposited on : 2016-08-03
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	101	<div> <div>3%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>10%</div> </div> </div>
1	B	101	<div> <div>%</div> <div> <div></div> <div>70%</div> <div>14%</div> <div>15%</div> </div> </div>
1	C	101	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>11%</div> </div> </div>
1	D	101	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>12%</div> </div> </div>
1	E	101	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>6%</div> <div>16%</div> </div> </div>
1	F	101	<div> <div></div> <div> <div></div> <div>80%</div> <div>9%</div> <div>11%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	101	<div><div><div></div><div></div><div></div></div><div><div>3%</div><div>78%</div><div>11%</div><div>11%</div></div></div>
1	H	101	<div><div><div></div><div></div><div></div></div><div><div>%</div><div>69%</div><div>18%</div><div>13%</div></div></div>
1	I	101	<div><div><div></div><div></div><div></div></div><div><div></div><div>74%</div><div>12%</div><div>14%</div></div></div>
1	J	101	<div><div><div></div><div></div><div></div></div><div><div></div><div>71%</div><div>17%</div><div>12%</div></div></div>
1	K	101	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>76%</div><div>9%</div><div>15%</div></div></div>
1	L	101	<div><div><div></div><div></div><div></div></div><div><div>3%</div><div>71%</div><div>14%</div><div>15%</div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8323 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MSM0272 - RMM microcompartment shell protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	91	Total	C	N	O	S	0	0	0
			634	393	111	129	1			
1	B	86	Total	C	N	O	S	0	0	0
			603	376	106	120	1			
1	C	90	Total	C	N	O	S	0	0	0
			628	390	110	127	1			
1	D	89	Total	C	N	O	S	0	0	0
			622	387	109	125	1			
1	E	85	Total	C	N	O	S	0	0	0
			597	373	105	118	1			
1	F	90	Total	C	N	O	S	0	0	0
			628	390	110	127	1			
1	G	90	Total	C	N	O	S	0	0	0
			628	390	110	127	1			
1	H	88	Total	C	N	O	S	0	0	0
			616	384	108	123	1			
1	I	87	Total	C	N	O	S	0	0	0
			609	379	107	122	1			
1	J	89	Total	C	N	O	S	0	0	0
			622	387	109	125	1			
1	K	86	Total	C	N	O	S	0	0	0
			603	376	106	120	1			
1	L	86	Total	C	N	O	S	0	0	0
			603	376	106	120	1			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	LEU	-	expression tag	UNP A0QP49
A	95	GLU	-	expression tag	UNP A0QP49
A	96	HIS	-	expression tag	UNP A0QP49
A	97	HIS	-	expression tag	UNP A0QP49
A	98	HIS	-	expression tag	UNP A0QP49

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	99	HIS	-	expression tag	UNP A0QP49
A	100	HIS	-	expression tag	UNP A0QP49
A	101	HIS	-	expression tag	UNP A0QP49
B	94	LEU	-	expression tag	UNP A0QP49
B	95	GLU	-	expression tag	UNP A0QP49
B	96	HIS	-	expression tag	UNP A0QP49
B	97	HIS	-	expression tag	UNP A0QP49
B	98	HIS	-	expression tag	UNP A0QP49
B	99	HIS	-	expression tag	UNP A0QP49
B	100	HIS	-	expression tag	UNP A0QP49
B	101	HIS	-	expression tag	UNP A0QP49
C	94	LEU	-	expression tag	UNP A0QP49
C	95	GLU	-	expression tag	UNP A0QP49
C	96	HIS	-	expression tag	UNP A0QP49
C	97	HIS	-	expression tag	UNP A0QP49
C	98	HIS	-	expression tag	UNP A0QP49
C	99	HIS	-	expression tag	UNP A0QP49
C	100	HIS	-	expression tag	UNP A0QP49
C	101	HIS	-	expression tag	UNP A0QP49
D	94	LEU	-	expression tag	UNP A0QP49
D	95	GLU	-	expression tag	UNP A0QP49
D	96	HIS	-	expression tag	UNP A0QP49
D	97	HIS	-	expression tag	UNP A0QP49
D	98	HIS	-	expression tag	UNP A0QP49
D	99	HIS	-	expression tag	UNP A0QP49
D	100	HIS	-	expression tag	UNP A0QP49
D	101	HIS	-	expression tag	UNP A0QP49
E	94	LEU	-	expression tag	UNP A0QP49
E	95	GLU	-	expression tag	UNP A0QP49
E	96	HIS	-	expression tag	UNP A0QP49
E	97	HIS	-	expression tag	UNP A0QP49
E	98	HIS	-	expression tag	UNP A0QP49
E	99	HIS	-	expression tag	UNP A0QP49
E	100	HIS	-	expression tag	UNP A0QP49
E	101	HIS	-	expression tag	UNP A0QP49
F	94	LEU	-	expression tag	UNP A0QP49
F	95	GLU	-	expression tag	UNP A0QP49
F	96	HIS	-	expression tag	UNP A0QP49
F	97	HIS	-	expression tag	UNP A0QP49
F	98	HIS	-	expression tag	UNP A0QP49
F	99	HIS	-	expression tag	UNP A0QP49
F	100	HIS	-	expression tag	UNP A0QP49

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	101	HIS	-	expression tag	UNP A0QP49
G	94	LEU	-	expression tag	UNP A0QP49
G	95	GLU	-	expression tag	UNP A0QP49
G	96	HIS	-	expression tag	UNP A0QP49
G	97	HIS	-	expression tag	UNP A0QP49
G	98	HIS	-	expression tag	UNP A0QP49
G	99	HIS	-	expression tag	UNP A0QP49
G	100	HIS	-	expression tag	UNP A0QP49
G	101	HIS	-	expression tag	UNP A0QP49
H	94	LEU	-	expression tag	UNP A0QP49
H	95	GLU	-	expression tag	UNP A0QP49
H	96	HIS	-	expression tag	UNP A0QP49
H	97	HIS	-	expression tag	UNP A0QP49
H	98	HIS	-	expression tag	UNP A0QP49
H	99	HIS	-	expression tag	UNP A0QP49
H	100	HIS	-	expression tag	UNP A0QP49
H	101	HIS	-	expression tag	UNP A0QP49
I	94	LEU	-	expression tag	UNP A0QP49
I	95	GLU	-	expression tag	UNP A0QP49
I	96	HIS	-	expression tag	UNP A0QP49
I	97	HIS	-	expression tag	UNP A0QP49
I	98	HIS	-	expression tag	UNP A0QP49
I	99	HIS	-	expression tag	UNP A0QP49
I	100	HIS	-	expression tag	UNP A0QP49
I	101	HIS	-	expression tag	UNP A0QP49
J	94	LEU	-	expression tag	UNP A0QP49
J	95	GLU	-	expression tag	UNP A0QP49
J	96	HIS	-	expression tag	UNP A0QP49
J	97	HIS	-	expression tag	UNP A0QP49
J	98	HIS	-	expression tag	UNP A0QP49
J	99	HIS	-	expression tag	UNP A0QP49
J	100	HIS	-	expression tag	UNP A0QP49
J	101	HIS	-	expression tag	UNP A0QP49
K	94	LEU	-	expression tag	UNP A0QP49
K	95	GLU	-	expression tag	UNP A0QP49
K	96	HIS	-	expression tag	UNP A0QP49
K	97	HIS	-	expression tag	UNP A0QP49
K	98	HIS	-	expression tag	UNP A0QP49
K	99	HIS	-	expression tag	UNP A0QP49
K	100	HIS	-	expression tag	UNP A0QP49
K	101	HIS	-	expression tag	UNP A0QP49
L	94	LEU	-	expression tag	UNP A0QP49

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	95	GLU	-	expression tag	UNP A0QP49
L	96	HIS	-	expression tag	UNP A0QP49
L	97	HIS	-	expression tag	UNP A0QP49
L	98	HIS	-	expression tag	UNP A0QP49
L	99	HIS	-	expression tag	UNP A0QP49
L	100	HIS	-	expression tag	UNP A0QP49
L	101	HIS	-	expression tag	UNP A0QP49

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Na 1 1	0	0
2	I	1	Total Na 1 1	0	0
2	A	1	Total Na 1 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	98	Total O 98 98	0	0
4	B	87	Total O 87 87	0	0
4	C	91	Total O 91 91	0	0
4	D	61	Total O 61 61	0	0
4	E	61	Total O 61 61	0	0
4	F	92	Total O 92 92	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	75	Total 75	O 75	0	0
4	H	83	Total 83	O 83	0	0
4	I	83	Total 83	O 83	0	0
4	J	83	Total 83	O 83	0	0
4	K	51	Total 51	O 51	0	0
4	L	60	Total 60	O 60	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MSM0272 - RMM microcompartment shell protein




- Molecule 1: MSM0272 - RMM microcompartment shell protein




- Molecule 1: MSM0272 - RMM microcompartment shell protein




- Molecule 1: MSM0272 - RMM microcompartment shell protein



- Molecule 1: MSM0272 - RMM microcompartment shell protein




- Molecule 1: MSM0272 - RMM microcompartment shell protein






- Molecule 1: MSM0272 - RMM microcompartment shell protein



- Molecule 1: MSM0272 - RMM microcompartment shell protein





- Molecule 1: MSM0272 - RMM microcompartment shell protein




- Molecule 1: MSM0272 - RMM microcompartment shell protein



- Molecule 1: MSM0272 - RMM microcompartment shell protein



- Molecule 1: MSM0272 - RMM microcompartment shell protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	93.60Å 93.60Å 192.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.20 48.13 – 1.97	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.13-2.20) 99.8 (48.13-1.97)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.97Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.200 , 0.233 0.167 , 0.191	Depositor DCC
R_{free} test set	4182 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.226 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8323	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/640	0.47	0/872
1	B	0.24	0/609	0.41	0/830
1	C	0.30	0/634	0.45	0/864
1	D	0.22	0/628	0.41	0/856
1	E	0.23	0/603	0.43	0/822
1	F	0.25	0/634	0.42	0/864
1	G	0.24	0/634	0.42	0/864
1	H	0.24	0/622	0.42	0/848
1	I	0.24	0/615	0.45	0/838
1	J	0.22	0/628	0.40	0/856
1	K	0.21	0/609	0.40	0/830
1	L	0.23	0/609	0.43	0/830
All	All	0.24	0/7465	0.43	0/10174

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	86	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	634	0	640	12	0
1	B	603	0	611	13	0
1	C	628	0	635	8	0
1	D	622	0	630	8	0
1	E	597	0	606	6	0
1	F	628	0	635	7	0
1	G	628	0	635	10	0
1	H	616	0	625	18	0
1	I	609	0	616	11	0
1	J	622	0	630	15	0
1	K	603	0	611	9	0
1	L	603	0	611	14	0
2	A	1	0	0	0	0
2	G	1	0	0	0	0
2	I	1	0	0	0	0
3	A	1	0	0	1	0
3	C	1	0	0	0	0
4	A	98	0	0	6	1
4	B	87	0	0	7	2
4	C	91	0	0	0	0
4	D	61	0	0	4	0
4	E	61	0	0	2	0
4	F	92	0	0	1	0
4	G	75	0	0	5	0
4	H	83	0	0	10	0
4	I	83	0	0	6	2
4	J	83	0	0	7	1
4	K	51	0	0	4	1
4	L	60	0	0	6	1
All	All	8323	0	7485	107	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:4:ASN:N	4:E:201:HOH:O	1.98	0.96
1:D:66:SER:O	4:D:201:HOH:O	1.88	0.92
1:H:4:ASN:ND2	4:H:201:HOH:O	1.93	0.91
1:B:22:ASP:OD2	4:B:201:HOH:O	1.93	0.87
1:G:83:GLU:OE1	4:G:301:HOH:O	1.93	0.86

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:K:224:HOH:O	4:L:231:HOH:O[3_654]	1.88	0.32
4:A:379:HOH:O	4:I:341:HOH:O[4_675]	2.06	0.14
4:B:208:HOH:O	4:I:347:HOH:O[2_765]	2.07	0.13
4:B:274:HOH:O	4:J:260:HOH:O[2_765]	2.12	0.08

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	89/101 (88%)	86 (97%)	2 (2%)	1 (1%)	17	14
1	B	84/101 (83%)	80 (95%)	3 (4%)	1 (1%)	16	12
1	C	88/101 (87%)	86 (98%)	2 (2%)	0	100	100
1	D	87/101 (86%)	87 (100%)	0	0	100	100
1	E	83/101 (82%)	80 (96%)	3 (4%)	0	100	100
1	F	88/101 (87%)	83 (94%)	5 (6%)	0	100	100
1	G	88/101 (87%)	87 (99%)	1 (1%)	0	100	100
1	H	86/101 (85%)	84 (98%)	2 (2%)	0	100	100
1	I	85/101 (84%)	82 (96%)	3 (4%)	0	100	100
1	J	87/101 (86%)	87 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	84/101 (83%)	81 (96%)	3 (4%)	0	100	100
1	L	84/101 (83%)	82 (98%)	2 (2%)	0	100	100
All	All	1033/1212 (85%)	1005 (97%)	26 (2%)	2 (0%)	52	59

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	84	LEU
1	A	87	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/75 (87%)	64 (98%)	1 (2%)	72	84
1	B	60/75 (80%)	60 (100%)	0	100	100
1	C	64/75 (85%)	63 (98%)	1 (2%)	70	82
1	D	63/75 (84%)	63 (100%)	0	100	100
1	E	59/75 (79%)	59 (100%)	0	100	100
1	F	64/75 (85%)	64 (100%)	0	100	100
1	G	64/75 (85%)	64 (100%)	0	100	100
1	H	62/75 (83%)	62 (100%)	0	100	100
1	I	61/75 (81%)	60 (98%)	1 (2%)	70	82
1	J	63/75 (84%)	63 (100%)	0	100	100
1	K	60/75 (80%)	60 (100%)	0	100	100
1	L	60/75 (80%)	59 (98%)	1 (2%)	68	81
All	All	745/900 (83%)	741 (100%)	4 (0%)	92	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	C	12	LYS
1	I	84	LEU
1	L	4	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	75	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	91/101 (90%)	-0.37	3 (3%) 50 49	12, 20, 40, 66	0
1	B	86/101 (85%)	-0.31	1 (1%) 81 80	18, 28, 53, 66	0
1	C	90/101 (89%)	-0.33	1 (1%) 82 82	18, 27, 50, 63	0
1	D	89/101 (88%)	-0.25	1 (1%) 82 82	24, 35, 55, 62	0
1	E	85/101 (84%)	-0.24	1 (1%) 81 80	19, 33, 63, 69	0
1	F	90/101 (89%)	-0.42	0 100 100	10, 23, 43, 52	0
1	G	90/101 (89%)	-0.20	3 (3%) 50 49	15, 28, 55, 68	0
1	H	88/101 (87%)	-0.34	1 (1%) 82 82	18, 28, 51, 65	0
1	I	87/101 (86%)	-0.34	0 100 100	19, 28, 50, 65	0
1	J	89/101 (88%)	-0.31	0 100 100	25, 34, 53, 65	0
1	K	86/101 (85%)	0.06	2 (2%) 64 63	33, 49, 69, 81	0
1	L	86/101 (85%)	-0.01	3 (3%) 48 46	23, 42, 64, 81	0
All	All	1057/1212 (87%)	-0.26	16 (1%) 76 75	10, 32, 58, 81	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	90	VAL	4.3
1	A	2	SER	4.2
1	L	3	SER	4.1
1	K	85	GLY	3.5
1	E	85	GLY	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NA	A	201	1/1	0.93	0.14	0.94	45,45,45,45	0
3	CL	A	202	1/1	1.00	0.12	0.21	32,32,32,32	0
2	NA	I	201	1/1	0.96	0.06	-1.85	30,30,30,30	0
2	NA	G	201	1/1	0.98	0.08	-3.06	16,16,16,16	0
3	CL	C	201	1/1	0.98	0.16	-	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.