Full wwPDB NMR Structure Validation Report (i)

Apr 27, 2016 – 12:23 AM BST

PDB ID	:	2L7M
Title	:	Solution Structure of the Pitx2 Homeodomain R24H mutant
Authors	:	Doerdelmann, T.; Rance, M.; Kojetin, D.J.; Baird-Titus, J.M.
Deposited on	:	2010-12-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	P	68	54% 10% 35%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *minimized average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	P:8-P:51 (44)	0.10	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 14, 17, 18, 19, 20
2	4, 9, 15, 16

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1181 atoms, of which 583 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pituitary homeobox 2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	P	68	1181	370	583	120	107	1	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	-1	GLY	-	EXPRESSION TAG	UNP Q99697
P	0	SER	-	EXPRESSION TAG	UNP Q99697
P	24	HIS	ARG	ENGINEERED MUTATION	UNP Q99697
P	61	GLU	-	SEE REMARK 999	UNP Q99697
P	62	PHE	-	SEE REMARK 999	UNP Q99697
P	63	ILE	-	SEE REMARK 999	UNP Q99697
P	64	VAL	-	SEE REMARK 999	UNP Q99697
P	65	THR	-	SEE REMARK 999	UNP Q99697
P	66	ASP	-	SEE REMARK 999	UNP Q99697

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Pituitary homeobox 2

Chain P: 54% 10% 35%

G-1 S0 Q1 R2 R3 Q4 R5 T6 H7 R22 T30 R31 I34 E42 A43 R44 V45 R46 R52 R53 A54 K55 W56 R57 K58 R59 E60 E61 F62 I63 V64 T65 D66

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Pituitary homeobox 2

Chain P: 57% 7% 35%

G-1 S0 Q1 R2 R3 Q4 R5 T6 H7 R22 T30 R31 I34 R44 R52 R53 A54 K55 W56 R57 K58 R59 E60 E61 F62 I63 V64 T65 D66

4.2.2 Score per residue for model 2

- Molecule 1: Pituitary homeobox 2

Chain P: 57% 7% 35%

G-1 S0 Q1 R2 R3 Q4 R5 T6 H7 R22 T30 R31 I34 R44 R52 R53 A54 K55 W56 R57 K58 R59 E60 E61 F62 I63 V64 T65 D66

4.2.3 Score per residue for model 3

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.4 Score per residue for model 4

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.5 Score per residue for model 5

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.6 Score per residue for model 6

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.7 Score per residue for model 7

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.8 Score per residue for model 8

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.9 Score per residue for model 9

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.10 Score per residue for model 10

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.11 Score per residue for model 11

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.12 Score per residue for model 12

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.13 Score per residue for model 13

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.14 Score per residue for model 14

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.15 Score per residue for model 15

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.16 Score per residue for model 16

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.17 Score per residue for model 17

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.18 Score per residue for model 18

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: Pituitary homeobox 2

Chain P:



4.2.20 Score per residue for model 20

- Molecule 1: Pituitary homeobox 2

Chain P:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	11
CYANA	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2l7m_cs.str
Number of chemical shift lists	1
Total number of shifts	854
Number of shifts mapped to atoms	854
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	P	0.76±0.01	0±0/389 (0.0±0.0%)	1.09±0.04	3±1/529 (0.5±0.2%)
All	All	0.76	0/7780 (0.0%)	1.09	57/10580 (0.5%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	P	22	ARG	NE-CZ-NH1	9.24	124.92	120.30	2	11
1	P	31	ARG	NE-CZ-NH1	7.57	124.09	120.30	20	18
1	P	46	ARG	NE-CZ-NH1	6.53	123.56	120.30	8	11
1	P	44	ARG	NE-CZ-NH1	6.40	123.50	120.30	14	15
1	P	31	ARG	NE-CZ-NH2	-5.15	117.73	120.30	7	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	7580	7180	7180	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	44/68 (65%)	44±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	880/1360 (65%)	879 (100%)	1 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	41/63 (65%)	38±1 (92±2%)	3±1 (8±2%)	20	66
All	All	820/1260 (65%)	756 (92%)	64 (8%)	20	66

All 5 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	P	30	THR	20
1	P	34	ILE	20
1	P	42	GLU	12
1	P	39	ASN	8
1	P	19	THR	4

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: 2l7m_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	854
Number of shifts mapped to atoms	854
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	8

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	67	-0.52 ± 0.14	Should be applied
$^{13}C_{\beta}$	67	0.22 ± 0.10	None needed (< 0.5 ppm)
$^{13}C'$	65	-0.38 ± 0.08	None needed (< 0.5 ppm)
^{15}N	65	-0.32 ± 0.25	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 522 atoms were assigned a chemical shift out of a possible 593. 0 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	217/218 (100%)	87/87 (100%)	87/88 (99%)	43/43 (100%)
Sidechain	250/309 (81%)	157/181 (87%)	85/108 (79%)	8/20 (40%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	55/66 (83%)	30/35 (86%)	23/28 (82%)	2/3 (67%)
Overall	522/593 (88%)	274/303 (90%)	195/224 (87%)	53/66 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 84%, i.e. 810 atoms were assigned a chemical shift out of a possible 968. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	328/338 (97%)	131/135 (97%)	132/136 (97%)	65/67 (97%)
Sidechain	403/536 (75%)	257/317 (81%)	136/174 (78%)	10/45 (22%)
Aromatic	79/94 (84%)	42/50 (84%)	34/39 (87%)	3/5 (60%)
Overall	810/968 (84%)	430/502 (86%)	302/349 (87%)	78/117 (67%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
???	P	52	ARG	HG2	-0.80	2.92 – 0.22	-8.8
???	P	52	ARG	HB3	-0.62	3.17 – 0.37	-8.5
???	P	16	LEU	HB2	-1.09	3.32 – -0.08	-8.0
???	P	26	PRO	HG2	-0.01	3.48 – 0.38	-6.3
???	P	52	ARG	HG3	-0.23	3.00 – 0.10	-6.1
???	P	16	LEU	HD11	-0.68	2.16 – -0.64	-5.2
???	P	16	LEU	HD13	-0.68	2.16 – -0.64	-5.2
???	P	16	LEU	HD12	-0.68	2.16 – -0.64	-5.2

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain P:


RCI
0.6
0.5
0.4
0.3
0.2
0.1
0
-1:GLY
1:GLN
3:ARG
5:ARG
7:HIS
9:THR
11:GLN
13:LEU
15:GLU
17:GLU
19:THR
21:GLN
23:ASN
25:TYR
27:ASP
29:SER
31:ARG
33:GLU
35:ALA
37:TRP
39:ASN
41:THR
43:ALA
45:VAL
47:VAL
49:PHE
51:ASN
53:ARG
55:LYS
57:ARG
59:ARG
61:GLU
63:ILE
65:THR