wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 12:25 AM BST

PDB ID	:	2L7W
Title	:	Solution structure of the human Raf-1 kinase inhibitor protein
Authors	:	Yi, C.Y.
Deposited on	:	2010-12-24

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 84%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	187	48% 47% 5% •

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:3-A:187 (185)	0.42	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 11, 13, 14, 16, 17, 18, 19, 20
2	8, 9, 10, 12, 15

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2953 atoms, of which 1465 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Phosphatidylethanolamine-binding protein 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	187	2953	946	1465	257	281	4	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Phosphatidylethanolamine-binding protein 1

Chain A: 48% 47% 5% •



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

- Molecule 1: Phosphatidylethanolamine-binding protein 1

Chain A: 43% 47% 9% •

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2l7w_cs.str
Number of chemical shift lists	1
Total number of shifts	2177
Number of shifts mapped to atoms	2177
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	84%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.37±0.02	0±0/1515 (0.0±0.0%)	0.48±0.01	0±0/2059 (0.0±0.0%)
All	All	0.37	2/30300 (0.0%)	0.48	0/41180 (0.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	158	TYR	CE1-CZ	-6.78	1.29	1.38	19	1
1	A	158	TYR	CE2-CZ	6.61	1.47	1.38	19	1

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1473	1447	1441	80±6
All	All	29460	28940	28820	1601

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

5 of 405 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:54:SER:HA	1:A:58:LEU:HD13	0.95	1.37	4	20
1:A:43:PRO:HD2	1:A:174:ASP:HB3	0.94	1.38	11	4
1:A:119:ARG:HB3	1:A:170:GLN:HG3	0.93	1.41	11	5
1:A:68:LEU:HB3	1:A:86:HIS:HB3	0.93	1.36	7	16
1:A:111:PRO:HG2	1:A:172:GLU:HA	0.90	1.39	11	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/187 (98%)	157±3 (85±2%)	23±3 (12±2%)	4±2 (2±1%)	12	51
All	All	3680/3740 (98%)	3139 (85%)	458 (12%)	83 (2%)	12	51

5 of 17 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	175	ASP	20
1	A	54	SER	17
1	A	56	ASP	8
1	A	20	GLN	6
1	A	110	GLY	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/164 (99%)	146±2 (90±1%)	17±2 (10±1%)	14	58
All	All	3240/3280 (99%)	2910 (90%)	330 (10%)	14	58

5 of 56 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	39	LYS	20
1	A	47	LYS	20
1	A	53	ILE	20
1	A	113	LYS	20
1	A	28	THR	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 84% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: 2l7w_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2177
Number of shifts mapped to atoms	2177
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	15

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	186	-1.15 ± 0.13	Should be applied
$^{13}C_{\beta}$	171	-0.71 ± 0.20	Should be applied
$^{13}C'$	183	-0.75 ± 0.21	Should be applied
^{15}N	172	-0.53 ± 0.53	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 84%, i.e. 1922 atoms were assigned a chemical shift out of a possible 2293. 1 out of 35 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	891/897 (99%)	355/356 (100%)	365/370 (99%)	171/171 (100%)
Sidechain	1016/1181 (86%)	635/698 (91%)	372/430 (87%)	9/53 (17%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	15/215 (7%)	15/109 (14%)	0/89 (0%)	0/17 (0%)
Overall	1922/2293 (84%)	1005/1163 (86%)	737/889 (83%)	180/241 (75%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	170	GLN	CG	44.89	39.38 – 28.18	9.9
1	A	146	ARG	HB2	-0.86	3.15 – 0.45	-9.8
1	A	113	LYS	HB2	0.09	3.03 – 0.53	-6.8
1	A	157	LYS	HG2	-0.29	2.67 – 0.07	-6.4
1	A	46	VAL	HG11	-0.79	2.13 – -0.47	-6.2
1	A	46	VAL	HG13	-0.79	2.13 – -0.47	-6.2
1	A	46	VAL	HG12	-0.79	2.13 – -0.47	-6.2
1	A	146	ARG	HD2	1.74	4.27 – 1.97	-6.0
1	A	113	LYS	HD2	0.32	2.76 – 0.46	-5.6
1	A	14	LEU	HB2	-0.28	3.32 – -0.08	-5.6
1	A	120	TYR	HB2	0.86	4.76 – 1.06	-5.5
1	A	20	GLN	HA	1.90	6.41 – 2.11	-5.5
1	A	68	LEU	HB2	-0.13	3.32 – -0.08	-5.2
1	A	21	PRO	HD3	1.66	5.52 – 1.72	-5.1
1	A	58	LEU	HB2	-0.11	3.32 – -0.08	-5.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:


RCI
0.3
0.2
0.1
0
1:MET
6:SER
11:PRO
16:GLU
21:PRO
26:HIS
31:GLY
36:GLU
41:LEU
46:VAL
51:THR
56:ASP
61:GLY
66:LEU
71:PRO
76:ARG
81:TYR
86:HIS
91:ASN
96:ASP
101:THR
106:TYR
111:PRO
116:GLY
121:VAL
126:GLU
131:LEU
136:PRO
141:ARG
146:ARG
151:VAL
156:LYS
161:ARG
166:GLY
171:ALA
176:TYR
181:TYR
186:GLY