



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:15 AM GMT

PDB ID : 3L70  
Title : Cytochrome BC1 complex from chicken with trifloxystrobin bound  
Authors : Huang, L.; Berry, E.A.  
Deposited on : 2009-12-27  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	<div> <div>2%</div> <div>64%</div> <div>33%</div> <div>.</div> </div>
1	N	446	<div> <div>2%</div> <div>61%</div> <div>35%</div> <div>..</div> </div>
2	B	441	<div> <div>4%</div> <div>54%</div> <div>38%</div> <div>5%</div> </div>
2	O	441	<div> <div>4%</div> <div>57%</div> <div>35%</div> <div>..</div> </div>
3	C	380	<div> <div>%</div> <div>75%</div> <div>24%</div> <div>.</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	P	380	
4	D	241	
4	Q	241	
5	E	196	
5	R	196	
6	F	110	
6	S	110	
7	G	81	
7	T	81	
8	H	77	
8	U	77	
9	I	47	
9	V	47	
10	J	61	
10	W	61	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	PEE	A	2008	-	-	-	X
11	PEE	C	2007	-	-	-	X
11	PEE	E	2005	-	-	-	X
11	PEE	N	3008	-	X	-	-
11	PEE	P	3007	-	-	-	X
11	PEE	R	3005	-	-	-	X
13	JZV	C	2001	-	-	-	X
13	JZV	P	3001	-	-	-	X
14	UQ	C	2002	-	-	-	X
14	UQ	P	3002	-	-	-	X
15	CDL	C	2004	-	-	-	X
15	CDL	D	2003	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
15	CDL	Q	3003	-	-	-	X
16	GOL	C	2011	-	-	-	X
16	GOL	P	3011	-	-	-	X
18	BOG	P	2010	-	-	-	X
19	FES	E	501	-	-	X	-
19	FES	R	501	-	-	X	-

## 2 Entry composition

There are 20 unique types of molecules in this entry. The entry contains 32653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondrial ubiquinol-cytochrome-c reductase complex core protein i.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3447	2160	607	659	21			
1	N	442	Total	C	N	O	S	0	0	0
			3437	2154	605	657	21			

- Molecule 2 is a protein called Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	420	Total	C	N	O	S	0	0	0
			3133	1968	544	612	9			
2	O	422	Total	C	N	O	S	0	0	0
			3147	1977	546	614	10			

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	380	Total	C	N	O	S	0	0	0
			3017	2022	478	505	12			
3	P	379	Total	C	N	O	S	0	0	0
			3012	2019	477	504	12			

- Molecule 4 is a protein called Mitochondrial cytochrome c1, heme protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			
4	Q	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			

- Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	196	Total	C	N	O	S	0	0	0
			1513	952	263	292	6			
5	R	196	Total	C	N	O	S	0	0	0
			1509	950	263	290	6			

- Molecule 6 is a protein called Mitochondrial ubiquinol-cytochrome c reductase 14 kda protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			
6	S	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			

- Molecule 7 is a protein called Mitochondrial ubiquinol-cytochrome c reductase ubiquinone-binding protein qp-c.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	G	80	Total	C	N	O	0	0	0
			672	437	119	116			
7	T	79	Total	C	N	O	0	0	0
			662	432	117	113			

- Molecule 8 is a protein called Mitochondrial ubiquinol-cytochrome c reductase 11 kda protein, complex iii subunit viii.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	70	Total	C	N	O	S	0	0	0
			574	350	105	114	5			
8	U	67	Total	C	N	O	S	0	0	0
			553	338	103	107	5			


- Molecule 9 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	46	Total	C	N	O	S	0	0	0
			287	171	58	56	2			
9	V	43	Total	C	N	O	S	0	0	0
			277	167	55	53	2			

- Molecule 10 is a protein called Mitochondrial ubiquinol-cytochrome c reductase 7.2 kda protein.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	J	61	Total	C	N	O	0	0	0
			497	321	87	89			
10	W	60	Total	C	N	O	0	0	1
			479	311	86	82			

- Molecule 11 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOETHANOLAMINE (three-letter code: PEE) (formula:  $C_{41}H_{83}NO_8P$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	O	P	0	0
			21	12	8	1		
11	C	1	Total	C	N	O	P	0
			49	39	1	8	1	
11	E	1	Total	C	N	O	P	0
			50	40	1	8	1	
11	N	1	Total	O	P		0	0
			5	4	1			
11	P	1	Total	C	N	O	P	0
			49	39	1	8	1	
11	R	1	Total	C	N	O	P	0
			50	40	1	8	1	

- Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




- Molecule 13 is METHYL (2E)-(METHOXYIMINO)(2-{|(({(1Z)-1-[3-(TRIFLUOROMETHYL)PHENYL]ETHYLIDENE}AMINO)OXY]METHYL}PHENYL)ETHANOATE (three-letter code: JZV) (formula: C<sub>20</sub>H<sub>19</sub>F<sub>3</sub>N<sub>2</sub>O<sub>4</sub>).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	C	1	Total	C	F	N	O	0	0
			29	20	3	2	4		
13	P	1	Total	C	F	N	O	0	0
			29	20	3	2	4		

- Molecule 14 is COENZYME Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-ISOMER (three-letter code: UQ) (formula: C<sub>59</sub>H<sub>90</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	C	1	Total	C	O	0	0
			19	15	4		
14	P	1	Total	C	O	0	0
			19	15	4		

- Molecule 15 is CARDIOLIPIN (three-letter code: CDL) (formula: C<sub>81</sub>H<sub>156</sub>O<sub>17</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	C	1	Total	C	O	P	0	0
			40	21	17	2		
15	D	1	Total	C	O	P	0	0
			42	23	17	2		
15	P	1	Total	C	O	P	0	0
			40	21	17	2		
15	Q	1	Total	C	O	P	0	0
			42	23	17	2		

- Molecule 16 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	C	1	Total C O 6 3 3	0	0
16	P	1	Total C O 6 3 3	0	0

- Molecule 17 is HEME C (three-letter code: HEC) (formula:  $\text{C}_{34}\text{H}_{34}\text{FeN}_4\text{O}_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
17	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
17	Q	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 18 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
18	D	1	Total	C	O	0	0
			20	14	6		
18	D	1	Total	C	O	0	0
			13	7	6		
18	P	1	Total	C	O	0	0
			12	6	6		
18	Q	1	Total	C	O	0	0
			20	14	6		
18	Q	1	Total	C	O	0	0
			13	7	6		

- Molecule 19 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula:  $\text{Fe}_2\text{S}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
19	E	1	Total 4	Fe 2	S 2	0	0
19	R	1	Total 4	Fe 2	S 2	0	0


- Molecule 20 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
20	C	8	Total 8	O 8	0	0
20	E	1	Total 1	O 1	0	0
20	P	9	Total 9	O 9	0	0
20	R	1	Total 1	O 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein i




- Molecule 1: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein i




• Molecule 2: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2




• Molecule 2: Mitochondrial ubiquinol-cytochrome-c reductase complex core protein 2




• Molecule 3: Cytochrome b








• Molecule 3: Cytochrome b




• Molecule 4: Mitochondrial cytochrome c1, heme protein



• Molecule 4: Mitochondrial cytochrome c1, heme protein




• Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial









- Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial




- Molecule 6: Mitochondrial ubiquinol-cytochrome c reductase 14 kda protein




- Molecule 6: Mitochondrial ubiquinol-cytochrome c reductase 14 kda protein




- Molecule 7: Mitochondrial ubiquinol-cytochrome c reductase ubiquinone-binding protein qp-c




- Molecule 7: Mitochondrial ubiquinol-cytochrome c reductase ubiquinone-binding protein qp-c




- Molecule 8: Mitochondrial ubiquinol-cytochrome c reductase 11 kda protein, complex iii subunit viii




- Molecule 8: Mitochondrial ubiquinol-cytochrome c reductase 11 kda protein, complex iii subunit viii




- Molecule 9: Cytochrome b-c1 complex subunit Rieske, mitochondrial




- Molecule 9: Cytochrome b-c1 complex subunit Rieske, mitochondrial



- Molecule 10: Mitochondrial ubiquinol-cytochrome c reductase 7.2 kda protein



- Molecule 10: Mitochondrial ubiquinol-cytochrome c reductase 7.2 kda protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.61Å 181.99Å 240.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.99 – 2.75 58.26 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.3 (24.99-2.75) 98.8 (58.26-2.68)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 2.69Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.267 , 0.297 0.259 , 0.286	Depositor DCC
$R_{free}$ test set	9575 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.4	Xtriage
Anisotropy	0.603	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 54.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	2 of 204662 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	32653	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CDL, UQ, FES, HEC, HEM, PEE, JZV, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/3518	0.65	0/4767
1	N	0.42	0/3508	0.64	0/4753
2	B	0.38	0/3187	0.63	0/4321
2	O	0.40	0/3202	0.63	0/4343
3	C	0.50	0/3119	0.66	0/4270
3	P	0.45	0/3114	0.64	0/4263
4	D	0.48	0/1956	0.64	0/2658
4	Q	0.39	0/1956	0.61	0/2658
5	E	0.37	0/1547	0.59	0/2103
5	R	0.37	0/1543	0.60	0/2098
6	F	0.54	1/911 (0.1%)	0.67	0/1219
6	S	0.43	0/911	0.61	0/1219
7	G	0.49	0/694	0.69	0/941
7	T	0.43	0/684	0.64	0/929
8	H	0.44	0/582	0.63	0/779
8	U	0.31	0/561	0.56	0/751
9	I	0.39	0/218	0.60	0/293
9	V	0.36	0/218	0.59	0/293
10	J	0.43	0/508	0.60	0/682
10	W	0.40	0/490	0.60	0/660
All	All	0.43	1/32427 (0.0%)	0.63	0/44000

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	10	GLY	N-CA	5.16	1.53	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3447	0	3362	126	0
1	N	3437	0	3349	143	0
2	B	3133	0	3130	187	0
2	O	3147	0	3146	181	0
3	C	3017	0	3063	76	0
3	P	3012	0	3058	95	0
4	D	1898	0	1846	55	0
4	Q	1898	0	1846	67	0
5	E	1513	0	1478	122	0
5	R	1509	0	1474	94	0
6	F	891	0	893	20	0
6	S	891	0	893	33	0
7	G	672	0	653	32	0
7	T	662	0	645	35	0
8	H	574	0	548	16	0
8	U	553	0	535	22	0
9	I	287	0	250	37	0
9	V	277	0	251	32	0
10	J	497	0	490	16	0
10	W	479	0	478	16	0
11	A	21	0	13	0	0
11	C	49	0	72	4	0
11	E	50	0	77	1	0
11	N	5	0	0	0	0
11	P	49	0	72	3	0
11	R	50	0	77	2	0
12	C	86	0	60	9	0
12	P	86	0	60	7	0
13	C	29	0	19	4	0
13	P	29	0	19	4	0
14	C	19	0	17	4	0
14	P	19	0	17	5	0
15	C	40	0	24	4	0
15	D	42	0	28	4	0
15	P	40	0	24	4	0
15	Q	42	0	28	3	0
16	C	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	P	6	0	8	0	0
17	D	43	0	30	3	0
17	Q	43	0	30	1	0
18	D	33	0	39	0	0
18	P	12	0	11	1	0
18	Q	33	0	39	1	0
19	E	4	0	0	2	0
19	R	4	0	0	2	0
20	C	8	0	0	1	0
20	E	1	0	0	0	0
20	P	9	0	0	1	0
20	R	1	0	0	0	0
All	All	32653	0	32160	1298	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1298 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:V:35:UNK:HG3	9:V:36:UNK:H	1.05	1.14
5:E:121:GLN:HG2	5:E:170:ARG:HD3	1.15	1.13
9:I:33:UNK:HG2	9:I:73:PRO:HB3	1.34	1.07
2:O:76:THR:HG22	2:O:82:SER:H	1.16	1.07
2:B:76:THR:HG22	2:B:82:SER:H	1.18	1.07

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/446 (99%)	414 (94%)	22 (5%)	6 (1%)	14	38
1	N	440/446 (99%)	414 (94%)	19 (4%)	7 (2%)	12	34
2	B	418/441 (95%)	358 (86%)	46 (11%)	14 (3%)	5	14
2	O	420/441 (95%)	370 (88%)	40 (10%)	10 (2%)	7	22
3	C	378/380 (100%)	363 (96%)	10 (3%)	5 (1%)	15	40
3	P	377/380 (99%)	358 (95%)	14 (4%)	5 (1%)	15	40
4	D	239/241 (99%)	227 (95%)	12 (5%)	0	100	100
4	Q	239/241 (99%)	223 (93%)	15 (6%)	1 (0%)	39	72
5	E	194/196 (99%)	152 (78%)	28 (14%)	14 (7%)	1	2
5	R	194/196 (99%)	165 (85%)	22 (11%)	7 (4%)	4	12
6	F	99/110 (90%)	95 (96%)	4 (4%)	0	100	100
6	S	99/110 (90%)	91 (92%)	8 (8%)	0	100	100
7	G	78/81 (96%)	69 (88%)	6 (8%)	3 (4%)	4	11
7	T	77/81 (95%)	69 (90%)	6 (8%)	2 (3%)	7	19
8	H	68/77 (88%)	64 (94%)	4 (6%)	0	100	100
8	U	65/77 (84%)	60 (92%)	3 (5%)	2 (3%)	5	15
9	I	29/47 (62%)	27 (93%)	2 (7%)	0	100	100
9	V	29/47 (62%)	28 (97%)	1 (3%)	0	100	100
10	J	59/61 (97%)	58 (98%)	1 (2%)	0	100	100
10	W	58/61 (95%)	54 (93%)	3 (5%)	1 (2%)	11	32
All	All	4002/4160 (96%)	3659 (91%)	266 (7%)	77 (2%)	10	28

5 of 77 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	218	GLY
2	B	21	ALA
2	B	24	LEU
2	B	26	ILE
2	B	29	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/368 (99%)	354 (97%)	11 (3%)	48	80
1	N	365/368 (99%)	352 (96%)	13 (4%)	42	75
2	B	331/347 (95%)	321 (97%)	10 (3%)	48	80
2	O	333/347 (96%)	324 (97%)	9 (3%)	52	83
3	C	328/329 (100%)	320 (98%)	8 (2%)	57	86
3	P	328/329 (100%)	322 (98%)	6 (2%)	66	90
4	D	200/200 (100%)	197 (98%)	3 (2%)	72	92
4	Q	200/200 (100%)	197 (98%)	3 (2%)	72	92
5	E	166/166 (100%)	161 (97%)	5 (3%)	48	80
5	R	165/166 (99%)	161 (98%)	4 (2%)	57	86
6	F	93/96 (97%)	88 (95%)	5 (5%)	27	58
6	S	93/96 (97%)	88 (95%)	5 (5%)	27	58
7	G	71/71 (100%)	69 (97%)	2 (3%)	51	82
7	T	70/71 (99%)	69 (99%)	1 (1%)	74	92
8	H	65/71 (92%)	65 (100%)	0	100	100
8	U	63/71 (89%)	62 (98%)	1 (2%)	70	91
9	I	23/26 (88%)	20 (87%)	3 (13%)	5	14
9	V	23/26 (88%)	20 (87%)	3 (13%)	5	14
10	J	49/49 (100%)	48 (98%)	1 (2%)	63	89
10	W	47/49 (96%)	46 (98%)	1 (2%)	61	88
All	All	3378/3446 (98%)	3284 (97%)	94 (3%)	51	82

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	G	4	PHE
1	N	106	MET
6	S	64	ARG
7	G	50	PRO
10	J	59	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 82 such sidechains are listed below:



Mol	Chain	Res	Type
7	G	44	GLN
1	N	267	ASN
5	R	107	ASN
7	G	73	ASN
1	N	85	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
11	PEE	A	2008	-	20,20,50	1.84	6 (30%)	21,25,55	0.68	0
13	JZV	C	2001	-	30,30,30	1.90	6 (20%)	37,41,41	3.37	10 (27%)
14	UQ	C	2002	-	19,19,63	2.51	11 (57%)	23,26,79	1.30	3 (13%)
15	CDL	C	2004	-	39,39,99	1.18	2 (5%)	41,51,111	1.14	4 (9%)
11	PEE	C	2007	-	48,48,50	1.29	7 (14%)	49,53,55	0.95	4 (8%)
16	GOL	C	2011	-	5,5,5	1.24	0	5,5,5	0.65	0
12	HEM	C	501	3	30,50,50	3.15	9 (30%)	24,82,82	2.31	10 (41%)
12	HEM	C	502	3	30,50,50	2.57	7 (23%)	24,82,82	2.28	7 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
15	CDL	D	2003	-	41,41,99	1.18	3 (7%)	43,53,111	1.07	2 (4%)
18	BOG	D	2009	-	20,20,20	0.92	1 (5%)	25,25,25	0.88	2 (8%)
18	BOG	D	2091	-	13,13,20	1.28	2 (15%)	18,18,25	1.05	2 (11%)
17	HEC	D	501	4	24,50,50	1.75	2 (8%)	19,82,82	3.20	6 (31%)
11	PEE	E	2005	-	49,49,50	1.46	9 (18%)	50,54,55	0.94	5 (10%)
19	FES	E	501	5	0,4,4	0.00	-	0,4,4	0.00	-
11	PEE	N	3008	-	4,4,50	3.75	4 (100%)	6,6,55	0.58	0
18	BOG	P	2010	-	12,12,20	1.44	3 (25%)	17,17,25	0.61	0
13	JZV	P	3001	-	30,30,30	1.95	6 (20%)	37,41,41	3.50	11 (29%)
14	UQ	P	3002	-	19,19,63	2.49	11 (57%)	23,26,79	1.28	3 (13%)
15	CDL	P	3004	-	39,39,99	1.18	2 (5%)	41,51,111	1.16	5 (12%)
11	PEE	P	3007	-	48,48,50	1.27	7 (14%)	49,53,55	0.92	4 (8%)
16	GOL	P	3011	-	5,5,5	1.34	0	5,5,5	0.69	0
12	HEM	P	501	3	30,50,50	3.16	9 (30%)	24,82,82	2.13	7 (29%)
12	HEM	P	502	3	30,50,50	2.60	10 (33%)	24,82,82	2.30	7 (29%)
15	CDL	Q	3003	-	41,41,99	1.18	2 (4%)	43,53,111	1.08	3 (6%)
18	BOG	Q	3009	-	20,20,20	0.96	1 (5%)	25,25,25	0.92	2 (8%)
18	BOG	Q	3091	-	13,13,20	1.37	2 (15%)	18,18,25	1.08	2 (11%)
17	HEC	Q	501	4	24,50,50	2.27	2 (8%)	19,82,82	3.33	6 (31%)
11	PEE	R	3005	-	49,49,50	1.44	10 (20%)	50,54,55	0.94	5 (10%)
19	FES	R	501	5	0,4,4	0.00	-	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	PEE	A	2008	-	-	0/24/24/54	0/0/0/0
13	JZV	C	2001	-	-	0/29/29/29	0/2/2/2
14	UQ	C	2002	-	-	0/11/35/87	0/1/1/1
15	CDL	C	2004	-	-	0/49/49/110	0/0/0/0
11	PEE	C	2007	-	-	0/52/52/54	0/0/0/0
16	GOL	C	2011	-	-	0/4/4/4	0/0/0/0
12	HEM	C	501	3	-	0/10/54/54	0/0/8/8
12	HEM	C	502	3	-	0/10/54/54	0/0/8/8
15	CDL	D	2003	-	-	0/51/51/110	0/0/0/0
18	BOG	D	2009	-	-	0/11/31/31	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
18	BOG	D	2091	-	-	0/4/24/31	0/1/1/1
17	HEC	D	501	4	-	0/6/54/54	0/0/8/8
11	PEE	E	2005	-	-	0/53/53/54	0/0/0/0
19	FES	E	501	5	-	0/0/4/4	0/1/1/1
11	PEE	N	3008	-	-	0/0/0/54	0/0/0/0
18	BOG	P	2010	-	-	0/2/22/31	0/1/1/1
13	JZV	P	3001	-	-	0/29/29/29	0/2/2/2
14	UQ	P	3002	-	-	0/11/35/87	0/1/1/1
15	CDL	P	3004	-	-	0/49/49/110	0/0/0/0
11	PEE	P	3007	-	-	0/52/52/54	0/0/0/0
16	GOL	P	3011	-	-	0/4/4/4	0/0/0/0
12	HEM	P	501	3	-	0/10/54/54	0/0/8/8
12	HEM	P	502	3	-	0/10/54/54	0/0/8/8
15	CDL	Q	3003	-	-	0/51/51/110	0/0/0/0
18	BOG	Q	3009	-	-	0/11/31/31	0/1/1/1
18	BOG	Q	3091	-	-	0/4/24/31	0/1/1/1
17	HEC	Q	501	4	-	0/6/54/54	0/0/8/8
11	PEE	R	3005	-	-	0/53/53/54	0/0/0/0
19	FES	R	501	5	-	0/0/4/4	0/1/1/1

The worst 5 of 134 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	P	501	HEM	C3B-C4B	-9.86	1.43	1.51
12	C	501	HEM	C3B-C4B	-8.71	1.44	1.51
17	Q	501	HEC	C3B-C2B	-7.38	1.33	1.40
17	Q	501	HEC	C3C-C2C	-7.29	1.33	1.40
12	P	501	HEM	C3B-CAB	-7.03	1.38	1.51

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
17	Q	501	HEC	CBB-CAB-C3B	-8.52	108.43	127.35
17	D	501	HEC	CBB-CAB-C3B	-8.51	108.44	127.35
13	C	2001	JZV	CAY-CAS-NAR	-7.63	108.61	123.72
13	P	3001	JZV	CAY-CAS-NAR	-7.20	109.46	123.72
17	Q	501	HEC	CBC-CAC-C3C	-6.99	111.83	127.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

22 monomers are involved in 66 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	C	2001	JZV	4	0
14	C	2002	UQ	4	0
15	C	2004	CDL	4	0
11	C	2007	PEE	4	0
12	C	501	HEM	5	0
12	C	502	HEM	4	0
15	D	2003	CDL	4	0
17	D	501	HEC	3	0
11	E	2005	PEE	1	0
19	E	501	FES	2	0
18	P	2010	BOG	1	0
13	P	3001	JZV	4	0
14	P	3002	UQ	5	0
15	P	3004	CDL	4	0
11	P	3007	PEE	3	0
12	P	501	HEM	4	0
12	P	502	HEM	3	0
15	Q	3003	CDL	3	0
18	Q	3091	BOG	1	0
17	Q	501	HEC	1	0
11	R	3005	PEE	2	0
19	R	501	FES	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	444/446 (99%)	0.07	11 (2%) 61 54	32, 58, 86, 100	0
1	N	442/446 (99%)	0.17	7 (1%) 74 70	41, 68, 93, 100	0
2	B	420/441 (95%)	0.28	17 (4%) 42 35	48, 76, 111, 133	0
2	O	422/441 (95%)	0.38	19 (4%) 37 30	39, 74, 103, 122	0
3	C	380/380 (100%)	0.07	5 (1%) 79 75	21, 39, 83, 123	0
3	P	379/380 (99%)	0.22	10 (2%) 59 53	29, 58, 88, 125	0
4	D	241/241 (100%)	-0.06	3 (1%) 81 76	29, 41, 81, 103	0
4	Q	241/241 (100%)	0.21	13 (5%) 29 22	46, 67, 97, 120	0
5	E	196/196 (100%)	1.76	77 (39%) 0 0	34, 135, 167, 175	0
5	R	196/196 (100%)	0.62	25 (12%) 5 3	40, 86, 132, 147	0
6	F	101/110 (91%)	-0.30	0 100 100	26, 43, 61, 94	0
6	S	101/110 (91%)	0.04	4 (3%) 42 35	50, 65, 102, 121	0
7	G	80/81 (98%)	0.14	2 (2%) 61 54	33, 52, 97, 111	0
7	T	79/81 (97%)	0.97	13 (16%) 2 1	46, 78, 139, 149	0
8	H	70/77 (90%)	-0.01	2 (2%) 55 48	34, 58, 81, 119	0
8	U	67/77 (87%)	1.03	13 (19%) 1 1	82, 108, 128, 130	0
9	I	31/47 (65%)	2.21	14 (45%) 0 0	72, 107, 133, 134	0
9	V	31/47 (65%)	2.24	17 (54%) 0 0	66, 107, 132, 136	0
10	J	61/61 (100%)	0.26	4 (6%) 22 16	43, 54, 95, 139	0
10	W	60/61 (98%)	0.57	3 (5%) 32 25	55, 68, 99, 107	0
All	All	4042/4160 (97%)	0.33	259 (6%) 23 17	21, 64, 121, 175	0

The worst 5 of 259 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
9	V	63	ASP	11.4
5	E	98	VAL	10.2
5	E	188	VAL	8.6
7	T	2	ILE	7.3
5	E	84	GLY	7.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
16	GOL	P	3011	6/6	0.80	0.40	7.95	73,75,76,77	0
11	PEE	C	2007	49/51	0.85	0.30	7.15	43,58,70,73	0
14	UQ	P	3002	19/63	0.76	0.38	5.30	119,124,125,126	0
18	BOG	P	2010	12/20	0.46	0.35	5.20	135,137,138,138	0
16	GOL	C	2011	6/6	0.85	0.34	4.92	67,70,72,72	0
11	PEE	R	3005	50/51	0.71	0.42	4.41	80,98,108,110	0
11	PEE	E	2005	50/51	0.82	0.35	4.34	68,89,95,97	0
15	CDL	Q	3003	42/100	0.72	0.29	3.78	109,121,139,139	0
11	PEE	A	2008	21/51	0.54	0.39	3.58	129,133,134,135	0
14	UQ	C	2002	19/63	0.84	0.30	3.54	88,90,90,91	0
13	JZV	C	2001	29/29	0.93	0.23	2.82	38,46,86,87	0
11	PEE	P	3007	49/51	0.83	0.33	2.56	53,79,93,94	0
15	CDL	D	2003	42/100	0.81	0.29	2.49	82,92,106,108	0
13	JZV	P	3001	29/29	0.89	0.24	2.31	59,66,104,106	0
15	CDL	C	2004	40/100	0.91	0.22	2.26	66,80,91,93	0
18	BOG	Q	3009	20/20	0.89	0.23	1.03	62,75,77,78	0
12	HEM	P	501	43/43	0.97	0.22	1.02	33,40,48,54	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
12	HEM	C	501	43/43	0.98	0.21	0.51	23,30,37,42	0
15	CDL	P	3004	40/100	0.86	0.21	0.49	96,102,111,111	0
18	BOG	D	2009	20/20	0.93	0.17	0.40	53,59,62,64	0
17	HEC	D	501	43/43	0.98	0.17	0.24	21,29,34,35	0
12	HEM	P	502	43/43	0.98	0.17	0.16	29,41,54,59	0
17	HEC	Q	501	43/43	0.96	0.18	0.12	52,56,64,65	0
12	HEM	C	502	43/43	0.98	0.17	-0.19	24,27,35,41	0
19	FES	E	501	4/4	0.93	0.11	-1.98	144,144,145,145	0
19	FES	R	501	4/4	0.98	0.07	-2.76	74,76,77,78	0
18	BOG	D	2091	13/20	0.27	0.61	-	200,201,201,201	0
18	BOG	Q	3091	13/20	0.23	0.61	-	183,186,187,187	0
11	PEE	N	3008	5/51	0.83	0.20	-	110,111,112,112	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.