



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 12:11 AM BST

PDB ID : 2L8W  
Title : r(CCGCUGCGG)2 UU Internal Loop Found in Myotonic Dystrophy Type 1  
- UU pair with two hydrogen bond pairs  
Authors : Parkesh, R.; Fountain, M.; Disney, M.  
Deposited on : 2011-01-26

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*



The overall completeness of chemical shifts assignment is 14%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric       | Whole archive<br>(#Entries) | NMR archive<br>(#Entries) |
|--------------|-----------------------------|---------------------------|
| Clashscore   | 114402                      | 11133                     |
| RNA backbone | 3027                        | 600                       |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

| Mol | Chain | Length | Quality of chain  |
|-----|-------|--------|---|
| 1   | A     | 9      | <br>56%                      44%                          |
| 1   | B     | 9      | <br>33%                      56%                      11% |

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 578 atoms, of which 200 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(\*CP\*CP\*GP\*CP\*UP\*GP\*CP\*GP\*G)-3').

| Mol | Chain | Residues | Atoms |    |     |    |    |   | Trace |
|-----|-------|----------|-------|----|-----|----|----|---|-------|
| 1   | A     | 9        | Total | C  | H   | N  | O  | P | 0     |
|     |       |          | 289   | 85 | 100 | 34 | 62 | 8 |       |
| 1   | B     | 9        | Total | C  | H   | N  | O  | P | 0     |
|     |       |          | 289   | 85 | 100 | 34 | 62 | 8 |       |

## 4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (5'-R(\*CP\*CP\*GP\*CP\*UP\*GP\*CP\*GP\*G)-3')

Chain A: 



- Molecule 1: RNA (5'-R(\*CP\*CP\*GP\*CP\*UP\*GP\*CP\*GP\*G)-3')

Chain B: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 25 calculated structures, 1 were deposited, based on the following criterion: *structures with acceptable covalent geometry*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification        | Version |
|---------------|-----------------------|---------|
| CYANA         | geometry optimization | 2.1     |
| AMBER         | refinement            | 11      |

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

|  |             |
|--|-------------|
| Chemical shift file(s)                       | 2l8w_cs.str |
| Number of chemical shift lists               | 1           |
| Total number of shifts                       | 48          |
| Number of shifts mapped to atoms             | 48          |
| Number of unparsed shifts                    | 0           |
| Number of shifts with mapping errors         | 0           |
| Number of shifts with mapping warnings       | 0           |
| Assignment completeness (well-defined parts) | 14%         |

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |                | Bond angles |                 |
|-----|-------|--------------|----------------|-------------|-----------------|
|     |       | RMSZ         | #Z>5           | RMSZ        | #Z>5            |
| 1   | A     | 3.17         | 29/210 (13.8%) | 3.55        | 48/326 (14.7%)  |
| 1   | B     | 3.23         | 26/210 (12.4%) | 3.58        | 53/326 (16.3%)  |
| All | All   | 3.20         | 55/420 (13.1%) | 3.56        | 101/652 (15.5%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | Chirality | Planarity |
|-----|-------|-----------|-----------|
| 1   | A     | 0         | 4         |
| 1   | B     | 0         | 6         |
| All | All   | 0         | 10        |

5 of 55 bond outliers are listed below. They are sorted according to the Z-score.

| Mol | Chain | Res | Type | Atoms   | Z      | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|--------|-------------|----------|
| 1   | B     | 12  | G    | N9-C8   | 11.62  | 1.46        | 1.37     |
| 1   | A     | 9   | G    | C2-N2   | -11.16 | 1.23        | 1.34     |
| 1   | A     | 9   | G    | N3-C4   | 10.06  | 1.42        | 1.35     |
| 1   | B     | 11  | C    | C4-N4   | -9.53  | 1.25        | 1.33     |
| 1   | B     | 10  | C    | C4'-O4' | -8.95  | 1.33        | 1.45     |

5 of 101 angle outliers are listed below. They are sorted according to the Z-score.

| Mol | Chain | Res | Type | Atoms      | Z      | Observed(°) | Ideal(°) |
|-----|-------|-----|------|------------|--------|-------------|----------|
| 1   | B     | 16  | C    | O4'-C1'-N1 | 13.79  | 119.23      | 108.20   |
| 1   | A     | 5   | U    | N1-C2-O2   | 12.50  | 131.55      | 122.80   |
| 1   | A     | 5   | U    | N3-C2-O2   | -12.49 | 113.46      | 122.20   |
| 1   | B     | 16  | C    | N3-C4-C5   | -11.92 | 117.13      | 121.90   |
| 1   | A     | 8   | G    | O4'-C1'-N9 | 11.89  | 117.71      | 108.20   |

There are no chirality outliers.

5 of 10 planar outliers are listed below.

| Mol | Chain | Res | Type | Group     |
|-----|-------|-----|------|-----------|
| 1   | B     | 18  | G    | Sidechain |
| 1   | B     | 11  | C    | Sidechain |
| 1   | B     | 13  | C    | Sidechain |
| 1   | B     | 15  | G    | Sidechain |
| 1   | A     | 8   | G    | Sidechain |

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| All | All   | 378   | 200      | 185      | -       |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

| Mol | Chain | Analysed    | Backbone Outliers | Pucker Outliers | Suiteness |
|-----|-------|-------------|-------------------|-----------------|-----------|
| 1   | A     | 8/9 (89%)   | 0 (0%)            | 0 (0%)          | 0.42      |
| 1   | B     | 8/9 (89%)   | 0 (0%)            | 1 (12%)         | 0.33      |
| All | All   | 16/18 (89%) | 0 (0%)            | 1 (6%)          | 0.38      |

The overall RNA backbone suiteness is 0.38.



There are no RNA backbone outliers to report.

All RNA pucker outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | B     | 11  | C    |

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 14% for the well-defined parts and 14% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2l8w\_cs.str

Chemical shift list name: *data.str*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

|   |    |
|---|----|
| Total number of shifts                  | 48 |
| Number of shifts mapped to atoms        | 48 |
| Number of unparsed shifts               | 0  |
| Number of shifts with mapping errors    | 0  |
| Number of shifts with mapping warnings  | 0  |
| Number of shift outliers (ShiftChecker) | 0  |

#### 7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 14%, i.e. 47 atoms were assigned a chemical shift out of a possible 346. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

|           | Total        | <sup>1</sup> H | <sup>13</sup> C | <sup>15</sup> N |
|-----------|--------------|----------------|-----------------|-----------------|
| Backbone  | 0/0 (—%)     | 0/0 (—%)       | 0/0 (—%)        | 0/0 (—%)        |
| Sidechain | 0/0 (—%)     | 0/0 (—%)       | 0/0 (—%)        | 0/0 (—%)        |
| Aromatic  | 0/0 (—%)     | 0/0 (—%)       | 0/0 (—%)        | 0/0 (—%)        |
| Overall   | 47/346 (14%) | 47/202 (23%)   | 0/118 (0%)      | 0/26 (0%)       |

#### 7.1.4 Statistically unusual chemical shifts

There are no statistically unusual chemical shifts.

### 7.1.5 Random Coil Index (RCI) plots

No *random coil index* (RCI) plot could be generated from the current chemical shift list (data.str). RCI is only applicable to proteins.