



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:19 PM GMT

PDB ID : 4LAD  
Title : Crystal Structure of the Ube2g2:RING-G2BR complex  
Authors : Liang, Y.-H.; Li, J.; Das, R.; Byrd, R.A.; Ji, X.  
Deposited on : 2013-06-19  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	
2	B	150	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-conjugating enzyme E2 G2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	1	0
			1231	792	199	230	10			

- Molecule 2 is a protein called E3 ubiquitin-protein ligase AMFR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	69	Total	C	N	O	S	0	0	0
			560	339	110	102	9			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	311	HIS	-	EXPRESSION TAG	UNP Q9UKV5
B	312	MET	-	EXPRESSION TAG	UNP Q9UKV5
B	534	GLY	-	LINKER	UNP Q9UKV5
B	535	GLY	-	LINKER	UNP Q9UKV5
B	536	GLY	-	LINKER	UNP Q9UKV5
B	537	GLY	-	LINKER	UNP Q9UKV5
B	538	GLY	-	LINKER	UNP Q9UKV5
B	539	GLY	-	LINKER	UNP Q9UKV5
B	540	GLY	-	LINKER	UNP Q9UKV5
B	541	SER	-	LINKER	UNP Q9UKV5
B	542	SER	-	LINKER	UNP Q9UKV5
B	543	GLY	-	LINKER	UNP Q9UKV5
B	544	SER	-	LINKER	UNP Q9UKV5
B	545	SER	-	LINKER	UNP Q9UKV5
B	546	GLY	-	LINKER	UNP Q9UKV5
B	547	GLY	-	LINKER	UNP Q9UKV5
B	548	SER	-	LINKER	UNP Q9UKV5
B	549	GLY	-	LINKER	UNP Q9UKV5
B	550	GLY	-	LINKER	UNP Q9UKV5
B	551	GLY	-	LINKER	UNP Q9UKV5

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	552	SER	-	LINKER	UNP Q9UKV5
B	553	GLY	-	LINKER	UNP Q9UKV5
B	554	SER	-	LINKER	UNP Q9UKV5
B	555	SER	-	LINKER	UNP Q9UKV5
B	556	SER	-	LINKER	UNP Q9UKV5
B	557	GLY	-	LINKER	UNP Q9UKV5
B	558	GLY	-	LINKER	UNP Q9UKV5
B	559	GLY	-	LINKER	UNP Q9UKV5
B	560	GLY	-	LINKER	UNP Q9UKV5
B	561	GLY	-	LINKER	UNP Q9UKV5
B	562	SER	-	LINKER	UNP Q9UKV5
B	563	GLY	-	LINKER	UNP Q9UKV5
B	564	GLY	-	LINKER	UNP Q9UKV5
B	565	GLY	-	LINKER	UNP Q9UKV5
B	566	SER	-	LINKER	UNP Q9UKV5
B	567	GLY	-	LINKER	UNP Q9UKV5
B	568	GLY	-	LINKER	UNP Q9UKV5
B	569	GLY	-	LINKER	UNP Q9UKV5
B	570	GLY	-	LINKER	UNP Q9UKV5
B	571	GLY	-	LINKER	UNP Q9UKV5
B	572	GLY	-	LINKER	UNP Q9UKV5
B	573	GLY	-	LINKER	UNP Q9UKV5

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Zn 2 2	0	0

- Molecule 4 is OXALATE ION (three-letter code: OXL) (formula: C<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 2 4	0	0
4	B	1	Total C O 6 2 4	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	26	Total O 26 26	0	0
5	B	13	Total O 13 13	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.25Å 58.25Å 158.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.12 – 2.30 39.12 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.0 (39.12-2.30) 97.0 (39.12-2.30)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.226 , 0.281 0.231 , 0.283	Depositor DCC
$R_{free}$ test set	998 reflections (8.70%)	DCC
Wilson B-factor (Å <sup>2</sup> )	56.1	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 12495 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1844	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/1264	0.63	0/1713
2	B	0.40	0/568	0.57	0/759
All	All	0.47	0/1832	0.61	0/2472

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1231	0	1204	43	0
2	B	560	0	540	17	0
3	B	2	0	0	0	0
4	B	12	0	0	0	0
5	A	26	0	0	0	0
5	B	13	0	0	1	0
All	All	1844	0	1744	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:TYR:CE2	1:A:89:CYS:SG	2.46	1.06
1:A:83:TYR:HE2	1:A:89:CYS:SG	1.79	1.05
1:A:70:LYS:HB3	1:A:87:ARG:HH12	1.37	0.89
1:A:83:TYR:HE2	1:A:89:CYS:HG	0.87	0.87
1:A:66:LEU:HD21	2:B:367:SER:HB3	1.59	0.84
1:A:45:GLU:CD	1:A:45:GLU:H	1.90	0.74
1:A:51:PHE:HE1	2:B:596:ARG:HH11	1.33	0.74
1:A:51:PHE:HE1	2:B:596:ARG:NH1	1.91	0.69
1:A:92:ILE:HD11	1:A:114:GLN:HG2	1.76	0.68
1:A:83:TYR:CE1	1:A:136:ALA:HA	2.28	0.67
1:A:51:PHE:CE1	2:B:596:ARG:NH1	2.64	0.66
1:A:133:GLU:HG3	1:A:133:GLU:O	1.96	0.65
1:A:45:GLU:HA	1:A:50:GLU:HG3	1.79	0.65
2:B:594:ARG:HD2	5:B:809:HOH:O	1.97	0.64
1:A:81:ASN:HD21	1:A:131:ASN:HB2	1.66	0.61
1:A:24:ILE:HD11	1:A:124:VAL:HG21	1.84	0.59
1:A:58:LEU:N	1:A:58:LEU:HD12	2.18	0.58
2:B:368:TRP:CD1	2:B:376:PRO:HG3	2.38	0.57
1:A:143:MET:HG2	1:A:151:PHE:HB2	1.86	0.57
1:A:111:SER:OG	1:A:113:VAL:HG12	2.08	0.54
1:A:114:GLN:N	1:A:114:GLN:HE21	2.05	0.54
2:B:378:CYS:O	2:B:379:ARG:HG2	2.08	0.54
2:B:365:LEU:HD22	2:B:369:LEU:HG	1.88	0.53
1:A:70:LYS:HB3	1:A:87:ARG:NH1	2.14	0.53
1:A:148:ARG:HG3	1:A:152:TYR:CE2	2.43	0.52
1:A:133:GLU:CG	1:A:133:GLU:O	2.57	0.52
1:A:79:HIS:ND1	1:A:80:PRO:HD2	2.25	0.52
2:B:372:ASP:OD1	2:B:374:SER:HB2	2.10	0.52
1:A:47:THR:O	1:A:50:GLU:HB2	2.10	0.51
1:A:112:PRO:HD3	2:B:368:TRP:CZ3	2.47	0.50
2:B:341:CYS:O	2:B:344:CYS:O	2.29	0.49
1:A:144:TRP:O	1:A:148:ARG:NE	2.38	0.49
1:A:12:GLU:O	1:A:16:LEU:HB2	2.13	0.49
1:A:50:GLU:HG2	1:A:51:PHE:CD2	2.49	0.47
2:B:352:ARG:NH2	2:B:366:ARG:HE	2.12	0.47
1:A:41:ILE:N	1:A:41:ILE:HD12	2.31	0.46
1:A:146:ASP:OD1	1:A:146:ASP:N	2.47	0.46
2:B:354:LEU:HB3	2:B:355:PRO:HD2	1.98	0.45
1:A:60:PHE:CD1	1:A:69:PRO:HB3	2.52	0.45
1:A:138:VAL:O	1:A:142:LYS:HG3	2.16	0.45
1:A:50:GLU:HG2	1:A:51:PHE:CE2	2.53	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:GLY:C	1:A:24:ILE:HD12	2.37	0.44
2:B:352:ARG:NH2	2:B:366:ARG:HH21	2.15	0.44
1:A:87:ARG:HH11	1:A:87:ARG:HG3	1.82	0.44
1:A:56:ALA:HB2	1:A:73:PHE:CD1	2.52	0.43
1:A:131:ASN:N	1:A:131:ASN:HD22	2.17	0.43
1:A:51:PHE:CE1	2:B:596:ARG:HB3	2.54	0.42
1:A:110:TRP:CZ3	1:A:114:GLN:HB3	2.54	0.42
1:A:79:HIS:HA	1:A:80:PRO:HD3	1.86	0.42
1:A:92:ILE:HD11	1:A:114:GLN:CG	2.47	0.42
2:B:354:LEU:N	2:B:354:LEU:HD12	2.35	0.41
2:B:355:PRO:O	2:B:357:GLY:N	2.54	0.41
1:A:83:TYR:HD2	1:A:87:ARG:HB3	1.86	0.41
1:A:64:TYR:CD1	1:A:65:PRO:HA	2.55	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/165 (92%)	139 (92%)	10 (7%)	2 (1%)	15	15
2	B	65/150 (43%)	55 (85%)	8 (12%)	2 (3%)	5	3
All	All	216/315 (69%)	194 (90%)	18 (8%)	4 (2%)	10	8

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	107	ALA
2	B	356	CYS
1	A	109	ARG
2	B	343	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/143 (95%)	125 (92%)	11 (8%)	15	18
2	B	62/109 (57%)	52 (84%)	10 (16%)	3	3
All	All	198/252 (79%)	177 (89%)	21 (11%)	8	9

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LEU
1	A	18	LEU
1	A	66	LEU
1	A	74	THR
1	A	113	VAL
1	A	114	GLN
1	A	126	MET
1	A	132	ASP
1	A	133	GLU
1	A	138	VAL
1	A	146	ASP
2	B	339	ASP
2	B	348	MET
2	B	352	ARG
2	B	359	LEU
2	B	360	PHE
2	B	365	LEU
2	B	370	GLU
2	B	373	THR
2	B	579	GLN
2	B	592	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	131	ASN
1	A	150	GLN
2	B	592	GLN
2	B	599	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	OXL	B	703	-	0,5,5	0.00	-	0,6,6	0.00	-
4	OXL	B	704	-	0,5,5	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OXL	B	703	-	-	0/0/4/4	0/0/0/0
4	OXL	B	704	-	-	0/0/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	154/165 (93%)	0.61	12 (7%) 16 22	39, 53, 80, 99	0
2	B	69/150 (46%)	0.91	8 (11%) 6 10	45, 73, 95, 98	0
All	All	223/315 (70%)	0.70	20 (8%) 12 17	39, 57, 89, 99	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	354	LEU	4.0
2	B	359	LEU	3.8
2	B	357	GLY	3.5
1	A	133	GLU	3.3
1	A	132	ASP	3.2
2	B	351	ALA	3.1
2	B	379	ARG	3.1
1	A	152	TYR	3.0
1	A	108	GLU	3.0
1	A	144	TRP	2.9
1	A	134	SER	2.9
2	B	573	GLY	2.9
1	A	138	VAL	2.8
2	B	347	SER	2.6
2	B	369	LEU	2.6
1	A	96	PRO	2.6
1	A	87	ARG	2.3
1	A	94	HIS	2.2
1	A	128	ALA	2.1
1	A	116	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	B	701	1/1	0.96	0.09	-	61,61,61,61	0
3	ZN	B	702	1/1	0.96	0.09	-	82,82,82,82	0
4	OXL	B	704	6/6	0.83	0.21	-	78,85,89,90	0
4	OXL	B	703	6/6	0.73	0.16	-	75,76,76,77	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.