



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2017 – 01:01 PM EST

PDB ID : 5LB3  
Title : Crystal structure of human RECQL5 helicase in complex with ADP/Mg.  
Authors : Newman, J.A.; Aitkenhead, H.; Savitsky, P.; Krojer, T.; Von Delft, F.; Arrow-smith, C.H.; Edwards, A.M.; Bountra, C.; Gileadi, O.; Structural Genomics Consortium (SGC)  
Deposited on : 2016-06-15  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	445	<div> <div>8%</div> <div>89%</div> <div>9%</div> </div>
1	E	445	<div> <div>7%</div> <div>91%</div> <div>7%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent DNA helicase Q5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	439	Total	C	N	O	S	0	7	0
			3461	2181	619	637	24			
1	E	439	Total	C	N	O	S	0	3	0
			3422	2161	607	630	24			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	MET	-	initiating methionine	UNP O94762
B	10	SER	-	expression tag	UNP O94762
E	9	MET	-	initiating methionine	UNP O94762
E	10	SER	-	expression tag	UNP O94762

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	E	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	E	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	E	1	Total Mg 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	522	Total O 522 522	0	0
5	E	469	Total O 469 469	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent DNA helicase Q5



- Molecule 1: ATP-dependent DNA helicase Q5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.76Å 62.94Å 104.70Å 90.00° 95.44° 90.00°	Depositor
Resolution (Å)	19.88 – 1.80 19.88 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.88-1.80) 99.8 (19.88-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.80Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1682)	Depositor
R, $R_{free}$	0.195 , 0.234 0.192 , 0.231	Depositor DCC
$R_{free}$ test set	4294 reflections (4.92%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtrriage
Anisotropy	0.329	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7932	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.52 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.8917e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.30	0/3527	0.43	1/4768 (0.0%)
1	E	0.31	0/3487	0.41	0/4714
All	All	0.30	0/7014	0.42	1/9482 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	392	ASP	CB-CG-OD1	-5.81	113.07	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3461	0	3460	33	0
1	E	3422	0	3422	18	0
2	B	1	0	0	0	0
2	E	1	0	0	0	0
3	B	27	0	12	0	0
3	E	27	0	12	2	0
4	B	1	0	0	0	0
4	E	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	522	0	0	11	4
5	E	469	0	0	8	3
All	All	7932	0	6906	53	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:272[B]:CYS:SG	1:B:316[B]:THR:HG23	1.78	1.24
1:B:272[B]:CYS:SG	1:B:316[B]:THR:CG2	2.51	0.98
1:B:201:GLU:OE1	5:B:601:HOH:O	2.02	0.76
1:B:152:SER:OG	5:B:602:HOH:O	2.04	0.75
1:B:395:THR:OG1	5:B:603:HOH:O	2.07	0.71
1:B:449:ARG:NH2	5:B:608:HOH:O	2.25	0.69
1:E:348:GLY:O	1:E:352:ARG:NH1	2.25	0.69
1:B:392:ASP:HA	5:B:603:HOH:O	1.91	0.69
1:B:397:MET:SD	5:B:1108:HOH:O	2.52	0.68
1:E:390:ALA:O	5:E:601:HOH:O	2.12	0.67
1:B:316[A]:THR:CG2	1:B:317:ILE:HG13	2.25	0.66
1:B:72:ILE:HG12	1:B:123:THR:O	1.96	0.65
1:B:270[A]:GLU:OE1	5:B:604:HOH:O	2.15	0.65
1:E:270:GLU:OE1	5:E:602:HOH:O	2.14	0.64
3:E:502:ADP:O3'	5:E:603:HOH:O	2.15	0.64
1:E:429:LYS:NZ	5:E:611:HOH:O	2.32	0.62
1:E:442:ARG:NH1	5:E:612:HOH:O	2.32	0.61
1:B:408:GLU:OE1	5:B:605:HOH:O	2.16	0.61
1:B:146:VAL:HG22	1:B:183:LEU:HD21	1.81	0.60
1:E:96:ARG:NH1	1:E:122:GLN:OE1	2.35	0.59
1:B:316[A]:THR:HG23	1:B:317:ILE:HG13	1.85	0.59
1:B:142:LEU:HD21	1:B:179:LEU:HD13	1.86	0.57
1:E:116:LEU:O	1:E:148:ARG:NH2	2.37	0.55
1:E:265:TYR:OH	1:E:349:ARG:NH1	2.41	0.54
1:B:289:TYR:CE1	1:B:298:ARG:HG3	2.44	0.52
1:E:162:VAL:HG12	1:E:173:TYR:HB3	1.91	0.51
1:B:316[A]:THR:HG21	5:B:883:HOH:O	2.10	0.51
1:B:180:ARG:NH2	1:B:183:LEU:O	2.38	0.51
3:E:502:ADP:O2'	5:E:606:HOH:O	2.19	0.51
1:E:243:LYS:HD3	1:E:282:ARG:HB3	1.93	0.51
1:B:10:SER:OG	1:B:11:ASP:N	2.43	0.51

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:316[A]:THR:HG22	1:B:317:ILE:HG13	1.92	0.50
1:B:331:VAL:HG23	1:B:350:ALA:HB2	1.94	0.50
1:B:275:LEU:HD11	1:B:334:TRP:CD1	2.47	0.49
1:B:441[A]:ARG:NH1	1:B:445:GLU:OE2	2.44	0.49
1:E:185:HIS:HB2	5:E:929:HOH:O	2.14	0.48
1:E:386:ARG:NE	5:E:605:HOH:O	2.18	0.48
1:B:272[B]:CYS:SG	1:B:316[B]:THR:HG22	2.49	0.47
1:B:101:ASN:HA	1:B:132:MET:HG3	1.96	0.47
1:E:404:THR:O	1:E:408:GLU:HB2	2.14	0.47
1:B:217:LYS:NZ	1:B:420:PHE:O	2.44	0.46
1:E:64:LEU:HB3	1:E:65:PRO:HD3	1.98	0.46
1:E:152:SER:O	1:E:187:PRO:HD2	2.17	0.45
1:B:115:ASP:O	1:B:118:ARG:HB3	2.17	0.45
1:B:429:LYS:NZ	5:B:634:HOH:O	2.50	0.45
1:B:30:LYS:NZ	1:B:55:GLY:O	2.46	0.43
1:E:72:ILE:HD11	1:E:121:PRO:HB2	1.99	0.43
1:B:142:LEU:O	1:B:146:VAL:HG23	2.19	0.42
1:B:238:PRO:HG3	5:B:637:HOH:O	2.19	0.42
1:B:279:LEU:HB3	1:B:284:VAL:HG13	2.02	0.42
1:E:180:ARG:HG2	1:E:207:LEU:O	2.21	0.41
1:B:180:ARG:NH1	1:B:210:LYS:HE2	2.36	0.41
1:E:331:VAL:HG23	1:E:350:ALA:HB2	2.03	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1001:HOH:O	5:E:872:HOH:O[2_755]	2.02	0.18
5:B:945:HOH:O	5:E:1015:HOH:O[2_755]	2.07	0.13
5:B:1117:HOH:O	5:E:1030:HOH:O[2_745]	2.14	0.06
5:B:957:HOH:O	5:B:965:HOH:O[2_756]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	442/445 (99%)	431 (98%)	11 (2%)	0	100	100
1	E	436/445 (98%)	428 (98%)	8 (2%)	0	100	100
All	All	878/890 (99%)	859 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	370/373 (99%)	368 (100%)	2 (0%)	92	91
1	E	366/373 (98%)	364 (100%)	2 (0%)	92	91
All	All	736/746 (99%)	732 (100%)	4 (0%)	90	91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	180	ARG
1	B	294	LYS
1	E	296	SER
1	E	453	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ADP	B	502	4	24,29,29	0.97	1 (4%)	23,45,45	1.78	2 (8%)
3	ADP	E	502	4	24,29,29	0.98	1 (4%)	23,45,45	1.83	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	B	502	4	-	0/12/32/32	0/3/3/3
3	ADP	E	502	4	-	0/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	502	ADP	C5-C4	2.95	1.47	1.40
3	B	502	ADP	C5-C4	3.05	1.47	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	502	ADP	N3-C2-N1	-6.74	123.58	128.87
3	E	502	ADP	N3-C2-N1	-6.59	123.69	128.87
3	B	502	ADP	C1'-N9-C4	-2.36	124.17	126.81
3	E	502	ADP	C1'-N9-C4	-2.01	124.57	126.81
3	E	502	ADP	O3B-PB-O2B	2.01	114.83	107.44
3	E	502	ADP	O4'-C1'-N9	2.35	112.55	108.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	502	ADP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	439/445 (98%)	0.33	37 (8%) 14 11	19, 31, 66, 90	0
1	E	439/445 (98%)	0.26	30 (6%) 20 16	18, 32, 62, 85	0
All	All	878/890 (98%)	0.29	67 (7%) 17 13	18, 32, 64, 90	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	323	VAL	6.6
1	B	185	HIS	5.9
1	E	253	GLU	5.8
1	B	82	LEU	5.7
1	E	257	GLY	5.1
1	E	12	PRO	4.8
1	B	453	TRP	4.7
1	E	147	SER	4.6
1	B	10	SER	4.2
1	E	82	LEU	3.8
1	B	253	GLU	3.7
1	E	258	LEU	3.7
1	B	92	THR	3.6
1	B	119	GLU	3.6
1	B	122	GLN	3.5
1	B	150	LEU	3.5
1	B	184	GLY	3.5
1	B	113	LEU	3.4
1	E	10	SER	3.4
1	E	11	ASP	3.4
1	B	118	ARG	3.3
1	B	81	ALA	3.3
1	B	297	GLU	3.3
1	E	106	ALA	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	185	HIS	3.1
1	E	318	SER	3.1
1	E	324	ASP	3.1
1	B	106	ALA	3.1
1	B	114	ALA	3.1
1	B	11	ASP	3.0
1	E	254	ALA	3.0
1	B	452	SER	2.9
1	B	80	ILE	2.9
1	E	165	TRP	2.9
1	E	388	ASN	2.9
1	B	324	ASP	2.8
1	E	107	GLN	2.7
1	E	120	LYS	2.6
1	B	79	LEU	2.6
1	E	319	PHE	2.5
1	B	147	SER	2.5
1	B	145	LEU	2.5
1	B	123	THR	2.5
1	B	70	LYS	2.5
1	E	81	ALA	2.4
1	B	110	LYS	2.4
1	B	317	ILE	2.4
1	E	102	SER	2.3
1	E	105	SER	2.3
1	B	140	PRO	2.3
1	B	88	ASP	2.3
1	B	136	SER	2.3
1	B	252	GLN	2.2
1	E	313	ILE	2.2
1	E	13	GLU	2.2
1	B	105	SER	2.2
1	B	149	HIS	2.2
1	E	252	GLN	2.2
1	E	114	ALA	2.1
1	E	389	LYS	2.1
1	B	183	LEU	2.1
1	B	138	PHE	2.1
1	E	390	ALA	2.0
1	E	108	GLU	2.0
1	B	303	ASN	2.0
1	B	296	SER	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	149	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ADP	E	502	27/27	0.96	0.08	-0.65	28,34,44,47	0
3	ADP	B	502	27/27	0.96	0.09	-1.08	32,36,43,47	0
2	ZN	B	501	1/1	1.00	0.06	-1.45	24,24,24,24	0
2	ZN	E	501	1/1	1.00	0.05	-2.98	21,21,21,21	0
4	MG	B	503	1/1	0.92	0.16	-	44,44,44,44	0
4	MG	E	503	1/1	0.77	0.23	-	46,46,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.