



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:15 AM GMT

PDB ID : 3LEM
Title : Crystal structure of fructosyltransferase (D191A) from *A. japonicus* in complex with Nystose
Authors : Chuankhayan, P.; Chen, C.J.; Chiang, C.M.
Deposited on : 2010-01-15
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	634	<div> <div>9%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	634	Total	C	N	O	S	0	0	0
			4883	3091	824	965	3			

- Molecule 2 is SUGAR (BETA-D-FRUCTOFURANOSYL-(2->1)-BETA-D-FRUCTOFURANOSYL-(2->1)-BETA-D-FRUCTOFURANOSYL ALPHA-D-GLUCOPYRANOSIDE) (three-letter code: NYT) (formula: C₂₄H₄₂O₂₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			45	24	21		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	219	Total	O	0	0
			219	219		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fructosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	98.68 Å 110.73 Å 66.00 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 27.25 – 1.94	Depositor EDS
% Data completeness (in resolution range)	96.1 (30.00-2.10) 90.8 (27.25-1.94)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 1.93 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.214 , 0.252 0.227 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 51998 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5147	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NYT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	3/5014 (0.1%)	0.70	2/6858 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	544	SER	C-N	5.54	1.46	1.34
1	A	547	SER	C-N	-5.30	1.21	1.34
1	A	608	ARG	C-N	5.27	1.46	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	344	GLY	C-N-CA	-5.72	107.41	121.70
1	A	456	GLN	O-C-N	-5.02	114.66	122.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	344	GLY	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4883	0	4609	94	0
2	A	45	0	42	3	0
3	A	219	0	0	1	0
All	All	5147	0	4651	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:ASN:HD21	1:A:373:GLY:H	1.16	0.91
1:A:61:PRO:O	1:A:372:ALA:HB2	1.78	0.83
1:A:591:ASP:H	1:A:606:ASN:HD21	1.20	0.83
1:A:33:LEU:H	1:A:562:ASN:HD21	1.29	0.81
1:A:61:PRO:O	1:A:372:ALA:CB	2.29	0.80
1:A:606:ASN:N	1:A:606:ASN:HD22	1.82	0.77
1:A:454:ASP:HB2	1:A:456:GLN:NE2	2.03	0.73
1:A:295:ASN:ND2	1:A:373:GLY:H	1.86	0.73
1:A:591:ASP:H	1:A:606:ASN:ND2	1.85	0.73
1:A:606:ASN:HD22	1:A:606:ASN:H	1.40	0.67
1:A:213:ASN:ND2	1:A:216:ALA:H	1.93	0.67
1:A:50:HIS:ND1	1:A:597:ASP:OD2	2.25	0.67
1:A:202:VAL:O	1:A:206:LEU:HG	1.96	0.65
1:A:162:ARG:HG3	1:A:169:ASP:OD2	1.96	0.65
1:A:60:ASP:OD2	2:A:3168:NYT:H13A	1.96	0.64
1:A:216:ALA:HA	1:A:219:GLN:OE1	1.97	0.64
1:A:545:ASN:O	1:A:546:GLU:C	2.37	0.62
1:A:66:THR:HG1	1:A:73:PHE:HD1	1.47	0.62
1:A:454:ASP:CB	1:A:456:GLN:NE2	2.64	0.61
1:A:439:GLU:H	1:A:439:GLU:CD	2.03	0.61
1:A:453:SER:HB2	1:A:458:ALA:HB2	1.83	0.60
1:A:475:LEU:HD22	1:A:646:LEU:HD22	1.84	0.60
1:A:33:LEU:H	1:A:562:ASN:ND2	1.97	0.60
1:A:556:SER:HB2	1:A:616:ARG:O	2.02	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:454:ASP:HB2	1:A:456:GLN:HE21	1.66	0.59
1:A:591:ASP:N	1:A:606:ASN:HD21	1.97	0.58
1:A:297:LEU:HD12	1:A:297:LEU:N	2.17	0.58
1:A:454:ASP:OD2	1:A:456:GLN:NE2	2.37	0.58
1:A:523:LEU:HD23	1:A:524:GLN:N	2.19	0.57
1:A:454:ASP:CB	1:A:456:GLN:HE21	2.18	0.57
1:A:531:ALA:O	1:A:624:GLN:HB2	2.05	0.57
1:A:320:SER:HA	1:A:330:SER:OG	2.05	0.57
1:A:295:ASN:HB2	1:A:314:THR:OG1	2.05	0.56
1:A:523:LEU:HD23	1:A:523:LEU:C	2.25	0.56
1:A:386:SER:OG	1:A:388:VAL:HG23	2.06	0.56
1:A:154:GLU:OE2	1:A:190:ARG:HD3	2.05	0.55
1:A:454:ASP:HB2	1:A:456:GLN:HG3	1.90	0.53
1:A:343:VAL:O	1:A:345:SER:N	2.41	0.53
1:A:315:LEU:N	1:A:315:LEU:HD23	2.24	0.52
1:A:244:PRO:HB2	1:A:291:PHE:CD1	2.44	0.52
1:A:57:GLN:HB2	1:A:418:THR:HB	1.92	0.52
1:A:582:ASN:C	1:A:584:GLN:N	2.62	0.52
1:A:452:GLU:O	1:A:458:ALA:HB1	2.09	0.52
1:A:456:GLN:NE2	1:A:457:THR:H	2.08	0.51
1:A:311:VAL:HG21	1:A:341:VAL:HG23	1.93	0.51
1:A:295:ASN:HD21	1:A:373:GLY:N	1.96	0.51
1:A:251:GLN:OE1	1:A:256:ALA:HA	2.10	0.51
1:A:277:TRP:HB3	1:A:284:ALA:HB3	1.93	0.51
1:A:344:GLY:O	1:A:345:SER:C	2.49	0.49
1:A:297:LEU:N	1:A:297:LEU:CD1	2.76	0.49
1:A:60:ASP:CG	2:A:3168:NYT:H13A	2.34	0.48
1:A:213:ASN:HD21	1:A:216:ALA:HB2	1.78	0.48
1:A:175:PRO:HG2	1:A:178:ALA:HB2	1.94	0.48
1:A:582:ASN:C	1:A:584:GLN:H	2.17	0.47
1:A:454:ASP:HB2	1:A:456:GLN:CG	2.44	0.47
1:A:239:VAL:HB	1:A:242:VAL:HB	1.95	0.47
1:A:606:ASN:N	1:A:606:ASN:ND2	2.54	0.47
1:A:45:TRP:CE2	1:A:616:ARG:HB3	2.50	0.47
1:A:118:PHE:HB2	1:A:136:THR:HB	1.97	0.47
1:A:206:LEU:CB	1:A:211:ALA:HB2	2.44	0.47
1:A:364:TRP:CE3	1:A:572:GLY:HA2	2.50	0.46
1:A:453:SER:HB2	1:A:458:ALA:CB	2.45	0.46
1:A:371:ALA:HB2	1:A:397:VAL:HG22	1.98	0.46
1:A:454:ASP:O	1:A:455:ASN:HB2	2.16	0.46
1:A:199:ARG:HH12	1:A:217:VAL:HG11	1.80	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:LEU:HD12	1:A:52:LEU:N	2.32	0.46
1:A:296:VAL:C	1:A:297:LEU:HD12	2.37	0.45
1:A:432:VAL:HB	1:A:460:LEU:HB2	1.98	0.45
1:A:437:ASP:OD2	1:A:442:ARG:NH2	2.33	0.45
1:A:423:LEU:O	1:A:425:ARG:HG3	2.17	0.45
1:A:538:ALA:HB3	1:A:551:ASP:HB3	1.99	0.45
1:A:454:ASP:CG	1:A:456:GLN:NE2	2.70	0.45
1:A:522:PRO:HB2	1:A:632:GLU:HG3	1.99	0.45
1:A:312:PHE:CE1	1:A:338:ALA:HB2	2.51	0.45
1:A:143:ILE:O	2:A:3168:NYT:H62	2.17	0.44
1:A:65:TYR:HB3	1:A:122:VAL:HB	2.00	0.44
1:A:50:HIS:NE2	1:A:602:GLU:OE2	2.50	0.44
1:A:453:SER:CB	1:A:458:ALA:HB2	2.46	0.44
1:A:300:THR:O	1:A:429:VAL:HB	2.17	0.44
1:A:32:ASN:ND2	1:A:35:THR:HG23	2.33	0.44
1:A:523:LEU:HD21	1:A:525:SER:OG	2.19	0.43
1:A:606:ASN:C	1:A:608:ARG:H	2.22	0.42
1:A:199:ARG:HH11	1:A:199:ARG:HG3	1.85	0.42
1:A:197:SER:HB2	1:A:298:PHE:CZ	2.55	0.42
1:A:526:GLY:HA3	1:A:539:ILE:O	2.20	0.42
1:A:108:PRO:HB3	1:A:116:ALA:HA	2.01	0.42
1:A:213:ASN:HD22	1:A:215:THR:HB	1.84	0.41
1:A:128:ASN:O	1:A:130:THR:HG23	2.20	0.41
1:A:454:ASP:HB2	1:A:456:GLN:CD	2.41	0.41
1:A:264:TYR:CZ	1:A:266:GLY:HA2	2.56	0.41
1:A:199:ARG:NH1	1:A:217:VAL:HG11	2.36	0.40
1:A:284:ALA:O	1:A:285:GLY:C	2.60	0.40
1:A:208:GLU:HG3	1:A:209:GLU:OE1	2.22	0.40
1:A:188:ALA:HB2	3:A:736:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	632/634 (100%)	600 (95%)	31 (5%)	1 (0%)	52 53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	607	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	516/516 (100%)	502 (97%)	14 (3%)	52 56

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LEU
1	A	55	GLU
1	A	57	GLN
1	A	119	ASP
1	A	181	PRO
1	A	209	GLU
1	A	228	ASN
1	A	268	TRP
1	A	297	LEU
1	A	347	GLN
1	A	427	LEU
1	A	456	GLN
1	A	476	LEU
1	A	606	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	57	GLN
1	A	156	GLN
1	A	213	ASN
1	A	295	ASN
1	A	434	ASN
1	A	456	GLN
1	A	562	ASN
1	A	606	ASN
1	A	639	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NYT	A	3168	-	48,48,48	2.06	9 (18%)	69,74,74	2.15	13 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NYT	A	3168	-	-	0/30/107/107	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	3168	NYT	O1-C1	2.01	1.50	1.40
2	A	3168	NYT	O21-C51	2.01	1.48	1.43
2	A	3168	NYT	C1-C2	2.40	1.59	1.52
2	A	3168	NYT	O5-C1	2.81	1.49	1.41
2	A	3168	NYT	O11-C11	3.17	1.48	1.42
2	A	3168	NYT	C11-C21	4.97	1.58	1.52
2	A	3168	NYT	O12-C12	5.58	1.52	1.42
2	A	3168	NYT	O12-C23	5.95	1.49	1.41
2	A	3168	NYT	O11-C22	6.27	1.50	1.41

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3168	NYT	C11-O11-C22	-5.01	107.78	116.04
2	A	3168	NYT	C62-C52-C42	-3.53	106.74	115.08
2	A	3168	NYT	C12-C22-C32	-2.95	108.74	115.17
2	A	3168	NYT	O21-C21-C31	-2.94	98.89	105.58
2	A	3168	NYT	O21-C21-C11	-2.79	101.17	107.59
2	A	3168	NYT	C63-C53-C43	-2.55	109.04	115.08
2	A	3168	NYT	O12-C23-C13	-2.30	102.45	109.67
2	A	3168	NYT	O1-C21-C11	-2.27	102.33	109.85
2	A	3168	NYT	O1-C21-O21	2.17	117.40	110.52
2	A	3168	NYT	O1-C21-C31	4.40	123.37	108.04
2	A	3168	NYT	O31-C31-C21	5.52	131.66	113.96
2	A	3168	NYT	O62-C62-C52	7.30	135.45	111.33
2	A	3168	NYT	C21-O1-C1	9.31	142.06	117.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3168	NYT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	634/634 (100%)	0.41	57 (8%)	12 16	20, 36, 66, 97	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	345	SER	10.0
1	A	211	ALA	8.1
1	A	21	TYR	8.0
1	A	212	ARG	7.5
1	A	20	SER	7.5
1	A	346	GLU	7.3
1	A	206	LEU	7.1
1	A	348	GLU	7.0
1	A	347	GLN	6.5
1	A	209	GLU	5.8
1	A	208	GLU	5.5
1	A	217	VAL	5.1
1	A	210	VAL	4.8
1	A	207	ASP	4.8
1	A	279	ASP	4.7
1	A	213	ASN	4.7
1	A	349	GLY	4.5
1	A	456	GLN	4.5
1	A	455	ASN	4.4
1	A	227	LYS	4.2
1	A	322	LEU	4.1
1	A	164	GLY	3.9
1	A	214	GLU	3.7
1	A	215	THR	3.6
1	A	216	ALA	3.4
1	A	582	ASN	3.4
1	A	452	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	218	GLN	3.1
1	A	453	SER	3.1
1	A	592	LEU	3.1
1	A	454	ASP	2.9
1	A	280	GLU	2.9
1	A	583	GLY	2.9
1	A	439	GLU	2.8
1	A	590	LEU	2.8
1	A	389	GLU	2.8
1	A	121	ALA	2.7
1	A	476	LEU	2.7
1	A	228	ASN	2.7
1	A	581	GLU	2.6
1	A	478	ASN	2.6
1	A	281	GLY	2.5
1	A	603	VAL	2.4
1	A	584	GLN	2.4
1	A	194	VAL	2.3
1	A	451	GLY	2.3
1	A	101	ASN	2.3
1	A	219	GLN	2.2
1	A	344	GLY	2.2
1	A	192	PRO	2.2
1	A	574	LEU	2.2
1	A	601	VAL	2.2
1	A	233	VAL	2.1
1	A	576	LEU	2.1
1	A	191	ALA	2.1
1	A	100	ASP	2.0
1	A	323	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NYT	A	3168	45/45	0.85	0.16	1.53	33,40,62,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.