



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2017 – 06:00 PM EST

PDB ID : 5LG4  
Title : Crystal structure of the Sec3/Sso2 complex at 2.9 angstrom resolution  
Authors : Zhang, Y.B.; Dong, G.  
Deposited on : 2016-07-05  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• •</div> </div> </div>
2	B	250	<div> <div>%</div> <div> <div></div> <div>60%</div> <div>9%</div> <div>30%</div> </div> </div>
2	C	250	<div> <div>6%</div> <div> <div></div> <div>57%</div> <div>11%</div> <div>•</div> <div>31%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4421 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein SSO2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	188	Total	C	N	O	S	0	0	0
			1531	945	266	314	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	GLY	-	expression tag	UNP P39926
A	33	SER	-	expression tag	UNP P39926
A	34	HIS	-	expression tag	UNP P39926
A	35	MET	-	expression tag	UNP P39926


- Molecule 2 is a protein called Exocyst complex component SEC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	174	Total	C	N	O	S	0	0	0
			1441	919	251	267	4			
2	C	172	Total	C	N	O	S	0	0	0
			1429	912	249	265	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	GLN	-	expression tag	UNP P33332
B	72	GLY	-	expression tag	UNP P33332
B	73	HIS	-	expression tag	UNP P33332
B	74	MET	-	expression tag	UNP P33332
C	71	GLN	-	expression tag	UNP P33332
C	72	GLY	-	expression tag	UNP P33332
C	73	HIS	-	expression tag	UNP P33332
C	74	MET	-	expression tag	UNP P33332

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.12Å 84.12Å 237.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.83 – 2.90 48.44 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.83-2.90) 99.5 (48.44-2.70)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, $R_{free}$	0.260 , 0.281 0.264 , 0.287	Depositor DCC
$R_{free}$ test set	1634 reflections (8.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	105.3	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 62.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4421	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	109.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.23	0/1545	0.35	0/2076
2	B	0.25	0/1473	0.42	0/1988
2	C	0.24	0/1461	0.40	0/1973
All	All	0.24	0/4479	0.39	0/6037

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1531	0	1496	14	0
2	B	1441	0	1441	14	0
2	C	1429	0	1430	16	0
3	A	10	0	0	0	0
3	B	10	0	0	1	0
All	All	4421	0	4367	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:157:ARG:NH2	3:B:401:SO4:O4	2.31	0.64
2:C:114:ILE:HD13	2:C:139:LEU:HG	1.82	0.59
2:C:161:ASP:OD2	2:C:161:ASP:N	2.37	0.58
1:A:172:ASN:OD1	1:A:172:ASN:N	2.37	0.57
2:B:118:SER:HB3	2:B:135:LYS:HE3	1.87	0.57
1:A:134:ARG:HH21	1:A:215:THR:HA	1.70	0.56
1:A:107:ARG:HA	1:A:110:LEU:HG	1.89	0.55
2:C:137:ARG:NH2	2:C:157:ARG:HE	2.06	0.54
2:C:81:GLN:HE21	2:C:113:ARG:HE	1.57	0.52
2:C:75:SER:O	2:C:77:PHE:N	2.42	0.51
2:C:108:TYR:OH	2:C:111:HIS:ND1	2.26	0.51
2:C:140:ILE:HB	2:C:154:HIS:HB2	1.91	0.51
1:A:62:ASP:HB2	1:A:136:ILE:HD13	1.92	0.51
2:B:184:ILE:HG22	2:B:186:GLU:H	1.76	0.51
1:A:181:LEU:HD13	1:A:194:LEU:HD13	1.92	0.50
1:A:57:ILE:HD12	1:A:94:LEU:HD12	1.93	0.50
1:A:195:ALA:HB2	2:C:182:LEU:HD12	1.92	0.49
2:B:168:ARG:HG2	2:B:170:TRP:CZ2	2.47	0.49
1:A:127:LEU:HD21	1:A:221:MET:HB2	1.95	0.48
2:B:114:ILE:HD12	2:B:198:TRP:CD1	2.47	0.48
2:B:114:ILE:HD13	2:B:139:LEU:HG	1.96	0.48
2:C:192:MET:HB2	2:C:193:SER:H	1.54	0.46
1:A:62:ASP:OD2	1:A:66:LYS:NZ	2.49	0.46
2:B:214:ILE:O	2:B:218:ILE:HG13	2.15	0.46
2:B:192:MET:HB2	2:B:193:SER:H	1.53	0.46
1:A:159:ALA:HB1	1:A:163:GLU:HB2	1.97	0.45
2:C:151:ILE:HB	2:C:172:LEU:HD22	1.98	0.45
2:C:180:LYS:HZ2	2:C:206:ARG:HH12	1.63	0.45
2:C:179:GLU:HB3	2:C:189:ILE:HB	1.99	0.45
2:B:142:SER:OG	2:B:154:HIS:NE2	2.46	0.45
2:B:158:GLU:HG2	2:B:164:PHE:CE1	2.52	0.45
2:C:218:ILE:HG12	2:C:225:VAL:HG12	1.98	0.45
2:C:135:LYS:HB3	2:C:137:ARG:NH1	2.32	0.44
2:C:239:ASP:H	2:C:242:SER:HG	1.65	0.44
1:A:171:VAL:HG23	1:A:172:ASN:HA	2.00	0.44
2:B:98:ASP:OD2	2:B:99:HIS:N	2.50	0.43
1:A:60:GLN:O	1:A:64:GLN:HG2	2.19	0.43
1:A:64:GLN:HB2	1:A:84:LEU:HD13	2.00	0.42
2:B:217:TYR:CD1	2:B:226:PRO:HD3	2.54	0.42
1:A:158:GLU:O	1:A:160:THR:N	2.44	0.41
2:C:144:LYS:HE3	2:C:152:GLN:NE2	2.36	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:184:ILE:HG22	2:B:187:GLY:H	1.86	0.40
2:B:87:LYS:HE2	2:B:87:LYS:HB3	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/196 (94%)	179 (97%)	4 (2%)	1 (0%)	34	71
2	B	170/250 (68%)	157 (92%)	13 (8%)	0	100	100
2	C	168/250 (67%)	159 (95%)	8 (5%)	1 (1%)	30	67
All	All	522/696 (75%)	495 (95%)	25 (5%)	2 (0%)	39	74

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	76	ASN
1	A	170	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/173 (97%)	160 (95%)	8 (5%)	31	67

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	162/230 (70%)	161 (99%)	1 (1%)	90	97
2	C	161/230 (70%)	159 (99%)	2 (1%)	78	94
All	All	491/633 (78%)	480 (98%)	11 (2%)	60	88

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	VAL
1	A	76	GLN
1	A	147	GLN
1	A	154	ILE
1	A	162	GLU
1	A	172	ASN
1	A	181	LEU
1	A	224	LEU
2	B	192	MET
2	C	161	ASP
2	C	192	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	301	1	4,4,4	0.18	0	6,6,6	0.08	0
3	SO4	A	302	-	4,4,4	0.21	0	6,6,6	0.06	0
3	SO4	B	401	-	4,4,4	0.16	0	6,6,6	0.12	0
3	SO4	B	402	-	4,4,4	0.20	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	301	1	-	0/0/0/0	0/0/0/0
3	SO4	A	302	-	-	0/0/0/0	0/0/0/0
3	SO4	B	401	-	-	0/0/0/0	0/0/0/0
3	SO4	B	402	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	SO4	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	188/196 (95%)	0.53	12 (6%) 23 16	87, 107, 143, 153	0
2	B	174/250 (69%)	0.19	3 (1%) 73 70	79, 95, 117, 127	0
2	C	172/250 (68%)	0.64	16 (9%) 11 7	99, 119, 144, 154	0
All	All	534/696 (76%)	0.46	31 (5%) 26 20	79, 107, 141, 154	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	TYR	5.8
1	A	170	ASP	4.5
2	C	96	ARG	4.3
2	C	156	ALA	4.0
2	C	158	GLU	3.7
1	A	75	GLU	3.7
1	A	162	GLU	3.5
1	A	153	THR	3.4
1	A	161	ASP	3.4
1	A	180	ALA	3.2
2	C	91	ASN	3.1
2	C	89	ILE	3.1
2	C	94	PHE	2.9
2	C	95	SER	2.7
2	C	134	LYS	2.7
2	C	132	GLU	2.4
2	B	103	GLU	2.4
2	C	241	ARG	2.4
1	A	157	PRO	2.4
2	C	131	LEU	2.4
1	A	78	MET	2.3
2	B	229	VAL	2.3
2	C	139	LEU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	156	GLN	2.3
2	C	167	GLY	2.3
2	C	118	SER	2.3
1	A	155	ILE	2.2
1	A	158	GLU	2.2
2	C	160	SER	2.1
2	B	146	ASN	2.1
2	C	128	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	B	401	5/5	0.94	0.22	-0.11	106,107,108,108	0
3	SO4	A	301	5/5	0.93	0.17	-1.20	97,97,98,98	0
3	SO4	B	402	5/5	0.89	0.15	-	106,106,108,108	0
3	SO4	A	302	5/5	0.95	0.12	-	113,113,114,116	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.