



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:16 AM GMT

PDB ID : 3LIJ
Title : Crystal structure of full length CpCDPK3 (cgd5_820) in complex with Ca²⁺ and AMPPNP
Authors : Qiu, W.; Hutchinson, A.; Wernimont, A.; Walker, J.R.; Sullivan, H.; Lin, Y.-H.; Mackenzie, F.; Kozieradzki, I.; Cossar, D.; Schapira, M.; Senisterra, G.; Vedadi, M.; Arrowsmith, C.H.; Bountra, C.; Weigelt, J.; Edwards, A.M.; Bochkarev, A.; Hui, R.; Amani, M.; Structural Genomics Consortium (SGC)
Deposited on : 2010-01-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calcium/calmodulin dependent protein kinase with a kinase domain and 4 calmodulin like EF hands.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	1	0
			3752	2385	629	719	19			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	EXPRESSION TAG	UNP Q5CS01
A	25	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	26	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	27	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	28	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	29	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	30	HIS	-	EXPRESSION TAG	UNP Q5CS01
A	31	SER	-	EXPRESSION TAG	UNP Q5CS01
A	32	SER	-	EXPRESSION TAG	UNP Q5CS01
A	33	GLY	-	EXPRESSION TAG	UNP Q5CS01
A	34	ARG	-	EXPRESSION TAG	UNP Q5CS01
A	35	GLU	-	EXPRESSION TAG	UNP Q5CS01
A	36	ASN	-	EXPRESSION TAG	UNP Q5CS01
A	37	LEU	-	EXPRESSION TAG	UNP Q5CS01
A	38	TYR	-	EXPRESSION TAG	UNP Q5CS01
A	39	PHE	-	EXPRESSION TAG	UNP Q5CS01
A	40	GLN	-	EXPRESSION TAG	UNP Q5CS01
A	41	GLY	-	EXPRESSION TAG	UNP Q5CS01

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 3 | A | 1 | Total Mg
1 1 | 0 | 0 |

-


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 6	O 12	P 3	0	0

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 5 | A | 252 | Total O
252 252 | 0 | 0 |

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Calcium/calmodulin dependent protein kinase with a kinase domain and 4 calmodulin like EF hands



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.08 Å 88.06 Å 58.61 Å 90.00° 118.07° 90.00°	Depositor
Resolution (Å)	19.56 – 1.90 19.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (19.56-1.90) 99.8 (19.40-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.90 Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.203 , 0.229 0.216 , 0.243	Depositor DCC
R_{free} test set	2013 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.7	EDS
Estimated twinning fraction	0.000 for -h-l,k,h 0.000 for l,k,-h-l 0.026 for h,-k,-h-l 0.034 for -h-l,-k,l 0.034 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40168 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4040	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3812	0.61	0/5108

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3752	0	3749	18	0
2	A	4	0	0	0	0
3	A	1	0	0	0	0
4	A	31	0	13	1	0
5	A	252	0	0	2	0
All	All	4040	0	3762	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:ARG:HG2	1:A:443:LYS:N	1.92	0.83
1:A:345:GLN:O	1:A:346:ASN:HB2	1.87	0.74
1:A:442:ARG:CG	1:A:443:LYS:N	2.63	0.60
1:A:345:GLN:O	1:A:346:ASN:CB	2.47	0.58
1:A:69:GLY:HA3	4:A:522:ANP:H5'1	1.92	0.51
1:A:300:GLN:HE21	1:A:302:ASP:H	1.59	0.50
1:A:68:LEU:HD13	1:A:78:LEU:HB2	1.93	0.49
1:A:172:VAL:HG21	1:A:246:ILE:HD11	1.95	0.49
1:A:243:VAL:HG11	1:A:307:ILE:O	2.17	0.45
1:A:94:ARG:HA	1:A:133:ASN:HD22	1.82	0.45
1:A:163:VAL:HG21	1:A:325:LYS:HG3	1.99	0.44
1:A:345:GLN:HG3	5:A:744:HOH:O	2.17	0.43
1:A:130:ASP:HB2	1:A:135:TYR:HE2	1.83	0.43
1:A:391:LEU:HD21	1:A:430:ILE:HD12	2.00	0.43
1:A:358:MET:HA	1:A:362:LEU:HB2	2.01	0.42
1:A:233:LEU:HD23	1:A:273:GLU:HG3	2.02	0.42
1:A:345:GLN:NE2	5:A:744:HOH:O	2.54	0.41
1:A:255:ALA:HB1	1:A:257:TYR:CE2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/494 (92%)	441 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/437 (94%)	388 (94%)	25 (6%)	23	11

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LEU
1	A	68	LEU
1	A	77	LEU
1	A	130	ASP
1	A	131	LYS
1	A	132	ARG
1	A	165	ILE
1	A	233	LEU
1	A	264	THR
1	A	281	SER
1	A	325	LYS
1	A	331	LEU
1	A	334	LEU
1	A	339	GLU
1	A	346	ASN
1	A	362	LEU
1	A	369	LYS
1	A	391	LEU
1	A	402	GLU
1	A	411	ILE
1	A	442	ARG
1	A	468	VAL
1	A	471	LEU
1	A	513	CYS
1	A	515	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	HIS
1	A	133	ASN
1	A	300	GLN
1	A	345	GLN
1	A	346	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	456	GLN
1	A	515	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ANP	A	522	3	27,33,33	2.81	5 (18%)	30,52,52	2.24	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	522	3	-	0/12/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	522	ANP	PG-O2G	-2.93	1.48	1.56
4	A	522	ANP	O4'-C1'	2.11	1.43	1.41
4	A	522	ANP	PB-O1B	2.22	1.48	1.46
4	A	522	ANP	PB-O3A	2.51	1.62	1.59
4	A	522	ANP	PG-O1G	13.20	1.61	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	522	ANP	N3-C2-N1	-9.31	121.77	128.89
4	A	522	ANP	PA-O3A-PB	-3.44	121.14	132.67
4	A	522	ANP	O1G-PG-N3B	-3.31	106.82	111.90
4	A	522	ANP	C4'-O4'-C1'	-3.18	106.23	109.72
4	A	522	ANP	O3G-PG-O1G	-3.05	105.39	113.49
4	A	522	ANP	C4-C5-N7	-2.38	107.29	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	522	ANP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	465/494 (94%)	0.63	54 (11%) 6 7	22, 39, 80, 102	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	SER	7.2
1	A	331	LEU	6.5
1	A	479	HIS	6.0
1	A	480	LEU	5.7
1	A	102	SER	5.6
1	A	333	SER	5.1
1	A	132	ARG	4.8
1	A	404	ALA	4.8
1	A	332	PRO	4.7
1	A	403	VAL	4.6
1	A	402	GLU	4.6
1	A	401	GLU	4.3
1	A	477	LEU	4.2
1	A	516	ASN	4.2
1	A	74	GLY	3.9
1	A	335	ALA	3.8
1	A	411	ILE	3.8
1	A	336	ASN	3.7
1	A	475	PHE	3.6
1	A	410	GLN	3.5
1	A	263	GLN	3.4
1	A	481	GLU	3.4
1	A	94	ARG	3.3
1	A	104	SER	3.1
1	A	103	ASN	3.1
1	A	513	CYS	3.0
1	A	57	HIS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	400	GLY	3.0
1	A	60	GLU	2.9
1	A	105	LYS	2.9
1	A	399	SER	2.7
1	A	197	LYS	2.5
1	A	406	PHE	2.5
1	A	264	THR	2.5
1	A	131	LYS	2.5
1	A	70	SER	2.5
1	A	474	VAL	2.4
1	A	59	SER	2.4
1	A	55	LYS	2.4
1	A	204	VAL	2.3
1	A	63	GLN	2.3
1	A	38	TYR	2.3
1	A	342	ARG	2.2
1	A	92	ILE	2.2
1	A	83	VAL	2.2
1	A	66	LYS	2.2
1	A	136	LEU	2.2
1	A	338	ILE	2.1
1	A	115	LEU	2.1
1	A	405	VAL	2.1
1	A	355	LEU	2.1
1	A	438	VAL	2.1
1	A	280	ASP	2.1
1	A	248	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ANP	A	522	31/31	0.92	0.12	-0.46	28,57,141,174	0
2	CA	A	1	1/1	0.99	0.02	-3.08	20,20,20,20	0
2	CA	A	520	1/1	0.99	0.04	-3.21	24,24,24,24	0
2	CA	A	519	1/1	0.99	0.03	-3.30	43,43,43,43	0
2	CA	A	518	1/1	0.99	0.02	-3.39	25,25,25,25	0
3	MG	A	521	1/1	0.95	0.22	-	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.