



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:26 PM GMT

PDB ID : 4LLH
Title : Substrate bound outward-open state of the symporter BetP
Authors : Perez, C.; Faust, B.; Ziegler, C.
Deposited on : 2013-07-09
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	566	<div> <div>9%</div> <div>47%</div> <div>35%</div> <div>9%</div> <div>7%</div> </div>
1	B	566	<div> <div>9%</div> <div>40%</div> <div>40%</div> <div>6%</div> <div>13%</div> </div>
1	C	566	<div> <div>7%</div> <div>45%</div> <div>38%</div> <div>•</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1Y8	C	603	-	-	-	X
3	NA	B	602	-	-	-	X
5	CM5	C	602	X	-	-	X
5	CM5	C	604	X	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11589 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycine betaine transporter BetP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			4001	2625	661	699	16			
1	B	492	Total	C	N	O	S	0	0	0
			3701	2443	590	652	16			
1	C	490	Total	C	N	O	S	0	0	0
			3704	2451	587	650	16			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582
B	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582
C	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582

- Molecule 2 is 2-(TRIMETHYL-LAMBDA 5 -ARSANYL)ETHANOL (three-letter code: 1Y8) (formula: C₅H₁₄AsO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	As	C	O	0	0
			7	1	5	1		
2	B	1	Total	As	C	O	0	0
			7	1	5	1		
2	C	1	Total	As	C	O	0	0
			7	1	5	1		


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is 5-CYCLOHEXYL-1-PENTYL-BETA-D-MALTOSE (three-letter code: CM5) (formula: C₂₃H₄₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			34	23	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			34	23	11		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	25	Total	O	0	0
			25	25		
6	B	26	Total	O	0	0
			26	26		
6	C	40	Total	O	0	0
			40	40		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glycine betaine transporter BetP





• Molecule 1: Glycine betaine transporter BetP



VAL
HIS
ASN
GLU
HIS
ARG
LYS
ARG
GLU
LEU
ALA
ALA
LYS
ARG
ARG
ARG
GLU
ARG
LYS
ALA
SER
GLY
ALA
GLY
LYS
ARG
ARG

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	115.47Å 129.69Å 164.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.30 – 2.80 36.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	82.5 (36.30-2.80) 75.8 (36.33-2.80)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.73 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.240 , 0.290 0.242 , 0.287	Depositor DCC
R_{free} test set	2350 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	55.4	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 74.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 46434 reflections	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	11589	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CM5, 1Y8, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/4102	0.42	0/5584
1	B	0.24	0/3798	0.41	0/5184
1	C	0.25	0/3801	0.41	0/5186
All	All	0.24	0/11701	0.41	0/15954

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4001	0	4032	240	0
1	B	3701	0	3723	260	0
1	C	3704	0	3730	200	0
2	A	7	0	3	0	0
2	B	7	0	3	2	0
2	C	7	0	3	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	C	1	0	0	1	0
5	C	68	0	84	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	25	0	0	3	0
6	B	26	0	0	3	0
6	C	40	0	0	5	0
All	All	11589	0	11578	693	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 693 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:TRP:HD1	1:A:388:ILE:HG22	1.30	0.94
1:A:295:ILE:HG22	1:A:296:SER:H	1.33	0.93
1:A:297:GLY:HA3	1:A:300:LYS:H	1.37	0.90
5:C:604:CM5:H17	5:C:604:CM5:H29	1.54	0.90
1:A:162:PRO:HG2	1:A:417:GLY:HA3	1.55	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	520/566 (92%)	445 (86%)	53 (10%)	22 (4%)	3	11
1	B	490/566 (87%)	424 (86%)	50 (10%)	16 (3%)	5	16
1	C	484/566 (86%)	429 (89%)	43 (9%)	12 (2%)	7	24
All	All	1494/1698 (88%)	1298 (87%)	146 (10%)	50 (3%)	5	16

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	ASP
1	A	138	THR
1	A	139	VAL
1	A	228	LYS
1	A	274	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/443 (93%)	359 (88%)	51 (12%)	6	17
1	B	381/443 (86%)	348 (91%)	33 (9%)	13	35
1	C	382/443 (86%)	351 (92%)	31 (8%)	15	39
All	All	1173/1329 (88%)	1058 (90%)	115 (10%)	10	28

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	60	TRP
1	B	217	LEU
1	C	377	TRP
1	B	80	PHE
1	B	127	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	481	GLN
1	B	482	HIS
1	C	346	GLN
1	B	343	ASN
1	C	331	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	1Y8	A	601	-	1,6,6	0.68	0	1,8,8	0.82	0
2	1Y8	B	601	-	1,6,6	0.64	0	1,8,8	0.92	0
5	CM5	C	602	-	36,36,36	1.45	6 (16%)	49,49,49	1.35	6 (12%)
2	1Y8	C	603	-	1,6,6	0.66	0	1,8,8	0.84	0
5	CM5	C	604	-	36,36,36	1.49	5 (13%)	49,49,49	1.59	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1Y8	A	601	-	-	0/3/4/4	0/0/0/0
2	1Y8	B	601	-	-	0/3/4/4	0/0/0/0
5	CM5	C	602	-	4/4/11/11	0/17/65/65	0/3/3/3
2	1Y8	C	603	-	-	0/3/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CM5	C	604	-	6/6/11/11	0/17/65/65	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	604	CM5	C17-C18	-4.80	1.39	1.52
5	C	602	CM5	C17-C18	-4.02	1.41	1.52
5	C	602	CM5	C17-C16	-3.71	1.41	1.52
5	C	604	CM5	C17-C16	-3.61	1.42	1.52
5	C	604	CM5	C28-C27	-3.60	1.42	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	602	CM5	O14-C15-C16	2.02	114.02	109.75
5	C	604	CM5	C18-C17-C16	2.07	114.14	109.60
5	C	602	CM5	O31-C30-C26	2.08	118.21	111.33
5	C	602	CM5	O12-C1-C2	2.16	118.49	109.88
5	C	604	CM5	O31-C30-C26	2.28	118.86	111.33

5 of 10 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	C	604	CM5	C18
5	C	604	CM5	C16
5	C	604	CM5	C24
5	C	604	CM5	C27
5	C	604	CM5	C13

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	1Y8	2	0
5	C	602	CM5	2	0
5	C	604	CM5	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/566 (92%)	0.32	49 (9%) 11 5	20, 68, 134, 227	0
1	B	492/566 (86%)	0.32	52 (10%) 8 4	16, 68, 151, 205	0
1	C	490/566 (86%)	0.01	37 (7%) 17 9	12, 50, 124, 192	0
All	All	1506/1698 (88%)	0.22	138 (9%) 11 5	12, 64, 136, 227	0

The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	298	VAL	13.2
1	A	297	GLY	11.8
1	A	82	ASP	10.6
1	B	511	GLY	10.0
1	C	296	SER	9.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NA	B	602	1/1	0.89	1.27	10.38	96,96,96,96	0
2	1Y8	C	603	7/7	0.90	0.64	5.58	48,89,134,236	0
5	CM5	C	604	34/34	0.77	0.23	2.60	94,126,142,144	0
5	CM5	C	602	34/34	0.95	0.27	2.53	42,67,71,71	0
3	NA	A	602	1/1	0.80	0.35	0.99	62,62,62,62	0
2	1Y8	B	601	7/7	0.97	0.28	-0.35	86,96,132,230	0
2	1Y8	A	601	7/7	0.95	0.27	-0.47	44,95,161,167	0
4	CL	C	601	1/1	0.98	0.06	-1.54	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.