



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:43 PM GMT

PDB ID : 1LOF
Title : X-RAY STRUCTURE OF A BIANENNARY OCTASACCHARIDE-LECT
IN COMPLEX AT 2.3 ANGSTROMS RESOLUTION
Authors : Bourne, Y.; Cambillau, C.
Deposited on : 1993-01-27
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	<div> <div>68%</div> <div>25%</div> <div>6%</div> </div>
2	B	52	<div> <div>58%</div> <div>31%</div> <div>10%</div> </div>
3	C	181	<div> <div>72%</div> <div>23%</div> </div>
4	D	52	<div> <div>65%</div> <div>31%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	504	-	-	X	X
5	NAG	A	505	-	-	-	X
6	GAL	A	506	-	-	X	X
8	CA	A	227	-	-	-	X
8	CA	C	457	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEGUME ISOLECTIN I (ALPHA CHAIN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	0	0	0
			1406	894	232	280			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	153	ALA	LYS	CONFLICT	UNP P04122

- Molecule 2 is a protein called LEGUME ISOLECTIN I (BETA CHAIN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	47	Total	C	N	O	0	0	0
			381	251	58	72			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	41	TYR	PHE	CONFLICT	UNP P12306

- Molecule 3 is a protein called LEGUME ISOLECTIN I (ALPHA CHAIN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	180	Total	C	N	O	0	0	0
			1401	893	231	277			

- Molecule 4 is a protein called LEGUME ISOLECTIN I (BETA CHAIN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	52	Total	C	N	O	0	0	0
			413	267	64	82			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	41	TYR	PHE	CONFLICT	UNP P12306
D	?	-	SER	DELETION	UNP P12306

- Molecule 5 is a polymer of unknown type called SUGAR (8-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	7	Total	C	N	O	0	0
			87	48	3	36		

- Molecule 6 is SUGAR (3-MER) (three-letter code: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 7 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	3	Total	C	N	O	0	0
			36	20	1	15		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Ca 1 1	0	0
8	C	1	Total Ca 1 1	0	0

- Molecule 9 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Mn 1 1	0	0
9	C	1	Total Mn 1 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	99	Total O 99 99	0	0
10	B	15	Total O 15 15	0	0
10	C	97	Total O 97 97	0	0
10	D	19	Total O 19 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LEGUME ISOLECTIN I (ALPHA CHAIN)




- Molecule 2: LEGUME ISOLECTIN I (BETA CHAIN)



- Molecule 3: LEGUME ISOLECTIN I (ALPHA CHAIN)



- Molecule 4: LEGUME ISOLECTIN I (BETA CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.30 Å 75.40 Å 103.90 Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.30 7.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-2.30) 85.3 (7.99-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.30 Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.190 , (Not available) 0.202 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.309	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 105.8	EDS
Estimated twinning fraction	0.023 for k,h,-l 0.021 for -k,-h,-l 0.021 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 22459 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3969	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CA, MN, GAL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	1/1440 (0.1%)	1.63	24/1965 (1.2%)
2	B	1.06	0/394	1.64	5/539 (0.9%)
3	C	0.95	0/1435	1.65	19/1958 (1.0%)
4	D	1.12	0/426	1.71	8/581 (1.4%)
All	All	0.97	1/3695 (0.0%)	1.65	56/5043 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	68	THR	CA-CB	5.46	1.67	1.53

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172	VAL	CG1-CB-CG2	-10.28	94.45	110.90
3	C	53	TRP	CD1-CG-CD2	9.37	113.80	106.30
3	C	152	TRP	CD1-CG-CD2	9.08	113.56	106.30
1	A	128	TRP	CE2-CD2-CG	-8.81	100.25	107.30
1	A	128	TRP	CD1-CG-CD2	8.71	113.27	106.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1352	33	0
2	B	381	0	357	16	0
3	C	1401	0	1354	36	0
4	D	413	0	383	14	0
5	A	87	0	74	14	0
6	A	11	0	10	7	0
7	C	36	0	31	8	0
8	A	1	0	0	0	0
8	C	1	0	0	0	0
9	A	1	0	0	0	0
9	C	1	0	0	0	0
10	A	99	0	0	12	0
10	B	15	0	0	1	0
10	C	97	0	0	7	0
10	D	19	0	0	1	0
All	All	3969	0	3561	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:504:NAG:C1	5:A:504:NAG:O7	2.11	0.99
3:C:39:ASN:HD21	7:C:515:NAG:H3	1.29	0.96
5:A:504:NAG:C4	6:A:506:GAL:C1	2.45	0.95
1:A:122:THR:HA	1:A:135:ARG:HG3	1.54	0.87
5:A:507:GAL:H61	10:A:851:HOH:O	1.74	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/181 (99%)	172 (96%)	6 (3%)	1 (1%)	30	36
2	B	45/52 (86%)	41 (91%)	3 (7%)	1 (2%)	8	6
3	C	178/181 (98%)	170 (96%)	8 (4%)	0	100	100
4	D	50/52 (96%)	45 (90%)	4 (8%)	1 (2%)	9	7
All	All	452/466 (97%)	428 (95%)	21 (5%)	3 (1%)	26	31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	51	SER
2	B	11	PRO
1	A	180	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/155 (100%)	134 (86%)	21 (14%)	5	4
2	B	40/44 (91%)	37 (92%)	3 (8%)	17	21
3	C	155/156 (99%)	145 (94%)	10 (6%)	21	27
4	D	44/44 (100%)	42 (96%)	2 (4%)	34	46
All	All	394/399 (99%)	358 (91%)	36 (9%)	12	13

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	ASN
1	A	181	ASN
3	C	178	THR
1	A	175	VAL
2	B	2	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	35	HIS
2	B	43	HIS
3	C	78	ASN
2	B	7	ASN
3	C	75	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MAN	A	501	5	11,11,12	0.95	0	14,15,17	1.70	3 (21%)
5	BMA	A	502	5	11,11,12	1.13	1 (9%)	14,15,17	1.73	4 (28%)
5	MAN	A	503	5	11,11,12	0.64	0	14,15,17	1.62	3 (21%)
5	NAG	A	504	5,6	14,14,15	1.02	1 (7%)	15,19,21	0.76	0
5	NAG	A	505	5	14,14,15	0.90	0	15,19,21	1.11	1 (6%)
5	GAL	A	507	5	11,11,12	0.67	0	14,15,17	1.20	0
5	NAG	A	508	5	15,15,15	1.14	2 (13%)	17,21,21	0.71	0
7	MAN	C	511	7	11,11,12	0.58	0	14,15,17	1.59	3 (21%)
7	MAN	C	512	7	11,11,12	0.95	0	14,15,17	2.34	4 (28%)
7	NAG	C	515	7	14,14,15	0.58	0	15,19,21	1.71	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	A	501	5	-	0/2/19/22	0/1/1/1
5	BMA	A	502	5	-	0/2/19/22	0/1/1/1
5	MAN	A	503	5	-	0/2/19/22	0/1/1/1
5	NAG	A	504	5,6	-	0/6/23/26	0/1/1/1
5	NAG	A	505	5	-	0/6/23/26	0/1/1/1
5	GAL	A	507	5	-	0/2/19/22	0/1/1/1
5	NAG	A	508	5	-	0/6/26/26	0/1/1/1
7	MAN	C	511	7	-	0/2/19/22	0/1/1/1
7	MAN	C	512	7	-	0/2/19/22	0/1/1/1
7	NAG	C	515	7	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	502	BMA	C2-C3	2.00	1.55	1.52
5	A	508	NAG	C4-C5	2.42	1.58	1.53
5	A	504	NAG	O4-C4	2.50	1.49	1.43
5	A	508	NAG	O4-C4	2.62	1.49	1.43

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	512	MAN	C6-C5-C4	-6.65	96.60	113.02
7	C	515	NAG	C6-C5-C4	-4.05	103.03	113.02
5	A	501	MAN	C2-C3-C4	-3.46	105.16	111.04
7	C	511	MAN	C6-C5-C4	-3.23	105.06	113.02
5	A	505	NAG	O4-C4-C3	-3.12	103.31	110.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	MAN	1	0
5	A	502	BMA	2	0
5	A	503	MAN	2	0
5	A	504	NAG	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	505	NAG	1	0
5	A	507	GAL	2	0
5	A	508	NAG	1	0
7	C	511	MAN	1	0
7	C	512	MAN	2	0
7	C	515	NAG	6	0

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GAL	A	506	5	11,11,12	0.69	0	14,15,17	1.10	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GAL	A	506	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
6	A	506	GAL	C1-C2-C3	-2.60	106.47	109.54
6	A	506	GAL	C2-C3-C4	-2.14	107.41	111.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	506	GAL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/181 (100%)	-0.63	2 (1%) 82 86	13, 28, 50, 94	0
2	B	47/52 (90%)	-0.58	1 (2%) 67 74	16, 25, 48, 92	0
3	C	180/181 (99%)	-0.57	0 100 100	12, 28, 51, 65	0
4	D	52/52 (100%)	-0.23	3 (5%) 26 35	13, 26, 69, 93	0
All	All	460/466 (98%)	-0.56	6 (1%) 79 84	12, 28, 53, 94	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	52	ASN	7.1
4	D	49	THR	5.3
4	D	50	SER	3.3
2	B	1	GLU	2.8
1	A	77	TYR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	NAG	A	504	14/15	0.61	0.40	5.26	81,89,94,98	0
5	NAG	A	505	14/15	0.88	0.15	2.39	52,57,63,72	0
7	MAN	C	511	11/12	0.88	0.17	1.80	38,46,55,56	0
7	NAG	C	515	14/15	0.87	0.19	1.59	51,65,73,81	0
5	MAN	A	501	11/12	0.93	0.14	1.31	38,42,50,50	0
5	MAN	A	503	11/12	0.59	0.33	-	84,86,90,91	0
5	BMA	A	502	11/12	0.80	0.21	-	56,64,71,78	0
7	MAN	C	512	11/12	0.87	0.21	-	67,82,89,98	0
5	GAL	A	507	11/12	0.62	0.29	-	80,86,91,94	0
5	NAG	A	508	15/15	0.52	0.40	-	81,89,94,98	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GAL	A	506	11/12	0.62	0.40	5.42	90,93,98,99	0
8	CA	A	227	1/1	0.55	0.25	5.18	34,34,34,34	0
8	CA	C	457	1/1	0.88	0.26	4.27	32,32,32,32	0
9	MN	C	458	1/1	0.86	0.08	-1.23	44,44,44,44	0
9	MN	A	228	1/1	0.95	0.04	-2.96	38,38,38,38	0

6.5 Other polymers ⓘ

There are no such residues in this entry.