



Full wwPDB X-ray Structure Validation Report i

Feb 15, 2017 – 01:08 PM EST

PDB ID : 5LP8
Title : Crystal structure of an asymmetric dimer of the ubiquitin ligase HUWE1
Authors : Sander, B.; Lorenz, S.G.
Deposited on : 2016-08-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 13785 atoms, of which 6770 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase HUWE1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	416	Total	C	H	N	O	S	0	0	0
			6693	2170	3285	593	629	16			
1	A	432	Total	C	H	N	O	S	0	0	0
			7060	2280	3485	616	663	16			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	3925	MET	-	initiating methionine	UNP Q7Z6Z7
B	3926	LYS	-	expression tag	UNP Q7Z6Z7
B	3927	HIS	-	expression tag	UNP Q7Z6Z7
B	3928	HIS	-	expression tag	UNP Q7Z6Z7
B	3929	HIS	-	expression tag	UNP Q7Z6Z7
B	3930	HIS	-	expression tag	UNP Q7Z6Z7
B	3931	HIS	-	expression tag	UNP Q7Z6Z7
B	3932	HIS	-	expression tag	UNP Q7Z6Z7
B	3933	PRO	-	expression tag	UNP Q7Z6Z7
B	3934	MET	-	expression tag	UNP Q7Z6Z7
B	3935	SER	-	expression tag	UNP Q7Z6Z7
B	3936	ASP	-	expression tag	UNP Q7Z6Z7
B	3937	TYR	-	expression tag	UNP Q7Z6Z7
B	3938	ASP	-	expression tag	UNP Q7Z6Z7
B	3939	ILE	-	expression tag	UNP Q7Z6Z7
B	3940	PRO	-	expression tag	UNP Q7Z6Z7
B	3941	THR	-	expression tag	UNP Q7Z6Z7
B	3942	THR	-	expression tag	UNP Q7Z6Z7
B	3943	GLU	-	expression tag	UNP Q7Z6Z7
B	3944	ASN	-	expression tag	UNP Q7Z6Z7
B	3945	LEU	-	expression tag	UNP Q7Z6Z7
B	3946	TYR	-	expression tag	UNP Q7Z6Z7
B	3947	PHE	-	expression tag	UNP Q7Z6Z7
B	3948	GLN	-	expression tag	UNP Q7Z6Z7
B	3949	GLY	-	expression tag	UNP Q7Z6Z7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	3950	ALA	-	expression tag	UNP Q7Z6Z7
A	3925	MET	-	initiating methionine	UNP Q7Z6Z7
A	3926	LYS	-	expression tag	UNP Q7Z6Z7
A	3927	HIS	-	expression tag	UNP Q7Z6Z7
A	3928	HIS	-	expression tag	UNP Q7Z6Z7
A	3929	HIS	-	expression tag	UNP Q7Z6Z7
A	3930	HIS	-	expression tag	UNP Q7Z6Z7
A	3931	HIS	-	expression tag	UNP Q7Z6Z7
A	3932	HIS	-	expression tag	UNP Q7Z6Z7
A	3933	PRO	-	expression tag	UNP Q7Z6Z7
A	3934	MET	-	expression tag	UNP Q7Z6Z7
A	3935	SER	-	expression tag	UNP Q7Z6Z7
A	3936	ASP	-	expression tag	UNP Q7Z6Z7
A	3937	TYR	-	expression tag	UNP Q7Z6Z7
A	3938	ASP	-	expression tag	UNP Q7Z6Z7
A	3939	ILE	-	expression tag	UNP Q7Z6Z7
A	3940	PRO	-	expression tag	UNP Q7Z6Z7
A	3941	THR	-	expression tag	UNP Q7Z6Z7
A	3942	THR	-	expression tag	UNP Q7Z6Z7
A	3943	GLU	-	expression tag	UNP Q7Z6Z7
A	3944	ASN	-	expression tag	UNP Q7Z6Z7
A	3945	LEU	-	expression tag	UNP Q7Z6Z7
A	3946	TYR	-	expression tag	UNP Q7Z6Z7
A	3947	PHE	-	expression tag	UNP Q7Z6Z7
A	3948	GLN	-	expression tag	UNP Q7Z6Z7
A	3949	GLY	-	expression tag	UNP Q7Z6Z7
A	3950	ALA	-	expression tag	UNP Q7Z6Z7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total O 2 2	0	0
2	A	30	Total O 30 30	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 ubiquitin-protein ligase HUWE1



- Molecule 1: E3 ubiquitin-protein ligase HUWE1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	177.46 Å 177.46 Å 106.26 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.15 – 2.70 46.15 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.15-2.70) 99.3 (46.15-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.25 (at 2.69 Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R , R_{free}	0.194 , 0.225 0.200 , 0.228	Depositor DCC
R_{free} test set	2549 reflections (4.90%)	DCC
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.035 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13785	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.34	0/3657	0.48	0/4933
1	B	0.50	6/3483 (0.2%)	0.53	0/4695
All	All	0.42	6/7140 (0.1%)	0.50	0/9628

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	3990	ARG	CZ-NH2	7.13	1.42	1.33
1	B	4063	ARG	CZ-NH2	7.05	1.42	1.33
1	B	4071	ARG	CZ-NH2	6.97	1.42	1.33
1	B	3990	ARG	CZ-NH1	6.95	1.42	1.33
1	B	4071	ARG	CZ-NH1	6.85	1.42	1.33
1	B	4063	ARG	CZ-NH1	6.74	1.41	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3575	3485	3484	14	0
1	B	3408	3285	3296	47	0
2	A	30	0	0	0	0
2	B	2	0	0	0	0
All	All	7015	6770	6780	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4027:PHE:HD1	1:B:4069:ILE:CD1	1.45	1.29
1:B:4027:PHE:CD1	1:B:4069:ILE:HD11	1.77	1.17
1:A:3939:ILE:HD11	1:A:3966:LEU:HD23	1.28	1.14
1:B:4073:MET:CE	1:B:4110:VAL:CG2	2.26	1.13
1:B:4027:PHE:HD1	1:B:4069:ILE:HD11	1.07	1.11
1:B:4073:MET:HE2	1:B:4110:VAL:HG21	1.30	1.11
1:B:4073:MET:HE3	1:B:4110:VAL:CG2	1.89	1.01
1:B:4027:PHE:CD1	1:B:4069:ILE:CD1	2.37	1.00
1:A:4134:LYS:HG2	1:A:4244:GLU:HG2	1.48	0.96
1:B:4056:GLN:HB2	1:B:4061:LEU:HD21	1.47	0.96
1:A:3939:ILE:CD1	1:A:3966:LEU:HD23	1.96	0.95
1:B:4073:MET:HE2	1:B:4110:VAL:CG2	1.92	0.94
1:B:4056:GLN:CB	1:B:4061:LEU:HD21	2.00	0.90
1:B:4073:MET:CE	1:B:4110:VAL:HG21	1.92	0.90
1:B:4026:VAL:HG21	1:B:4068:ILE:CG2	2.03	0.89
1:B:4026:VAL:HG21	1:B:4068:ILE:HG21	1.54	0.88
1:B:4057:ASP:O	1:B:4061:LEU:HG	1.77	0.84
1:B:3990:ARG:NH1	1:B:4374:ALA:OXT	2.14	0.81
1:B:4056:GLN:HB2	1:B:4061:LEU:CD2	2.12	0.78
1:B:4073:MET:CE	1:B:4110:VAL:HG23	2.11	0.78
1:B:4073:MET:HE3	1:B:4110:VAL:HG22	1.69	0.74
1:B:3958:PHE:CE2	1:A:3939:ILE:HG23	2.23	0.73
1:B:4198:VAL:O	1:B:4198:VAL:HG13	1.89	0.72
1:A:3939:ILE:CD1	1:A:3966:LEU:CD2	2.69	0.71
1:B:4027:PHE:HD1	1:B:4069:ILE:HD12	1.51	0.69
1:B:3951:ASP:HB3	1:A:3941:THR:HG22	1.75	0.69
1:B:4027:PHE:CE1	1:B:4069:ILE:HD11	2.30	0.66
1:B:4026:VAL:HG21	1:B:4068:ILE:HG22	1.80	0.64
1:B:4056:GLN:HB3	1:B:4061:LEU:HD21	1.79	0.64
1:B:4071:ARG:HH11	1:B:4071:ARG:HG2	1.67	0.60
1:B:4231:TYR:CZ	1:B:4236:LYS:HG3	2.39	0.58
1:B:4231:TYR:CE1	1:B:4236:LYS:HA	2.40	0.57
1:B:3958:PHE:CE2	1:A:3939:ILE:CG2	2.90	0.55
1:B:4063:ARG:HH11	1:B:4063:ARG:HG2	1.73	0.54
1:A:3939:ILE:HD11	1:A:3966:LEU:CD2	2.18	0.52
1:A:4134:LYS:CG	1:A:4244:GLU:HG2	2.30	0.49
1:A:4211:CYS:O	1:A:4215:MET:HG2	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4056:GLN:HB2	1:B:4061:LEU:HD11	1.95	0.47
1:A:4245:GLN:HB3	1:A:4298:GLN:OE1	2.13	0.47
1:A:3937:TYR:CD2	1:A:3999:LYS:HE3	2.49	0.47
1:B:4061:LEU:N	1:B:4061:LEU:HD23	2.30	0.47
1:B:3988:TYR:O	1:B:3991:VAL:HG22	2.14	0.47
1:B:4231:TYR:OH	1:B:4236:LYS:HG3	2.15	0.47
1:B:4027:PHE:CD1	1:B:4069:ILE:HD12	2.36	0.46
1:B:4146:MET:HG2	1:B:4211:CYS:SG	2.56	0.46
1:B:4053:GLU:HG3	1:B:4054:GLU:N	2.32	0.45
1:B:4063:ARG:NH1	1:B:4063:ARG:HG2	2.31	0.44
1:B:4056:GLN:HB2	1:B:4061:LEU:CD1	2.48	0.44
1:B:4066:TYR:OH	1:B:4116:LYS:HG3	2.17	0.44
1:B:4198:VAL:CG1	1:B:4198:VAL:O	2.60	0.43
1:B:4071:ARG:NH1	1:B:4071:ARG:HG2	2.31	0.42
1:B:4073:MET:SD	1:B:4110:VAL:HG23	2.59	0.42
1:B:4243:THR:HG22	1:B:4373:LEU:HG	2.01	0.42
1:B:4048:ILE:HG13	1:B:4065:TRP:CB	2.50	0.41
1:B:4027:PHE:HA	1:B:4069:ILE:HD13	2.01	0.41
1:A:3937:TYR:CD2	1:A:3999:LYS:CE	3.03	0.41
1:A:4141:VAL:HG21	1:A:4215:MET:HG3	2.02	0.41
1:B:4098:HIS:ND1	1:B:4181:PHE:HA	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	428/450 (95%)	421 (98%)	7 (2%)	0	100 100
1	B	410/450 (91%)	398 (97%)	12 (3%)	0	100 100
All	All	838/900 (93%)	819 (98%)	19 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/405 (96%)	386 (100%)	2 (0%)	92	98
1	B	362/405 (89%)	362 (100%)	0	100	100
All	All	750/810 (93%)	748 (100%)	2 (0%)	94	99

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4244	GLU
1	A	4334	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	432/450 (96%)	0.16	6 (1%) 78 77	27, 52, 97, 149	0
1	B	416/450 (92%)	0.52	39 (9%) 11 8	30, 74, 133, 166	0
All	All	848/900 (94%)	0.34	45 (5%) 30 28	27, 61, 124, 166	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4173	THR	5.4
1	B	4204	LYS	5.2
1	B	4203	LYS	5.0
1	B	4209	LEU	4.9
1	B	4168	LEU	4.9
1	B	4205	GLU	4.6
1	B	4201	GLU	4.2
1	B	4184	CYS	3.8
1	B	4199	THR	3.7
1	B	4202	ASN	3.7
1	B	4160	LEU	3.7
1	B	4165	VAL	3.5
1	B	4174	PHE	3.4
1	B	4189	LEU	3.4
1	B	4170	TYR	3.2
1	B	4175	SER	3.1
1	B	4050	PHE	3.0
1	B	4207	VAL	2.9
1	A	4370	GLY	2.8
1	B	4178	VAL	2.8
1	A	4371	PHE	2.8
1	B	4176	THR	2.8
1	B	4211	CYS	2.7
1	A	4054	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	4200	GLU	2.6
1	B	4198	VAL	2.6
1	B	4161	LEU	2.5
1	B	4169	GLY	2.4
1	B	4177	GLU	2.4
1	B	4098	HIS	2.4
1	B	4179	GLN	2.4
1	B	4218	ALA	2.3
1	B	4158	VAL	2.3
1	A	3953	GLN	2.2
1	B	4097	SER	2.2
1	B	4099	CYS	2.2
1	B	4163	ASN	2.2
1	A	3939	ILE	2.2
1	B	4054	GLU	2.2
1	B	4032	ARG	2.2
1	B	4214	ARG	2.1
1	A	3950	ALA	2.1
1	B	4056	GLN	2.0
1	B	4142	ARG	2.0
1	B	4223	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.