



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:44 PM GMT

PDB ID : 1LQT
Title : A covalent modification of NADP⁺ revealed by the atomic resolution structure of FprA, a Mycobacterium tuberculosis oxidoreductase
Authors : Bossi, R.T.; Aliverti, A.; Raimondi, D.; Fischer, F.; Zanetti, G.; Ferrari, D.; Tahallah, N.; Maier, C.S.; Heck, A.J.R.; Rizzi, M.; Mattevi, A.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2002-05-13
Resolution : 1.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1147 (1.12-1.00)
Ramachandran outliers	100387	1086 (1.12-1.00)
Sidechain outliers	100360	1084 (1.12-1.00)
RSRZ outliers	91569	1080 (1.12-1.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	
1	B	456	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	1866	-	-	X	X
2	ACT	A	1869	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	1871	-	-	-	X
2	ACT	A	1872	-	-	-	X
2	ACT	B	1867	-	-	X	X
2	ACT	B	1868	-	-	X	X
2	ACT	B	1873	-	-	X	X
2	ACT	B	1874	-	-	X	-
2	ACT	B	1875	-	-	X	X
2	ACT	B	1876	-	-	X	X
4	ODP	A	2458	X	-	-	-
4	ODP	B	3458	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 20280 atoms, of which 11064 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FprA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	452	Total	C	H	N	O	S	187	23	0
			7214	2261	3629	641	673	10			
1	B	454	Total	C	H	N	O	S	182	12	0
			7073	2222	3554	627	662	8			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	B	1	Total	C	H	O	0	0
			7	2	3	2		
2	B	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	B	1	Total	C	H	O	0	0
			7	2	3	2		
2	B	1	Total	C	H	O	0	0
			7	2	3	2		
2	B	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 86	C 27	H 33	N 9	O 15	P 2	9	0
3	B	1	Total 86	C 27	H 33	N 9	O 15	P 2	9	0

- Molecule 4 is 4-OXO-NICOTINAMIDE-ADENINE DINUCLEOTIDE PHOSPHATE (three-letter code: ODP) (formula: $C_{21}H_{29}N_7O_{18}P_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	10	0
			76	21	27	7	18	3		
4	B	1	Total	C	H	N	O	P	10	0
			76	21	27	7	18	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	971	Total	H	O	1942	0
			2913	1942	971		
5	B	893	Total	H	O	1786	0
			2679	1786	893		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FprA



• Molecule 1: FprA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.33Å 89.22Å 160.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.05 10.00 – 1.05	Depositor EDS
% Data completeness (in resolution range)	95.4 (40.00-1.05) 95.5 (10.00-1.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.05Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.134 , 0.153 0.145 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	9.5	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.49 , 61.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 440131 reflections	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	20280	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ODP, FAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3704	0.87	11/5024 (0.2%)
1	B	0.64	0/3632	0.85	4/4934 (0.1%)
All	All	0.65	0/7336	0.86	15/9958 (0.2%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	441	ARG	NE-CZ-NH1	10.29	125.45	120.30
1	A	33	ASP	CB-CG-OD2	6.68	124.32	118.30
1	A	441	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	B	265	ARG	NE-CZ-NH2	-6.28	117.16	120.30
1	B	401	ASP	CB-CG-OD2	6.28	123.96	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3585	3629	3620	76	0
1	B	3519	3554	3551	67	0
2	A	20	15	15	9	0
2	B	24	18	17	27	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	53	33	31	3	0
3	B	53	33	31	0	0
4	A	49	27	24	4	0
4	B	49	27	24	13	0
5	A	971	1942	0	38	8
5	B	893	1786	0	46	6
All	All	9216	11064	7313	165	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 165 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349[A]:ASN:ND2	1:B:342[A]:PRO:HG3	1.44	1.29
1:A:60[B]:ILE:HG13	5:A:2865:HOH:O	1.42	1.20
1:A:344[B]:VAL:HG23	1:A:349[B]:ASN:ND2	1.57	1.17
1:B:361:LYS:HD2	5:B:4329:HOH:O	1.47	1.14
2:B:1876:ACT:H2	4:B:3458:ODP:O1A	1.43	1.12

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1868:ACT:OXT	5:A:3320:HOH:O[1_565]	0.65	1.55
2:B:1868:ACT:C	5:A:3320:HOH:O[1_565]	1.82	0.38
5:A:3325:HOH:O	5:B:3803:HOH:O[3_545]	1.84	0.36
5:A:2970:HOH:O	5:A:3396:HOH:O[4_445]	1.99	0.21
5:A:2684:HOH:O	5:B:4333:HOH:O[1_545]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/456 (103%)	461 (98%)	9 (2%)	1 (0%)	52	19
1	B	464/456 (102%)	452 (97%)	11 (2%)	1 (0%)	52	19
All	All	935/912 (102%)	913 (98%)	20 (2%)	2 (0%)	52	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	TRP
1	B	262	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/365 (105%)	383 (100%)	1 (0%)	94	76
1	B	377/365 (103%)	375 (100%)	2 (0%)	92	70
All	All	761/730 (104%)	758 (100%)	3 (0%)	93	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	TYR
1	B	132	TYR
1	B	435	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	405	GLN
1	B	264	HIS
1	B	388	ASN
1	A	402	HIS
1	B	385	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	1866	-	1,3,3	0.29	0	0,3,3	0.00	-
2	ACT	A	1869	-	1,3,3	0.11	0	0,3,3	0.00	-
2	ACT	A	1870	-	1,3,3	0.50	0	0,3,3	0.00	-
2	ACT	A	1871	-	1,3,3	1.67	0	0,3,3	0.00	-
2	ACT	A	1872	-	1,3,3	1.35	0	0,3,3	0.00	-
3	FAD	A	2457	-	48,58,58	0.92	2 (4%)	53,89,89	1.23	6 (11%)
4	ODP	A	2458	-	41,53,53	1.07	2 (4%)	48,82,82	1.29	4 (8%)
2	ACT	B	1867	-	1,3,3	0.03	0	0,3,3	0.00	-
2	ACT	B	1868	-	1,3,3	0.53	0	0,3,3	0.00	-
2	ACT	B	1873	1	1,3,3	0.83	0	0,3,3	0.00	-
2	ACT	B	1874	-	1,3,3	2.38	1 (100%)	0,3,3	0.00	-
2	ACT	B	1875	-	1,3,3	0.62	0	0,3,3	0.00	-
2	ACT	B	1876	4	1,3,3	1.25	0	0,3,3	0.00	-
3	FAD	B	3457	-	48,58,58	0.91	3 (6%)	53,89,89	1.20	4 (7%)
4	ODP	B	3458	2	41,53,53	0.83	2 (4%)	48,82,82	1.27	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	1866	-	-	0/0/0/0	0/0/0/0
2	ACT	A	1869	-	-	0/0/0/0	0/0/0/0
2	ACT	A	1870	-	-	0/0/0/0	0/0/0/0
2	ACT	A	1871	-	-	0/0/0/0	0/0/0/0
2	ACT	A	1872	-	-	0/0/0/0	0/0/0/0
3	FAD	A	2457	-	-	0/30/50/50	0/6/6/6
4	ODP	A	2458	-	1/1/18/18	0/26/80/80	0/4/5/5
2	ACT	B	1867	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1868	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1873	1	-	0/0/0/0	0/0/0/0
2	ACT	B	1874	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1875	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1876	4	-	0/0/0/0	0/0/0/0
3	FAD	B	3457	-	-	0/30/50/50	0/6/6/6
4	ODP	B	3458	2	1/1/18/18	0/26/80/80	0/4/5/5

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2457	FAD	C6-C5X	-2.63	1.37	1.41
2	B	1874	ACT	CH3-C	-2.38	1.45	1.48
4	B	3458	ODP	O4D-C1D	-2.01	1.39	1.41
4	A	2458	ODP	O4D-C1D	2.25	1.43	1.41
3	B	3457	FAD	C2A-N3A	2.27	1.36	1.32

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2458	ODP	N3A-C2A-N1A	-5.14	124.96	128.89
4	B	3458	ODP	N3A-C2A-N1A	-4.16	125.71	128.89
3	B	3457	FAD	N3A-C2A-N1A	-3.60	126.14	128.89
4	B	3458	ODP	C4B-O4B-C1B	-3.31	106.08	109.72
3	A	2457	FAD	N3A-C2A-N1A	-3.04	126.57	128.89

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	2458	ODP	C3N

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
4	B	3458	ODP	C3N

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 41 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1866	ACT	7	0
2	A	1870	ACT	1	0
2	A	1872	ACT	1	0
3	A	2457	FAD	3	0
4	A	2458	ODP	4	0
2	B	1867	ACT	2	0
2	B	1868	ACT	0	2
2	B	1873	ACT	2	0
2	B	1874	ACT	4	0
2	B	1875	ACT	7	0
2	B	1876	ACT	12	0
4	B	3458	ODP	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/456 (99%)	-0.09	12 (2%) 58 50	8, 12, 22, 36	0
1	B	454/456 (99%)	-0.00	21 (4%) 36 31	7, 13, 24, 36	0
All	All	906/912 (99%)	-0.05	33 (3%) 46 39	7, 13, 23, 36	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	393	ALA	8.2
1	B	263	GLY	7.7
1	B	282	ARG	6.3
1	B	231	ASP	5.8
1	B	281	LYS	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACT	B	1876	4/4	0.79	0.38	35.47	25,30,30,31	0
2	ACT	B	1875	4/4	0.77	0.31	20.71	27,30,30,31	0
2	ACT	B	1868	4/4	0.90	0.12	14.41	19,20,20,21	0
2	ACT	A	1869	4/4	0.72	0.23	10.95	27,29,29,29	0
2	ACT	A	1866	4/4	0.79	0.36	10.79	25,27,28,28	0
2	ACT	B	1867	4/4	0.91	0.20	4.15	14,16,24,24	0
2	ACT	A	1872	4/4	0.74	0.19	4.09	24,27,27,27	0
2	ACT	A	1871	4/4	0.94	0.09	3.55	15,16,16,18	0
2	ACT	B	1873	4/4	0.21	0.28	3.49	47,47,47,48	0
3	FAD	B	3457	53/53	0.98	0.08	1.87	4,7,9,10	13
2	ACT	A	1870	4/4	0.80	0.19	1.70	29,31,31,33	0
2	ACT	B	1874	4/4	0.92	0.08	1.16	15,16,18,18	0
3	FAD	A	2457	53/53	0.98	0.09	0.91	5,8,10,12	13
4	ODP	B	3458	49/49	0.99	0.07	0.31	7,10,13,14	10
4	ODP	A	2458	49/49	0.99	0.06	-0.22	6,10,13,16	10

6.5 Other polymers ⓘ

There are no such residues in this entry.