



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:32 PM GMT

PDB ID : 4LW2  
Title : Structural changes during cysteine desulfurase CsdA and sulfur-acceptor CsdE interactions provide insight into the trans-persulfuration  
Authors : Kim, S.; Park, S.Y.  
Deposited on : 2013-07-26  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	<div> <div>2%</div> <div>86% 11% ..</div> </div>
1	B	404	<div> <div>2%</div> <div>84% 13% ..</div> </div>
1	C	404	<div> <div>2%</div> <div>83% 14% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	502	-	-	X	X
3	GOL	B	502	-	-	X	X
3	GOL	C	502	-	-	X	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine sulfinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	2	0
			3043	1930	531	570	12			
1	B	399	Total	C	N	O	S	0	3	0
			3046	1932	529	573	12			
1	C	398	Total	C	N	O	S	0	2	0
			3033	1925	526	570	12			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q46925
A	-1	SER	-	EXPRESSION TAG	UNP Q46925
A	0	HIS	-	EXPRESSION TAG	UNP Q46925
B	-2	GLY	-	EXPRESSION TAG	UNP Q46925
B	-1	SER	-	EXPRESSION TAG	UNP Q46925
B	0	HIS	-	EXPRESSION TAG	UNP Q46925
C	-2	GLY	-	EXPRESSION TAG	UNP Q46925
C	-1	SER	-	EXPRESSION TAG	UNP Q46925
C	0	HIS	-	EXPRESSION TAG	UNP Q46925

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	386	Total	O	0	0
			386	386		
4	B	363	Total	O	0	0
			363	363		
4	C	358	Total	O	0	0
			358	358		



- Molecule 1: Cysteine sulfinatase desulfinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	168.30 Å   91.44 Å   83.89 Å 90.00°   97.26°   90.00°	Depositor
Resolution (Å)	50.00 – 1.80 37.89 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-1.80) 99.2 (37.89-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.56 (at 1.79 Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.165   ,   0.212 0.169   ,   0.213	Depositor DCC
$R_{free}$ test set	5817 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.9	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 116265 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10292	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.33	7/3116 (0.2%)	1.01	2/4245 (0.0%)
1	B	1.28	9/3122 (0.3%)	1.03	6/4254 (0.1%)
1	C	1.26	2/3106 (0.1%)	1.02	1/4232 (0.0%)
All	All	1.29	18/9344 (0.2%)	1.02	9/12731 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	267	TRP	CE3-CZ3	5.96	1.48	1.38
1	B	195	VAL	CB-CG1	-5.89	1.40	1.52
1	B	379	TYR	CD1-CE1	5.80	1.48	1.39
1	B	288	TRP	CB-CG	5.78	1.60	1.50
1	A	204	PHE	CG-CD1	5.78	1.47	1.38

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	60	ARG	NE-CZ-NH2	-8.01	116.30	120.30
1	B	246	LEU	CB-CG-CD2	5.60	120.53	111.00
1	C	151	ASP	CB-CG-OD2	-5.47	113.38	118.30
1	B	322	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	B	24	ASP	CB-CG-OD2	-5.31	113.52	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3043	0	3010	39	0
1	B	3046	0	3009	45	0
1	C	3033	0	2997	51	0
2	A	15	0	6	2	0
2	B	15	0	7	0	0
2	C	15	0	6	1	0
3	A	6	0	7	5	0
3	B	6	0	8	8	0
3	C	6	0	8	4	0
4	A	386	0	0	6	0
4	B	363	0	0	8	0
4	C	358	0	0	12	0
All	All	10292	0	9058	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:8:GLN:HG2	4:C:688:HOH:O	1.52	1.07
1:C:17:GLN:HG3	4:C:854:HOH:O	1.53	1.05
1:B:373:ARG:HH22	3:B:502:GOL:H31	1.22	1.02
1:C:66:GLU:HG2	4:C:791:HOH:O	1.62	0.97
1:A:261:THR:HG23	4:A:756:HOH:O	1.66	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/404 (99%)	388 (97%)	11 (3%)	0	100	100
1	B	400/404 (99%)	386 (96%)	13 (3%)	1 (0%)	46	29
1	C	398/404 (98%)	390 (98%)	8 (2%)	0	100	100
All	All	1197/1212 (99%)	1164 (97%)	32 (3%)	1 (0%)	56	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	159	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/317 (99%)	304 (96%)	11 (4%)	43	25
1	B	316/317 (100%)	308 (98%)	8 (2%)	55	39
1	C	314/317 (99%)	300 (96%)	14 (4%)	34	16
All	All	945/951 (99%)	912 (96%)	33 (4%)	43	25

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	246	LEU
1	B	379	TYR
1	C	375	SER
1	B	250	LYS
1	B	252	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	77	ASN
1	B	122	ASN
1	C	324	GLN
1	B	99	GLN
1	B	131	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PLP	A	501	1	15,15,16	3.15	5 (33%)	21,22,23	2.46	11 (52%)
3	GOL	A	502	-	5,5,5	1.50	1 (20%)	5,5,5	1.51	1 (20%)
2	PLP	B	501	1	15,15,16	2.19	3 (20%)	21,22,23	2.01	5 (23%)
3	GOL	B	502	-	5,5,5	1.13	0	5,5,5	1.97	1 (20%)
2	PLP	C	501	1	15,15,16	2.62	5 (33%)	21,22,23	1.71	7 (33%)
3	GOL	C	502	-	5,5,5	1.14	0	5,5,5	2.42	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1
3	GOL	A	502	-	-	0/4/4/4	0/0/0/0
2	PLP	B	501	1	-	0/6/6/8	0/1/1/1
3	GOL	B	502	-	-	0/4/4/4	0/0/0/0
2	PLP	C	501	1	-	0/6/6/8	0/1/1/1
3	GOL	C	502	-	-	0/4/4/4	0/0/0/0

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	PLP	C4A-C4	-9.12	1.33	1.51
2	C	501	PLP	C4A-C4	-6.70	1.38	1.51
2	B	501	PLP	C4A-C4	-5.83	1.39	1.51
2	B	501	PLP	C2A-C2	-3.64	1.43	1.50
2	A	501	PLP	O3-C3	-3.01	1.30	1.37

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PLP	C5-C6-N1	-4.50	116.04	123.86
2	A	501	PLP	C3-C2-N1	-3.81	115.35	120.61
3	C	502	GOL	O2-C2-C3	-3.61	92.09	108.65
2	C	501	PLP	C3-C2-N1	-2.98	116.50	120.61
2	A	501	PLP	C4A-C4-C5	-2.97	117.78	120.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PLP	2	0
3	A	502	GOL	5	0
3	B	502	GOL	8	0
2	C	501	PLP	1	0
3	C	502	GOL	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	399/404 (98%)	-0.31	4 (1%) 84 82	11, 19, 32, 49	0
1	B	399/404 (98%)	-0.19	8 (2%) 68 64	12, 21, 37, 49	0
1	C	398/404 (98%)	-0.08	10 (2%) 61 56	12, 22, 38, 45	0
All	All	1196/1212 (98%)	-0.19	22 (1%) 71 67	11, 21, 36, 49	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	400	VAL	3.6
1	C	159	PRO	3.4
1	C	160	ARG	3.1
1	A	155	GLU	3.1
1	C	337	HIS	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	C	502	6/6	0.85	0.28	4.65	34,37,39,42	0
3	GOL	A	502	6/6	0.84	0.21	3.85	21,27,35,37	0
3	GOL	B	502	6/6	0.88	0.20	3.35	27,32,33,40	0
2	PLP	A	501	15/16	0.99	0.12	-0.28	12,14,16,17	0
2	PLP	B	501	15/16	0.99	0.09	-1.24	13,15,18,18	0
2	PLP	C	501	15/16	0.98	0.10	-1.59	13,18,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.