



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:27 AM GMT

PDB ID : 3LY8
Title : Crystal structure of mutant D471E of the periplasmic domain of CadC
Authors : Eichinger, A.; Skerra, A.
Deposited on : 2010-02-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional activator cadC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2566	1628	435	493	10	149	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	GLY	-	EXPRESSION TAG	UNP P23890
A	142	SER	-	EXPRESSION TAG	UNP P23890
A	143	GLY	-	EXPRESSION TAG	UNP P23890
A	144	MET	-	EXPRESSION TAG	UNP P23890
A	145	LYS	-	EXPRESSION TAG	UNP P23890
A	146	GLU	-	EXPRESSION TAG	UNP P23890
A	147	THR	-	EXPRESSION TAG	UNP P23890
A	148	ALA	-	EXPRESSION TAG	UNP P23890
A	149	ALA	-	EXPRESSION TAG	UNP P23890
A	150	ALA	-	EXPRESSION TAG	UNP P23890
A	151	LYS	-	EXPRESSION TAG	UNP P23890
A	152	PHE	-	EXPRESSION TAG	UNP P23890
A	153	GLU	-	EXPRESSION TAG	UNP P23890
A	154	ARG	-	EXPRESSION TAG	UNP P23890
A	155	GLN	-	EXPRESSION TAG	UNP P23890
A	156	HIS	-	EXPRESSION TAG	UNP P23890
A	157	MET	-	EXPRESSION TAG	UNP P23890
A	158	ASP	-	EXPRESSION TAG	UNP P23890
A	159	SER	-	EXPRESSION TAG	UNP P23890
A	160	PRO	-	EXPRESSION TAG	UNP P23890
A	161	ASP	-	EXPRESSION TAG	UNP P23890
A	162	LEU	-	EXPRESSION TAG	UNP P23890
A	163	GLY	-	EXPRESSION TAG	UNP P23890
A	164	THR	-	EXPRESSION TAG	UNP P23890
A	165	ASP	-	EXPRESSION TAG	UNP P23890
A	166	ASP	-	EXPRESSION TAG	UNP P23890
A	167	ASP	-	EXPRESSION TAG	UNP P23890

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	ASP	-	EXPRESSION TAG	UNP P23890
A	169	LYS	-	EXPRESSION TAG	UNP P23890
A	170	ALA	-	EXPRESSION TAG	UNP P23890
A	171	MET	-	EXPRESSION TAG	UNP P23890
A	172	ALA	-	EXPRESSION TAG	UNP P23890
A	173	HIS	-	EXPRESSION TAG	UNP P23890
A	174	HIS	-	EXPRESSION TAG	UNP P23890
A	175	HIS	-	EXPRESSION TAG	UNP P23890
A	176	HIS	-	EXPRESSION TAG	UNP P23890
A	177	HIS	-	EXPRESSION TAG	UNP P23890
A	178	HIS	-	EXPRESSION TAG	UNP P23890
A	179	SER	-	EXPRESSION TAG	UNP P23890
A	180	SER	-	EXPRESSION TAG	UNP P23890
A	181	GLY	-	EXPRESSION TAG	UNP P23890
A	182	HIS	-	EXPRESSION TAG	UNP P23890
A	183	ILE	-	EXPRESSION TAG	UNP P23890
A	184	GLU	-	EXPRESSION TAG	UNP P23890
A	185	GLY	-	EXPRESSION TAG	UNP P23890
A	186	ARG	-	EXPRESSION TAG	UNP P23890
A	187	HIS	-	EXPRESSION TAG	UNP P23890
A	471	GLU	ASP	ENGINEERED	UNP P23890


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	234	Total O 234 234	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional activator cadC



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	83.63 Å 83.63 Å 199.49 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	72.43 – 1.90 34.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (72.43-1.90) 99.9 (34.95-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.81 (at 1.89 Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.202 , 0.231 0.193 , 0.224	Depositor DCC
R_{free} test set	1684 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 33530 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2800	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/2616	0.64	0/3552

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2566	0	2551	14	0
2	A	234	0	0	1	0
All	All	2800	0	2551	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:ARG:HH12	1:A:484:ASN:HD22	1.31	0.79
1:A:384:ILE:HD12	1:A:425:VAL:HG12	1.74	0.69
1:A:384:ILE:CD1	1:A:425:VAL:HG12	2.22	0.69
1:A:192:ILE:HD11	1:A:252:SER:HA	1.75	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:THR:O	1:A:296:ASN:O	2.14	0.65
1:A:318:TRP:CE3	1:A:322:MET:SD	3.06	0.49
1:A:471:GLU:HG2	2:A:6:HOH:O	2.12	0.49
1:A:247:ILE:HD11	1:A:249:GLU:HB3	1.95	0.49
1:A:192:ILE:HD11	1:A:252:SER:CA	2.44	0.45
1:A:373:THR:HG22	1:A:413:LEU:HD11	1.99	0.44
1:A:334:GLY:HA2	1:A:337:LEU:HD12	1.98	0.44
1:A:292:TYR:CD1	1:A:302:ILE:HD12	2.53	0.43
1:A:204:VAL:HG13	1:A:260:ALA:HB2	2.00	0.42
1:A:295:THR:O	1:A:296:ASN:C	2.58	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/372 (86%)	310 (97%)	5 (2%)	6 (2%)	10 2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	SER
1	A	213	SER
1	A	296	ASN
1	A	209	ASN
1	A	212	SER
1	A	211	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	285/324 (88%)	276 (97%)	9 (3%)	46 35

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	191	ARG
1	A	192	ILE
1	A	204	VAL
1	A	211	TRP
1	A	247	ILE
1	A	296	ASN
1	A	384	ILE
1	A	471	GLU
1	A	486	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	484	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	315/372 (84%)	0.04	11 (3%)	48 51	11, 20, 35, 45	26 (8%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	215	TYR	4.9
1	A	204	VAL	4.1
1	A	246	ASN	3.6
1	A	269	ALA	3.6
1	A	296	ASN	3.1
1	A	511	SER	3.0
1	A	510	ALA	2.7
1	A	247	ILE	2.6
1	A	205	ASN	2.5
1	A	297	GLY	2.3
1	A	261	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.