



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:47 PM GMT

PDB ID : 1LZS
Title : STRUCTURAL CHANGES OF THE ACTIVE SITE CLEFT AND DIFFERENT SACCHARIDE BINDING MODES IN HUMAN LYSOZYME CO-CRYSTALLIZED WITH HEXA-N-ACETYL-CHITOHXAOSE AT PH 4.0
Authors : Matsushima, M.; Song, H.
Deposited on : 1994-09-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	<div> <div>2%</div> <div>73%</div> <div>20%</div> <div>.</div> </div>
1	B	130	<div> <div>78%</div> <div>21%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	136	-	-	-	X
2	NAG	A	139	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	139	-	-	-	X
3	NAG	A	140	-	-	-	X
3	NAG	A	141	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	130	Total	C	N	O	S	0	0	0
			1029	633	200	186	10			
1	B	130	Total	C	N	O	S	0	0	0
			1029	633	200	186	10			

- Molecule 2 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	4	Total	C	N	O	0	0
			57	32	4	21		
2	B	4	Total	C	N	O	0	0
			57	32	4	21		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			29	16	2	11		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	129	Total	O	0	0
			129	129		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	142	Total	O	0	0
			142	142		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HUMAN LYSOZYME



• Molecule 1: HUMAN LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.99 Å 162.25 Å 32.52 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.60 6.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-1.60) 72.3 (6.00-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.170 , (Not available) 0.163 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	13.9	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.65 , 95.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ¹	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 23982 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2474	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1049	1.68	18/1416 (1.3%)
1	B	0.92	1/1049 (0.1%)	1.81	22/1416 (1.6%)
All	All	0.89	1/2098 (0.0%)	1.74	40/2832 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	10	ARG	CD-NE	-5.22	1.37	1.46

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	10	ARG	NE-CZ-NH1	20.16	130.38	120.30
1	A	41	ARG	NE-CZ-NH1	14.42	127.51	120.30
1	B	10	ARG	NE-CZ-NH2	-14.39	113.11	120.30
1	A	5	ARG	NE-CZ-NH1	13.48	127.04	120.30
1	B	10	ARG	CD-NE-CZ	12.04	140.45	123.60
1	B	5	ARG	NE-CZ-NH2	-10.83	114.89	120.30
1	B	107	ARG	NE-CZ-NH1	9.19	124.89	120.30
1	B	119	ARG	NE-CZ-NH2	-8.52	116.04	120.30
1	B	122	ARG	NE-CZ-NH1	8.37	124.48	120.30
1	B	18	ASP	CB-CG-OD2	-8.29	110.84	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	5	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	B	115	ARG	NE-CZ-NH1	8.09	124.34	120.30
1	A	113	ARG	NE-CZ-NH1	8.08	124.34	120.30
1	A	62	ARG	NE-CZ-NH1	8.08	124.34	120.30
1	A	41	ARG	NE-CZ-NH2	-8.01	116.29	120.30
1	B	10	ARG	CG-CD-NE	7.78	128.13	111.80
1	B	18	ASP	CB-CG-OD1	7.31	124.88	118.30
1	B	98	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	A	104	GLN	CB-CG-CD	7.10	130.05	111.60
1	A	5	ARG	NE-CZ-NH2	-6.91	116.84	120.30
1	A	107	ARG	NE-CZ-NH1	6.69	123.65	120.30
1	B	115	ARG	NE-CZ-NH2	-6.57	117.02	120.30
1	A	21	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	50	ARG	NE-CZ-NH1	6.25	123.42	120.30
1	A	115	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	A	10	ARG	NE-CZ-NH2	-5.91	117.35	120.30
1	A	87	ASP	CB-CG-OD1	5.86	123.58	118.30
1	B	113	ARG	NE-CZ-NH2	-5.83	117.39	120.30
1	A	122	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	B	107	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	B	101	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	A	36	SER	N-CA-CB	-5.51	102.24	110.50
1	B	53	ASP	CB-CG-OD2	-5.48	113.36	118.30
1	B	14	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	63	TYR	N-CA-C	5.41	125.60	111.00
1	A	115	ARG	CD-NE-CZ	-5.33	116.14	123.60
1	B	124	TYR	CB-CG-CD1	-5.22	117.87	121.00
1	B	63	TYR	CB-CG-CD1	-5.15	117.91	121.00
1	B	7	GLU	CG-CD-OE1	5.10	128.50	118.30
1	A	107	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	41	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1029	0	992	19	1
1	B	1029	0	992	12	0
2	A	57	0	51	0	0
2	B	57	0	51	0	0
3	A	29	0	27	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	129	0	0	2	0
5	B	142	0	0	0	1
All	All	2474	0	2113	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ARG:HE	1:B:126:GLN:NE2	1.67	0.91
1:A:48:GLY:O	1:A:49:ASP:HB3	1.85	0.75
1:B:122:ARG:HE	1:B:126:GLN:HE22	1.35	0.74
1:B:50:ARG:NH1	1:B:71:PRO:HG3	2.11	0.65
1:A:64:TRP:O	1:A:77:CYS:HB2	1.96	0.65
1:A:23:ILE:HD13	1:A:106:ILE:HD12	1.82	0.62
1:A:110:VAL:HG22	1:A:113:ARG:NH2	2.17	0.60
1:A:104:GLN:HB2	5:A:452:HOH:O	2.07	0.54
1:A:23:ILE:HD12	1:A:100:VAL:HG11	1.92	0.52
1:A:23:ILE:CD1	1:A:100:VAL:HG11	2.40	0.51
1:B:122:ARG:HE	1:B:126:GLN:HE21	1.56	0.50
1:A:69:LYS:HE2	5:A:552:HOH:O	2.11	0.50
1:A:49:ASP:O	1:A:62:ARG:NH1	2.35	0.49
1:A:123:GLN:O	1:A:126:GLN:HG3	2.14	0.48
1:A:45:TYR:CE2	1:A:47:ALA:HA	2.49	0.48
1:A:58:GLN:OE1	3:A:140:NAG:H83	2.15	0.46
1:B:122:ARG:NE	1:B:126:GLN:NE2	2.49	0.46
1:A:62:ARG:NH2	1:A:71:PRO:O	2.48	0.46
1:A:64:TRP:CD2	1:A:99:VAL:HG22	2.51	0.46
1:A:45:TYR:CE1	1:A:50:ARG:HA	2.51	0.45
1:B:64:TRP:CE3	1:B:76:ALA:HB3	2.52	0.45
1:B:50:ARG:HH11	1:B:50:ARG:HD3	1.58	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:ASP:OD2	1:B:62:ARG:HD2	2.17	0.44
1:B:112:TRP:CD1	1:B:116:CYS:HB2	2.54	0.42
1:A:67:ASP:HB3	1:A:81:CYS:SG	2.60	0.41
1:A:52:THR:HB	1:A:54:TYR:CE1	2.55	0.41
1:A:21:ARG:HD2	1:B:50:ARG:NH2	2.35	0.41
1:B:13:LYS:NZ	1:B:18:ASP:OD2	2.54	0.41
1:A:64:TRP:CE2	1:A:99:VAL:HG22	2.55	0.41
1:B:12:LEU:HB2	1:B:25:LEU:HD11	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:GLN:CB	5:B:456:HOH:O[1_554]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/130 (98%)	124 (97%)	3 (2%)	1 (1%)	24	6
1	B	128/130 (98%)	123 (96%)	5 (4%)	0	100	100
All	All	256/260 (98%)	247 (96%)	8 (3%)	1 (0%)	39	17

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/105 (100%)	98 (93%)	7 (7%)	20	4
1	B	105/105 (100%)	103 (98%)	2 (2%)	65	39
All	All	210/210 (100%)	201 (96%)	9 (4%)	35	11

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ARG
1	A	49	ASP
1	A	50	ARG
1	A	62	ARG
1	A	69	LYS
1	A	100	VAL
1	A	102	ASP
1	B	100	VAL
1	B	102	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	B	78	HIS
1	B	86	GLN
1	B	117	GLN
1	B	126	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	136	2	15,15,15	0.65	0	17,21,21	1.51	2 (11%)
2	NAG	A	137	2	13,14,15	0.60	0	14,19,21	1.28	1 (7%)
2	NAG	A	138	2	13,14,15	0.83	0	14,19,21	1.52	4 (28%)
2	NAG	A	139	2	13,14,15	1.07	1 (7%)	14,19,21	1.56	2 (14%)
3	NAG	A	140	3	15,15,15	0.79	0	17,21,21	1.40	2 (11%)
3	NAG	A	141	3	13,14,15	0.64	0	14,19,21	1.72	3 (21%)
2	NAG	B	136	2	15,15,15	0.73	0	17,21,21	2.34	6 (35%)
2	NAG	B	137	2	13,14,15	0.48	0	14,19,21	1.12	0
2	NAG	B	138	2	13,14,15	0.71	0	14,19,21	1.34	3 (21%)
2	NAG	B	139	2	13,14,15	0.66	0	14,19,21	1.43	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	136	2	-	0/6/26/26	0/1/1/1
2	NAG	A	137	2	-	0/6/22/26	0/1/1/1
2	NAG	A	138	2	-	0/6/22/26	0/1/1/1
2	NAG	A	139	2	-	0/6/22/26	0/1/1/1
3	NAG	A	140	3	-	0/6/26/26	0/1/1/1
3	NAG	A	141	3	-	0/6/22/26	0/1/1/1
2	NAG	B	136	2	-	0/6/26/26	0/1/1/1
2	NAG	B	137	2	-	0/6/22/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	138	2	-	0/6/22/26	0/1/1/1
2	NAG	B	139	2	-	0/6/22/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	139	NAG	C1-C2	-2.58	1.50	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	136	NAG	O3-C3-C4	-3.07	103.43	110.34
2	B	138	NAG	O7-C7-C8	-3.03	116.49	122.06
2	A	136	NAG	O3-C3-C4	-2.92	103.76	110.34
2	A	137	NAG	O5-C5-C4	-2.86	104.64	109.97
2	B	136	NAG	O3-C3-C2	-2.84	103.87	109.66
2	A	138	NAG	C1-O5-C5	-2.78	108.34	113.47
2	A	138	NAG	O7-C7-C8	-2.53	117.42	122.06
2	B	139	NAG	C3-C4-C5	-2.48	106.38	111.40
2	B	139	NAG	O1-C1-O5	-2.31	103.94	110.25
2	B	136	NAG	C1-O5-C5	-2.19	109.42	113.47
2	A	138	NAG	O1-C1-O5	-2.18	104.28	110.25
2	A	139	NAG	C3-C4-C5	-2.08	107.19	111.40
2	A	138	NAG	O7-C7-N2	2.04	126.03	121.86
3	A	141	NAG	O6-C6-C5	2.18	118.03	111.66
3	A	140	NAG	O7-C7-C8	2.19	126.08	122.06
2	B	138	NAG	O7-C7-N2	2.24	126.44	121.86
2	B	138	NAG	C4-C5-C6	2.26	116.55	112.38
2	B	136	NAG	O4-C4-C3	2.54	116.06	110.34
2	B	139	NAG	O5-C5-C6	2.96	110.24	106.62
2	B	136	NAG	C3-C2-N2	3.20	117.28	110.66
3	A	141	NAG	O5-C5-C6	3.75	111.21	106.62
2	A	139	NAG	C1-O5-C5	3.75	120.38	113.47
3	A	140	NAG	C4-C3-C2	3.97	115.93	110.43
3	A	141	NAG	C4-C3-C2	3.98	115.91	111.47
2	A	136	NAG	C4-C3-C2	4.31	116.40	110.43
2	B	136	NAG	C4-C3-C2	6.68	119.70	110.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	140	NAG	1	0

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	130/130 (100%)	-0.35	3 (2%) 64 62	11, 18, 31, 38	0
1	B	130/130 (100%)	-0.64	0 100 100	9, 14, 22, 28	0
All	All	260/260 (100%)	-0.50	3 (1%) 81 81	9, 16, 28, 38	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	VAL	2.4
1	A	48	GLY	2.2
1	A	126	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	141	14/15	0.79	0.20	17.62	34,36,37,37	0
2	NAG	A	136	15/15	0.80	0.18	8.75	28,33,36,36	0
2	NAG	A	139	14/15	0.84	0.20	4.23	30,32,36,37	0
3	NAG	A	140	15/15	0.82	0.17	4.06	35,36,37,38	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	B	139	14/15	0.94	0.11	2.89	14,17,21,22	0
2	NAG	B	136	15/15	0.87	0.14	0.81	26,30,32,33	0
2	NAG	A	138	14/15	0.96	0.07	0.35	14,18,23,24	0
2	NAG	B	138	14/15	0.97	0.07	-0.03	12,14,16,18	0
2	NAG	B	137	14/15	0.94	0.08	-	17,20,24,24	0
2	NAG	A	137	14/15	0.94	0.09	-	21,24,27,27	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NA	B	131	1/1	0.99	0.04	-1.14	18,18,18,18	0
4	NA	A	131	1/1	0.99	0.03	-1.61	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.