



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:27 AM GMT

PDB ID : 3M1Y  
Title : Crystal Structure of a Phosphoserine phosphatase (SerB) from *Helicobacter pylori*  
Authors : Syed Ibrahim, B.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2010-03-05  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*





The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	 65% 26% • 6%
1	B	217	 61% 31% • 5%
1	C	217	 55% 36% 5% •
1	D	217	 51% 40% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	C	301	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoserine phosphatase (SerB).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	Se	0	0	0
			1580	1015	267	290	3	5			
1	B	206	Total	C	N	O	S	Se	0	0	0
			1599	1026	270	294	3	6			
1	C	208	Total	C	N	O	S	Se	0	0	0
			1611	1035	273	294	3	6			
1	D	206	Total	C	N	O	S	Se	0	0	0
			1598	1024	271	294	3	6			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP O25367
A	0	SER	-	expression tag	UNP O25367
A	1	LEU	-	expression tag	UNP O25367
A	208	GLU	-	expression tag	UNP O25367
A	209	GLY	-	expression tag	UNP O25367
A	210	HIS	-	expression tag	UNP O25367
A	211	HIS	-	expression tag	UNP O25367
A	212	HIS	-	expression tag	UNP O25367
A	213	HIS	-	expression tag	UNP O25367
A	214	HIS	-	expression tag	UNP O25367
A	215	HIS	-	expression tag	UNP O25367
B	-1	MSE	-	expression tag	UNP O25367
B	0	SER	-	expression tag	UNP O25367
B	1	LEU	-	expression tag	UNP O25367
B	208	GLU	-	expression tag	UNP O25367
B	209	GLY	-	expression tag	UNP O25367
B	210	HIS	-	expression tag	UNP O25367
B	211	HIS	-	expression tag	UNP O25367
B	212	HIS	-	expression tag	UNP O25367
B	213	HIS	-	expression tag	UNP O25367
B	214	HIS	-	expression tag	UNP O25367

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	215	HIS	-	expression tag	UNP O25367
C	-1	MSE	-	expression tag	UNP O25367
C	0	SER	-	expression tag	UNP O25367
C	1	LEU	-	expression tag	UNP O25367
C	208	GLU	-	expression tag	UNP O25367
C	209	GLY	-	expression tag	UNP O25367
C	210	HIS	-	expression tag	UNP O25367
C	211	HIS	-	expression tag	UNP O25367
C	212	HIS	-	expression tag	UNP O25367
C	213	HIS	-	expression tag	UNP O25367
C	214	HIS	-	expression tag	UNP O25367
C	215	HIS	-	expression tag	UNP O25367
D	-1	MSE	-	expression tag	UNP O25367
D	0	SER	-	expression tag	UNP O25367
D	1	LEU	-	expression tag	UNP O25367
D	208	GLU	-	expression tag	UNP O25367
D	209	GLY	-	expression tag	UNP O25367
D	210	HIS	-	expression tag	UNP O25367
D	211	HIS	-	expression tag	UNP O25367
D	212	HIS	-	expression tag	UNP O25367
D	213	HIS	-	expression tag	UNP O25367
D	214	HIS	-	expression tag	UNP O25367
D	215	HIS	-	expression tag	UNP O25367

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	2	Total Mg 2 2	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Cl 2	0	0
3	D	1	Total 1	Cl 1	0	0
3	C	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total 30	O 30	0	0
4	B	24	Total 24	O 24	0	0
4	C	35	Total 35	O 35	0	0
4	D	13	Total 13	O 13	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Phosphoserine phosphatase (SerB)




#### • Molecule 1: Phosphoserine phosphatase (SerB)




#### • Molecule 1: Phosphoserine phosphatase (SerB)





● Molecule 1: Phosphoserine phosphatase (SerB)





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.61Å 84.72Å 204.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.07 – 2.40 44.07 – 2.39	Depositor EDS
% Data completeness (in resolution range)	94.0 (44.07-2.40) 93.4 (44.07-2.39)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.60 (at 2.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.229 , 0.271 0.234 , 0.274	Depositor DCC
$R_{free}$ test set	1673 reflections (5.18%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.7	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 39.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 35568 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6500	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/1600	0.71	2/2150 (0.1%)
1	B	0.48	1/1620 (0.1%)	0.64	0/2177
1	C	0.51	0/1631	0.72	1/2190 (0.0%)
1	D	0.49	0/1617	0.68	1/2170 (0.0%)
All	All	0.50	1/6468 (0.0%)	0.69	4/8687 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	136	MSE	CG-SE	-5.69	1.76	1.95

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	205	PRO	N-CA-CB	5.97	110.47	103.30
1	C	91	LYS	N-CA-C	-5.58	95.94	111.00
1	A	91	LYS	N-CA-C	-5.49	96.18	111.00
1	A	205	PRO	N-CA-CB	5.14	109.46	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	0	1624	45	0
1	B	1599	0	1637	49	0
1	C	1611	0	1655	69	0
1	D	1598	0	1630	72	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	30	0	0	0	0
4	B	24	0	0	1	0
4	C	35	0	0	1	0
4	D	13	0	0	1	0
All	All	6500	0	6546	228	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 228 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ASN:ND2	1:A:167:ASN:H	1.66	0.89
1:A:161:VAL:HG13	1:A:171:MSE:HE3	1.53	0.87
1:C:17:GLU:HB3	1:C:20:GLU:HG2	1.59	0.85
1:D:161:VAL:HG21	1:D:171:MSE:HB3	1.58	0.82
1:C:38:LYS:HA	1:C:41:ASN:ND2	1.97	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/217 (92%)	190 (96%)	8 (4%)	1 (0%)	34	48
1	B	204/217 (94%)	195 (96%)	7 (3%)	2 (1%)	19	28
1	C	204/217 (94%)	192 (94%)	10 (5%)	2 (1%)	19	28
1	D	202/217 (93%)	183 (91%)	19 (9%)	0	100	100
All	All	809/868 (93%)	760 (94%)	44 (5%)	5 (1%)	30	43

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	THR
1	B	42	GLY
1	C	124	ASN
1	B	198	PRO
1	C	198	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/181 (97%)	165 (94%)	10 (6%)	25	40
1	B	176/181 (97%)	157 (89%)	19 (11%)	8	11
1	C	177/181 (98%)	160 (90%)	17 (10%)	10	15
1	D	175/181 (97%)	164 (94%)	11 (6%)	22	35
All	All	703/724 (97%)	646 (92%)	57 (8%)	15	22

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	184	LYS
1	C	30	ASP
1	D	110	LEU
1	B	187	LEU
1	B	202	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	2	GLN
1	C	89	ASN
1	D	47	HIS
1	B	196	ASN
1	D	89	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/217 (91%)	-0.12	1 (0%) 91 91	18, 29, 48, 60	0
1	B	200/217 (92%)	0.12	4 (2%) 68 68	22, 40, 58, 64	0
1	C	202/217 (93%)	0.06	2 (0%) 84 83	17, 31, 61, 69	0
1	D	200/217 (92%)	0.33	9 (4%) 37 38	29, 48, 65, 70	0
All	All	800/868 (92%)	0.10	16 (2%) 68 68	17, 37, 61, 70	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	201	ALA	3.5
1	D	64	LEU	3.2
1	D	166	ALA	3.2
1	A	33	LYS	2.9
1	B	42	GLY	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	C	301	1/1	0.89	0.24	6.21	45,45,45,45	0
2	MG	A	300	1/1	0.85	0.14	-0.25	31,31,31,31	0
3	CL	C	302	1/1	0.90	0.12	-0.72	55,55,55,55	0
2	MG	D	300	1/1	0.95	0.13	-0.77	44,44,44,44	0
3	CL	D	302	1/1	0.93	0.08	-1.69	55,55,55,55	0
3	CL	B	302	1/1	0.94	0.08	-1.82	55,55,55,55	0
2	MG	B	300	1/1	0.91	0.09	-1.94	28,28,28,28	0
2	MG	C	300	1/1	0.69	0.10	-2.99	36,36,36,36	0
3	CL	A	302	1/1	0.97	0.06	-3.87	38,38,38,38	0
3	CL	A	303	1/1	0.94	0.15	-	46,46,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.