



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 02:29 AM BST

PDB ID : 2M45
Title : NMR solution structure of the C-terminus of the minichromosome maintenance protein MCM from *Sulfolobus solfataricus*
Authors : Wiedemann, C.; Ohlenschlager, O.; Medagli, B.; Onesti, S.; Gorlach, M.
Deposited on : 2013-01-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	87	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:610-A:618 (9)	0.13	4
2	A:621-A:686 (66)	0.38	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	2, 5, 6, 10, 16, 17, 18, 20
2	1, 4, 7, 8, 13, 15
3	3, 9, 19
Single-model clusters	11; 12; 14

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1384 atoms, of which 714 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Minichromosome maintenance protein MCM.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	87	1384	419	714	112	132	7	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	600	GLY	-	EXPRESSION TAG	UNP Q9UXG1
A	601	SER	-	EXPRESSION TAG	UNP Q9UXG1
A	602	HIS	-	EXPRESSION TAG	UNP Q9UXG1
A	603	MET	-	EXPRESSION TAG	UNP Q9UXG1
A	604	GLY	-	EXPRESSION TAG	UNP Q9UXG1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Minichromosome maintenance protein MCM




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Minichromosome maintenance protein MCM




4.2.2 Score per residue for model 2

- Molecule 1: Minichromosome maintenance protein MCM




4.2.3 Score per residue for model 3

- Molecule 1: Minichromosome maintenance protein MCM




4.2.4 Score per residue for model 4

- Molecule 1: Minichromosome maintenance protein MCM




4.2.5 Score per residue for model 5

- Molecule 1: Minichromosome maintenance protein MCM




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: Minichromosome maintenance protein MCM




4.2.7 Score per residue for model 7

- Molecule 1: Minichromosome maintenance protein MCM




4.2.8 Score per residue for model 8

- Molecule 1: Minichromosome maintenance protein MCM




4.2.9 Score per residue for model 9

- Molecule 1: Minichromosome maintenance protein MCM




4.2.10 Score per residue for model 10

- Molecule 1: Minichromosome maintenance protein MCM




4.2.11 Score per residue for model 11

- Molecule 1: Minichromosome maintenance protein MCM




4.2.12 Score per residue for model 12

- Molecule 1: Minichromosome maintenance protein MCM




4.2.13 Score per residue for model 13

- Molecule 1: Minichromosome maintenance protein MCM




4.2.14 Score per residue for model 14

- Molecule 1: Minichromosome maintenance protein MCM




4.2.15 Score per residue for model 15

- Molecule 1: Minichromosome maintenance protein MCM




4.2.16 Score per residue for model 16

- Molecule 1: Minichromosome maintenance protein MCM




4.2.17 Score per residue for model 17

- Molecule 1: Minichromosome maintenance protein MCM




4.2.18 Score per residue for model 18

- Molecule 1: Minichromosome maintenance protein MCM




4.2.19 Score per residue for model 19

- Molecule 1: Minichromosome maintenance protein MCM



4.2.20 Score per residue for model 20

- Molecule 1: Minichromosome maintenance protein MCM



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	3
OPAL	refinement	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m45_cs.str
Number of chemical shift lists	1
Total number of shifts	1004
Number of shifts mapped to atoms	1004
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.53±0.01	0±0/588 (0.0±0.0%)	0.99±0.05	1±1/781 (0.1±0.1%)
All	All	0.53	0/11760 (0.0%)	0.99	20/15620 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.6±1.1
All	All	0	32

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	623	ARG	NE-CZ-NH2	-8.71	115.94	120.30	15	2
1	A	676	TYR	CB-CG-CD2	-7.64	116.42	121.00	5	2
1	A	623	ARG	NE-CZ-NH1	6.73	123.67	120.30	7	3
1	A	676	TYR	CA-CB-CG	5.87	124.55	113.40	5	1
1	A	670	ARG	NE-CZ-NH2	-5.86	117.37	120.30	20	2
1	A	629	ILE	CB-CG1-CD1	5.60	129.59	113.90	9	1
1	A	666	LEU	CB-CG-CD1	5.44	120.25	111.00	16	2
1	A	670	ARG	CA-CB-CG	5.33	125.13	113.40	16	1
1	A	670	ARG	CB-CG-CD	5.26	125.28	111.60	16	1
1	A	623	ARG	CD-NE-CZ	5.24	130.93	123.60	19	2
1	A	670	ARG	CD-NE-CZ	5.16	130.83	123.60	2	1
1	A	680	PRO	C-N-CA	5.08	134.39	121.70	3	1
1	A	613	THR	CA-CB-CG2	5.06	119.48	112.40	10	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	670	ARG	Sidechain	12
1	A	676	TYR	Sidechain	7
1	A	623	ARG	Sidechain	7
1	A	683	TYR	Sidechain	6

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	586	627	627	1±1
All	All	11720	12540	12540	19

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:626:MET:CE	1:A:665:LEU:HD11	0.60	2.25	14	1
1:A:626:MET:HA	1:A:629:ILE:HG13	0.54	1.78	9	1
1:A:632:ILE:HG21	1:A:651:GLU:OE1	0.53	2.03	14	1
1:A:625:LYS:O	1:A:629:ILE:HD12	0.52	2.05	13	1
1:A:632:ILE:HD13	1:A:651:GLU:OE2	0.49	2.08	14	1
1:A:632:ILE:HD13	1:A:651:GLU:CD	0.48	2.29	14	1
1:A:675:ILE:HG22	1:A:685:LYS:HA	0.47	1.87	19	2
1:A:658:GLU:OE2	1:A:659:LYS:HE2	0.46	2.11	1	1
1:A:636:LEU:CD1	1:A:647:ASP:CG	0.46	2.84	14	1
1:A:651:GLU:O	1:A:655:VAL:HG13	0.45	2.12	11	1
1:A:629:ILE:HD13	1:A:665:LEU:HD22	0.44	1.89	13	1
1:A:660:SER:HA	1:A:663:GLU:HG3	0.43	1.90	17	1
1:A:626:MET:HA	1:A:629:ILE:CG1	0.42	2.44	9	1
1:A:644:LYS:H	1:A:644:LYS:CD	0.42	2.27	11	1
1:A:645:VAL:HG23	1:A:666:LEU:CD1	0.42	2.44	17	1
1:A:658:GLU:OE1	1:A:660:SER:OG	0.42	2.34	2	1
1:A:626:MET:HE3	1:A:665:LEU:HD11	0.41	1.90	14	1
1:A:675:ILE:HG22	1:A:685:LYS:CB	0.41	2.46	2	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	74/87 (85%)	67±2 (91±3%)	6±2 (8±3%)	1±1 (1±1%)	29	74
All	All	1480/1740 (85%)	1349 (91%)	119 (8%)	12 (1%)	29	74

All 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	681	GLU	3
1	A	678	ALA	2
1	A	659	LYS	2
1	A	640	SER	1
1	A	680	PRO	1
1	A	621	SER	1
1	A	622	ALA	1
1	A	617	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/76 (88%)	58±2 (86±3%)	9±2 (14±3%)	8	48
All	All	1340/1520 (88%)	1151 (86%)	189 (14%)	8	48

All 45 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	670	ARG	13
1	A	644	LYS	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	615	MET	11
1	A	610	ASP	9
1	A	613	THR	9
1	A	647	ASP	8
1	A	618	LYS	8
1	A	638	VAL	8
1	A	659	LYS	8
1	A	624	GLU	6
1	A	681	GLU	6
1	A	685	LYS	6
1	A	664	LYS	6
1	A	636	LEU	5
1	A	684	LYS	4
1	A	633	ILE	4
1	A	671	LYS	4
1	A	677	GLU	4
1	A	676	TYR	4
1	A	623	ARG	4
1	A	686	VAL	4
1	A	666	LEU	4
1	A	630	ILE	3
1	A	654	GLN	3
1	A	658	GLU	3
1	A	650	LYS	3
1	A	661	ASN	3
1	A	660	SER	3
1	A	634	ASP	3
1	A	651	GLU	2
1	A	682	CYS	2
1	A	653	GLN	2
1	A	679	LYS	2
1	A	646	LYS	2
1	A	626	MET	1
1	A	631	GLU	1
1	A	665	LEU	1
1	A	669	MET	1
1	A	627	MET	1
1	A	612	ASP	1
1	A	625	LYS	1
1	A	683	TYR	1
1	A	655	VAL	1
1	A	668	ASP	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	628	LYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: 2m45_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1004
Number of shifts mapped to atoms	1004
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	87	-0.40 \pm 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	81	0.07 \pm 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	83	-0.29 \pm 0.14	None needed (< 0.5 ppm)
^{15}N	82	0.22 \pm 0.43	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 831 atoms were assigned a chemical shift out of a possible 936. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	371/373 (99%)	149/149 (100%)	148/150 (99%)	74/74 (100%)
Sidechain	444/547 (81%)	259/320 (81%)	185/206 (90%)	0/21 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	16/16 (100%)	8/8 (100%)	8/8 (100%)	0/0 (—%)
Overall	831/936 (89%)	416/477 (87%)	341/364 (94%)	74/95 (78%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 946 atoms were assigned a chemical shift out of a possible 1069. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	421/431 (98%)	169/172 (98%)	170/174 (98%)	82/85 (96%)
Sidechain	505/615 (82%)	297/362 (82%)	208/230 (90%)	0/23 (0%)
Aromatic	20/23 (87%)	10/12 (83%)	10/10 (100%)	0/1 (0%)
Overall	946/1069 (88%)	476/546 (87%)	388/414 (94%)	82/109 (75%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

