



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 02:27 AM BST

PDB ID : 2M47  
Title : Solution NMR structure of the Polyketide\_cyc-like protein Cgl2372 from Corynebacterium glutamicum, Northeast Structural Genomics Consortium Target CgR160  
Authors : Yang, Y.; Ramelot, T.A.; Lee, D.; Ciccocanti, C.; Sapin, A.; Janjua, H.; Nair, R.; Rost, B.; Acton, T.B.; Xiao, R.; Everett, J.K.; Montelione, G.T.; Kennedy, M.A.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2013-01-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 85%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	163	 78% 21%

## 2 Ensemble composition and analysis i

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:5-A:25, A:40-A:56, A:66-A:156 (129)	0.78	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 6 single-model clusters were found.

Cluster number	Models
1	7, 14, 18
2	2, 4, 6
3	1, 12
4	8, 9
5	3, 11
6	16, 17
Single-model clusters	5; 10; 13; 15; 19; 20

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2599 atoms, of which 1272 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Uncharacterized protein Cgl2373.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	163	2599	842	1272	228	252	5	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	156	LEU	-	EXPRESSION TAG	UNP Q8NN40
A	157	GLU	-	EXPRESSION TAG	UNP Q8NN40
A	158	HIS	-	EXPRESSION TAG	UNP Q8NN40
A	159	HIS	-	EXPRESSION TAG	UNP Q8NN40
A	160	HIS	-	EXPRESSION TAG	UNP Q8NN40
A	161	HIS	-	EXPRESSION TAG	UNP Q8NN40
A	162	HIS	-	EXPRESSION TAG	UNP Q8NN40
A	163	HIS	-	EXPRESSION TAG	UNP Q8NN40

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Uncharacterized protein Cgl2373




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.2 Score per residue for model 2

- Molecule 1: Uncharacterized protein Cgl2373



H161  
H162  
H163

### 4.2.3 Score per residue for model 3

- Molecule 1: Uncharacterized protein Cgl2373

Chain A:  75% 21%

M1 P2 K3 S4 D26 I27 T28 R29 T30 G31 E32 H33 S34 P35 V36 C37 E38 K39 M42 D43 E44 P48 R57 N58 V59 T60 P61 E62 R63 T64 M65 R78 E106 E157 H158 H159 H160 H161 H162 H163

### 4.2.4 Score per residue for model 4

- Molecule 1: Uncharacterized protein Cgl2373

Chain A:  74% 6% 21%

M1 P2 K3 S4 I11 L23 D26 I27 T28 R29 T30 G31 E32 H33 S34 P35 V36 C37 E38 K39 C40 R57 N58 V59 T60 P61 E62 R63 T64 M65 E70 D86 N95 E106 K121 L145 E157 H158 H159 H160 H161 H162 H163

### 4.2.5 Score per residue for model 5

- Molecule 1: Uncharacterized protein Cgl2373

Chain A:  69% 10% 21%

M1 P2 K3 S4 L5 E8 V20 D26 I27 T28 R29 T30 G31 E32 H33 S34 P35 V36 C37 E38 K39 C40 D43 F54 R57 N58 V59 T60 P61 E62 R63 T64 M65 R68 R78 T85 D86 T102 E106 F118 S126 I130 Q149 E153

E157  
H158  
H159  
H160  
H161  
H162  
H163

### 4.2.6 Score per residue for model 6

- Molecule 1: Uncharacterized protein Cgl2373

Chain A:  75% 21%

M1 P2 K3 S4 L5 D26 I27 T28 R29 T30 G31 E32 H33 S34 P35 V36 C37 E38 K39 R57 N58 V59 T60 P61 E62 R63 T64 M65 S83 D86 G87 E106 F112 K113 E157 H158 H159 H160 H161 H162 H163


### 4.2.7 Score per residue for model 7

- Molecule 1: Uncharacterized protein Cgl2373




### 4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: Uncharacterized protein Cgl2373




### 4.2.9 Score per residue for model 9

- Molecule 1: Uncharacterized protein Cgl2373




### 4.2.10 Score per residue for model 10

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.11 Score per residue for model 11

- Molecule 1: Uncharacterized protein Cgl2373




#### 4.2.12 Score per residue for model 12

- Molecule 1: Uncharacterized protein Cgl2373




#### 4.2.13 Score per residue for model 13

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.14 Score per residue for model 14

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.15 Score per residue for model 15

- Molecule 1: Uncharacterized protein Cgl2373






#### 4.2.16 Score per residue for model 16

- Molecule 1: Uncharacterized protein Cgl2373




#### 4.2.17 Score per residue for model 17

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.18 Score per residue for model 18

- Molecule 1: Uncharacterized protein Cgl2373




#### 4.2.19 Score per residue for model 19

- Molecule 1: Uncharacterized protein Cgl2373



#### 4.2.20 Score per residue for model 20

- Molecule 1: Uncharacterized protein Cgl2373



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CNS	structure solution	
CNS	geometry optimization	
CYANA	refinement	3.0
CYANA	geometry optimization	3.0
CYANA	structure solution	3.0
AutoStructure	refinement	2.1
TALOS+	geometry optimization	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m47_cs.str
Number of chemical shift lists	1
Total number of shifts	1842
Number of shifts mapped to atoms	1842
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	85%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1034	998	996	2±2
All	All	20680	19960	19920	48

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:ILE:HB	1:A:97:PRO:HB3	0.54	1.80	15	1
1:A:11:ILE:HG22	1:A:145:LEU:HD12	0.53	1.79	20	4
1:A:42:TRP:HB2	1:A:45:ASP:HB2	0.52	1.82	19	1
1:A:106:GLU:HB3	1:A:145:LEU:HD21	0.50	1.82	13	2
1:A:67:THR:HG22	1:A:85:THR:HG22	0.50	1.81	20	1
1:A:112:PRO:HA	1:A:115:GLN:HB3	0.49	1.84	7	2
1:A:92:ILE:HB	1:A:107:SER:HB2	0.48	1.85	16	2
1:A:95:MET:N	1:A:95:MET:SD	0.48	2.87	20	2
1:A:139:THR:O	1:A:143:GLU:HG2	0.47	2.09	9	1
1:A:43:ASP:HB2	1:A:54:PHE:HB3	0.46	1.87	5	2
1:A:5:LEU:HD21	1:A:130:ILE:HB	0.46	1.88	5	1
1:A:68:ARG:HH21	1:A:86:ASP:HB2	0.46	1.69	16	1
1:A:54:PHE:HB2	1:A:69:SER:HB2	0.46	1.87	2	1
1:A:24:VAL:HG12	1:A:148:ILE:HD13	0.45	1.87	1	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:THR:HG22	1:A:109:GLU:HA	0.45	1.89	1	1
1:A:148:ILE:O	1:A:152:LEU:HG	0.44	2.12	10	1
1:A:83:SER:HB2	1:A:87:GLY:HA2	0.44	1.89	6	1
1:A:42:TRP:HB3	1:A:44:GLU:HG2	0.44	1.90	3	1
1:A:104:LEU:HD12	1:A:145:LEU:HD11	0.44	1.90	20	1
1:A:71:VAL:HA	1:A:82:TRP:HA	0.43	1.88	9	1
1:A:118:PHE:HB3	1:A:126:SER:HB3	0.43	1.91	5	1
1:A:137:ALA:O	1:A:141:ILE:HG12	0.43	2.13	10	1
1:A:106:GLU:HB3	1:A:141:ILE:HG23	0.42	1.92	17	1
1:A:53:HIS:HB3	1:A:70:GLU:HG3	0.42	1.91	1	1
1:A:19:GLN:HB3	1:A:152:LEU:HD22	0.41	1.92	17	1
1:A:94:SER:HB3	1:A:105:THR:HB	0.41	1.92	13	1
1:A:95:MET:SD	1:A:95:MET:N	0.41	2.93	16	1
1:A:128:GLU:O	1:A:132:LYS:HE2	0.41	2.15	7	1
1:A:40:CYS:SG	1:A:56:GLY:HA2	0.41	2.55	19	1
1:A:20:VAL:HG21	1:A:102:THR:HB	0.41	1.92	5	3
1:A:122:PHE:HB3	1:A:125:LYS:HD3	0.41	1.91	13	1
1:A:71:VAL:HG11	1:A:74:ALA:HB2	0.41	1.93	19	1
1:A:84:VAL:HB	1:A:89:VAL:HB	0.41	1.93	20	1
1:A:90:LYS:HB2	1:A:109:GLU:HB2	0.41	1.93	7	1
1:A:92:ILE:HB	1:A:107:SER:HB3	0.41	1.93	9	1
1:A:14:ALA:O	1:A:156:LEU:HD11	0.40	2.17	10	1
1:A:149:GLN:O	1:A:153:GLU:HG2	0.40	2.17	5	1
1:A:94:SER:HB2	1:A:105:THR:HB	0.40	1.93	17	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/163 (79%)	116±2 (90±2%)	12±2 (9±2%)	2±1 (1±1%)	21	68
All	All	2580/3260 (79%)	2316 (90%)	232 (9%)	32 (1%)	21	68

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	86	ASP	10
1	A	76	PRO	4
1	A	78	ARG	3
1	A	112	PRO	3
1	A	48	PRO	2
1	A	85	THR	2
1	A	5	LEU	2
1	A	113	LYS	2
1	A	71	VAL	1
1	A	47	GLY	1
1	A	87	GLY	1
1	A	49	VAL	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	112/145 (77%)	108±2 (97±2%)	4±2 (3±2%)	49 89
All	All	2240/2900 (77%)	2165 (97%)	75 (3%)	49 89

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	106	GLU	12
1	A	115	GLN	4
1	A	78	ARG	4
1	A	66	GLU	4
1	A	8	GLU	4
1	A	95	MET	3
1	A	150	ARG	3
1	A	54	PHE	3
1	A	121	LYS	3
1	A	42	TRP	3
1	A	23	LEU	2
1	A	109	GLU	2
1	A	100	GLU	2
1	A	82	TRP	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	68	ARG	2
1	A	129	GLU	2
1	A	116	ARG	2
1	A	5	LEU	1
1	A	19	GLN	1
1	A	70	GLU	1
1	A	130	ILE	1
1	A	75	GLU	1
1	A	111	THR	1
1	A	43	ASP	1
1	A	53	HIS	1
1	A	18	ASN	1
1	A	120	ASP	1
1	A	77	ASN	1
1	A	93	TYR	1
1	A	133	ARG	1
1	A	88	ASN	1
1	A	85	THR	1
1	A	145	LEU	1
1	A	90	LYS	1
1	A	44	GLU	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 85% for the well-defined parts and 80% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2m47\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1842
Number of shifts mapped to atoms	1842
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	14

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	156	$-0.37 \pm 0.18$	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	146	$-0.10 \pm 0.14$	None needed (< 0.5 ppm)
$^{13}\text{C}'$	142	$0.10 \pm 0.14$	None needed (< 0.5 ppm)
$^{15}\text{N}$	142	$-0.02 \pm 0.39$	None needed (< 0.5 ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 85%, i.e. 1347 atoms were assigned a chemical shift out of a possible 1588. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	613/633 (97%)	249/252 (99%)	245/258 (95%)	119/123 (97%)
Sidechain	655/800 (82%)	402/465 (86%)	248/305 (81%)	5/30 (17%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	79/155 (51%)	43/81 (53%)	29/65 (45%)	7/9 (78%)
Overall	1347/1588 (85%)	694/798 (87%)	522/628 (83%)	131/162 (81%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 80%, i.e. 1638 atoms were assigned a chemical shift out of a possible 2048. 0 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	743/797 (93%)	303/317 (96%)	298/326 (91%)	142/154 (92%)
Sidechain	806/1024 (79%)	498/600 (83%)	303/382 (79%)	5/42 (12%)
Aromatic	89/227 (39%)	48/117 (41%)	32/87 (37%)	9/23 (39%)
Overall	1638/2048 (80%)	849/1034 (82%)	633/795 (80%)	156/219 (71%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	69	SER	HB3	1.21	5.25 – 2.45	-9.4
1	A	58	ASN	HB3	0.40	4.41 – 1.11	-7.2
1	A	55	THR	HB	1.89	5.82 – 2.52	-6.9
1	A	27	ILE	HG13	-1.38	3.26 – -0.84	-6.3
1	A	58	ASN	HB2	0.90	4.36 – 1.26	-6.2
1	A	25	SER	HB3	2.22	5.25 – 2.45	-5.8
1	A	137	ALA	HB2	-0.05	2.61 – 0.11	-5.6
1	A	137	ALA	HB1	-0.05	2.61 – 0.11	-5.6
1	A	137	ALA	HB3	-0.05	2.61 – 0.11	-5.6
1	A	134	ARG	HG3	3.16	3.00 – 0.10	5.6
1	A	76	PRO	HA	2.62	6.05 – 2.75	-5.4
1	A	27	ILE	HD12	-0.83	2.13 – -0.77	-5.2
1	A	27	ILE	HD13	-0.83	2.13 – -0.77	-5.2
1	A	27	ILE	HD11	-0.83	2.13 – -0.77	-5.2

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble

composition.

Random coil index (RCI) for chain A:

