



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:49 PM GMT

PDB ID : 1M6Z
Title : Crystal structure of reduced recombinant cytochrome c4 from *Pseudomonas stutzeri*
Authors : Noergaard, A.; Harris, P.; Larsen, S.; Christensen, H.E.M.
Deposited on : 2002-07-18
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2199 (1.40-1.32)
Clashscore	102246	2337 (1.40-1.32)
Ramachandran outliers	100387	2280 (1.40-1.32)
Sidechain outliers	100360	2279 (1.40-1.32)
RSRZ outliers	91569	2199 (1.40-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>2%</div> <div>89%</div> <div>11%</div> <div>.</div> </div>
1	B	190	<div> <div>5%</div> <div>87%</div> <div>11%</div> <div>..</div> </div>
1	C	190	<div> <div>3%</div> <div>93%</div> <div>6%</div> <div>.</div> </div>
1	D	190	<div> <div>9%</div> <div>91%</div> <div>9%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TRS	D	201	-	-	X	X
4	GOL	C	201	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	9	0
			1414	873	254	276	11			
1	B	190	Total	C	N	O	S	0	2	0
			1384	853	248	274	9			
1	C	190	Total	C	N	O	S	0	8	0
			1402	864	251	278	9			
1	D	190	Total	C	N	O	S	0	5	0
			1391	858	248	276	9			

- Molecule 2 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	C	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	C	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	D	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	D	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O		
			8	4	1	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	188	Total	O	0	0
			188	188		
5	B	135	Total	O	0	0
			135	135		
5	C	201	Total	O	0	0
			201	201		
5	D	169	Total	O	0	0
			169	169		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome c4





• Molecule 1: Cytochrome c4



• Molecule 1: Cytochrome c4



• Molecule 1: Cytochrome c4



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.11Å 57.56Å 83.75Å 90.00° 104.32° 90.00°	Depositor
Resolution (Å)	30.00 – 1.35 28.78 – 1.35	Depositor EDS
% Data completeness (in resolution range)	92.6 (30.00-1.35) 92.6 (28.78-1.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.38 (at 1.35Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.140 , 0.191 0.145 , 0.158	Depositor DCC
R_{free} test set	7157 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	11.2	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 141227 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6648	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 84.77 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5182e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TRS, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/1477	1.20	11/1986 (0.6%)
1	B	0.48	0/1417	1.24	13/1906 (0.7%)
1	C	0.54	0/1461	1.18	13/1966 (0.7%)
1	D	0.54	0/1437	1.14	9/1934 (0.5%)
All	All	0.53	0/5792	1.19	46/7792 (0.6%)

There are no bond length outliers.

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	38	ARG	NE-CZ-NH2	21.52	131.06	120.30
1	C	168	ARG	NE-CZ-NH1	10.72	125.66	120.30
1	D	94	TYR	CB-CG-CD2	10.69	127.41	121.00
1	D	108	ARG	NE-CZ-NH2	-9.68	115.46	120.30
1	A	38[A]	ARG	CD-NE-CZ	9.28	136.59	123.60
1	A	38[B]	ARG	CD-NE-CZ	9.28	136.59	123.60
1	C	168	ARG	NE-CZ-NH2	-9.19	115.71	120.30
1	D	38	ARG	CD-NE-CZ	8.88	136.04	123.60
1	A	131	ASP	CB-CG-OD2	-8.62	110.54	118.30
1	D	94	TYR	CB-CG-CD1	-8.50	115.90	121.00
1	C	38[A]	ARG	CD-NE-CZ	8.49	135.48	123.60
1	C	38[B]	ARG	CD-NE-CZ	8.49	135.48	123.60
1	B	38	ARG	NE-CZ-NH1	-8.48	116.06	120.30
1	D	131	ASP	CB-CG-OD2	-7.98	111.11	118.30
1	C	154	ARG	NE-CZ-NH2	-7.50	116.55	120.30
1	C	38[A]	ARG	NE-CZ-NH2	-7.17	116.71	120.30
1	C	38[B]	ARG	NE-CZ-NH2	-7.17	116.71	120.30
1	C	163	ASP	CB-CG-OD1	7.14	124.72	118.30
1	C	38[A]	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	C	38[B]	ARG	NE-CZ-NH1	7.10	123.85	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	61	ARG	NE-CZ-NH2	-7.00	116.80	120.30
1	B	38	ARG	CD-NE-CZ	6.63	132.89	123.60
1	B	108	ARG	CA-CB-CG	6.62	127.97	113.40
1	A	23[A]	ASN	O-C-N	-6.48	112.33	122.70
1	A	23[B]	ASN	O-C-N	-6.48	112.33	122.70
1	B	77	ASP	CB-CG-OD2	-6.33	112.60	118.30
1	B	126	ASN	CB-CG-ND2	6.28	131.76	116.70
1	A	38[A]	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	A	38[B]	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	D	108	ARG	NE-CZ-NH1	6.12	123.36	120.30
1	B	158	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	D	38	ARG	NE-CZ-NH2	-6.08	117.26	120.30
1	B	155	GLU	OE1-CD-OE2	-6.03	116.07	123.30
1	B	155	GLU	CG-CD-OE1	6.03	130.35	118.30
1	D	96	ASP	CB-CG-OD2	6.03	123.72	118.30
1	B	38	ARG	NH1-CZ-NH2	-5.93	112.88	119.40
1	A	38[A]	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	A	38[B]	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	77	ASP	CB-CG-OD1	5.83	123.55	118.30
1	B	108	ARG	CD-NE-CZ	5.57	131.39	123.60
1	B	96	ASP	CB-CG-OD1	5.49	123.24	118.30
1	D	102	GLN	CG-CD-OE1	5.42	132.44	121.60
1	A	94	TYR	CB-CG-CD2	-5.14	117.92	121.00
1	C	23[A]	ASN	C-N-CA	5.09	134.42	121.70
1	C	23[B]	ASN	C-N-CA	5.09	134.42	121.70
1	C	145	TYR	CB-CG-CD2	5.07	124.05	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1414	0	1383	13	0
1	B	1384	0	1352	13	0
1	C	1402	0	1367	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1391	0	1361	14	0
2	A	86	0	60	6	0
2	B	86	0	60	3	0
2	C	86	0	60	0	0
2	D	86	0	60	1	0
3	D	8	0	11	7	0
4	A	6	0	7	0	0
4	C	6	0	7	2	0
5	A	188	0	0	3	0
5	B	135	0	0	4	0
5	C	201	0	0	5	0
5	D	169	0	0	5	0
All	All	6648	0	5728	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:76:GLN:HB3	4:C:201:GOL:H32	1.58	0.83
1:D:137[A]:LYS:HG2	3:D:201:TRS:H22	1.62	0.79
1:D:137[B]:LYS:HZ2	3:D:201:TRS:H12	1.50	0.76
1:D:139:GLY:O	3:D:201:TRS:H21	1.88	0.74
1:A:69[A]:MET:HE1	2:A:199:HEC:HMC2	1.70	0.74
1:D:137[B]:LYS:HG2	3:D:201:TRS:H22	1.69	0.73
1:C:173:LYS:HE3	5:C:267:HOH:O	1.91	0.70
1:A:62:LYS:HE3	5:A:351:HOH:O	1.94	0.67
1:B:24:SER:HB3	5:B:216:HOH:O	1.95	0.65
1:D:92[A]:VAL:HG12	5:D:365:HOH:O	1.96	0.65
1:D:173:LYS:HE3	5:D:281:HOH:O	1.97	0.63
1:A:102:GLN:HG2	5:A:380:HOH:O	2.00	0.61
1:B:114:GLN:O	1:B:173:LYS:HE2	2.00	0.61
1:D:112:LEU:HG	5:D:254:HOH:O	2.01	0.61
1:D:137[B]:LYS:NZ	3:D:201:TRS:H12	2.16	0.60
1:A:114:GLN:O	1:A:173:LYS:HE2	2.03	0.59
1:B:165:MET:HG2	1:B:168:ARG:CZ	2.34	0.57
1:A:135:PHE:CD1	2:A:200:HEC:HMD3	2.42	0.54
1:C:20[B]:VAL:HG12	5:C:366:HOH:O	2.09	0.52
1:B:135:PHE:CD1	2:B:200:HEC:HMD3	2.45	0.52
1:B:126:ASN:ND2	1:B:128:VAL:HG13	2.25	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:HIS:HA	5:B:299:HOH:O	2.09	0.51
1:D:137[A]:LYS:HG2	3:D:201:TRS:C2	2.37	0.50
2:A:200:HEC:HBC3	2:A:200:HEC:HMC1	1.94	0.50
1:B:25:PRO:HD2	5:B:238:HOH:O	2.12	0.49
1:D:62:LYS:HE3	5:D:305:HOH:O	2.12	0.48
1:A:190:HIS:HE1	5:A:294:HOH:O	1.95	0.48
2:B:200:HEC:HMC1	2:B:200:HEC:HBC3	1.97	0.47
1:A:5:GLU:CD	1:B:101:LYS:HD3	2.35	0.47
1:C:38[A]:ARG:NE	5:C:256:HOH:O	2.49	0.46
1:A:69[B]:MET:HE2	2:A:199:HEC:HMC2	1.97	0.46
1:A:126:ASN:OD1	1:A:128[B]:VAL:HG22	2.17	0.45
1:C:45:GLN:NE2	5:C:296:HOH:O	2.49	0.45
1:C:76:GLN:CB	4:C:201:GOL:H32	2.38	0.45
1:D:137[B]:LYS:HG2	3:D:201:TRS:C2	2.41	0.45
1:A:69[B]:MET:CE	2:A:199:HEC:HMC2	2.46	0.44
1:D:128[A]:VAL:HG12	5:D:363:HOH:O	2.16	0.44
1:C:57:GLU:O	1:C:57:GLU:HG2	2.16	0.44
1:C:137[A]:LYS:HD3	5:C:402:HOH:O	2.18	0.43
1:A:29:PHE:CD1	2:A:199:HEC:HMD3	2.54	0.43
1:A:48:LYS:NZ	1:A:73:LEU:O	2.49	0.43
1:B:18:HIS:C	1:B:24:SER:HB2	2.40	0.42
1:B:165:MET:HG2	1:B:168:ARG:NH2	2.35	0.42
1:C:47:ILE:HA	1:C:61:ARG:O	2.20	0.42
1:D:126:ASN:OD1	1:D:128[A]:VAL:HG22	2.19	0.41
1:B:29:PHE:CD1	2:B:199:HEC:HMD3	2.54	0.41
1:D:135:PHE:CD1	2:D:200:HEC:HMD3	2.56	0.41
1:B:113:ASP:OD1	1:B:113:ASP:N	2.54	0.40
1:A:113:ASP:OD1	1:A:113:ASP:N	2.51	0.40
1:B:100:ALA:HA	5:B:225:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/190 (104%)	192 (98%)	5 (2%)	0	100	100
1	B	190/190 (100%)	187 (98%)	3 (2%)	0	100	100
1	C	196/190 (103%)	191 (97%)	5 (3%)	0	100	100
1	D	193/190 (102%)	190 (98%)	3 (2%)	0	100	100
All	All	776/760 (102%)	760 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/138 (106%)	143 (97%)	4 (3%)	52	15
1	B	140/138 (101%)	134 (96%)	6 (4%)	35	5
1	C	146/138 (106%)	145 (99%)	1 (1%)	88	68
1	D	143/138 (104%)	138 (96%)	5 (4%)	43	9
All	All	576/552 (104%)	560 (97%)	16 (3%)	55	14

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	VAL
1	A	102	GLN
1	A	108[A]	ARG
1	A	108[B]	ARG
1	B	92	VAL
1	B	101	LYS
1	B	102	GLN
1	B	114	GLN
1	B	126	ASN
1	B	130	ASN
1	C	57	GLU
1	D	45	GLN
1	D	57	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	89	LYS
1	D	92[A]	VAL
1	D	92[B]	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	HIS
1	B	126	ASN
1	B	130	ASN
1	B	157	ASN
1	C	45	GLN
1	D	45	GLN
1	D	102	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	199	1	24,50,50	2.26	6 (25%)	19,82,82	3.10	8 (42%)
2	HEC	A	200	1	24,50,50	1.87	3 (12%)	19,82,82	2.77	5 (26%)
4	GOL	A	201	-	5,5,5	0.80	0	5,5,5	1.91	2 (40%)
2	HEC	B	199	1	24,50,50	2.13	6 (25%)	19,82,82	3.27	11 (57%)
2	HEC	B	200	1	24,50,50	1.95	5 (20%)	19,82,82	2.64	5 (26%)
2	HEC	C	199	1	24,50,50	2.05	6 (25%)	19,82,82	3.09	10 (52%)
2	HEC	C	200	1	24,50,50	1.90	5 (20%)	19,82,82	2.62	4 (21%)
4	GOL	C	201	-	5,5,5	1.11	0	5,5,5	3.48	3 (60%)
2	HEC	D	199	1	24,50,50	2.02	7 (29%)	19,82,82	2.92	9 (47%)
2	HEC	D	200	1	24,50,50	1.83	5 (20%)	19,82,82	2.64	5 (26%)
3	TRS	D	201	-	7,7,7	2.73	3 (42%)	9,9,9	2.83	6 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	199	1	-	0/6/54/54	0/0/8/8
2	HEC	A	200	1	-	0/6/54/54	0/0/8/8
4	GOL	A	201	-	-	0/4/4/4	0/0/0/0
2	HEC	B	199	1	-	0/6/54/54	0/0/8/8
2	HEC	B	200	1	-	0/6/54/54	0/0/8/8
2	HEC	C	199	1	-	0/6/54/54	0/0/8/8
2	HEC	C	200	1	-	0/6/54/54	0/0/8/8
4	GOL	C	201	-	-	0/4/4/4	0/0/0/0
2	HEC	D	199	1	-	0/6/54/54	0/0/8/8
2	HEC	D	200	1	-	0/6/54/54	0/0/8/8
3	TRS	D	201	-	-	0/9/9/9	0/0/0/0

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	199	HEC	C3C-C2C	-6.92	1.33	1.40
2	B	200	HEC	C3B-C2B	-6.87	1.33	1.40
2	C	200	HEC	C3B-C2B	-6.74	1.33	1.40
2	A	200	HEC	C3B-C2B	-6.49	1.34	1.40
3	D	201	TRS	C2-C	-6.24	1.41	1.53
2	C	199	HEC	C3C-C2C	-6.06	1.34	1.40
2	D	200	HEC	C3B-C2B	-6.04	1.34	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	199	HEC	C3C-C2C	-5.90	1.34	1.40
2	D	199	HEC	C3C-C2C	-5.64	1.34	1.40
2	A	199	HEC	C3B-C2B	-4.67	1.35	1.40
2	C	199	HEC	C3B-C2B	-4.26	1.36	1.40
2	D	199	HEC	C3B-C2B	-4.13	1.36	1.40
2	B	199	HEC	C3B-C2B	-3.79	1.36	1.40
2	D	200	HEC	C3C-C2C	-3.73	1.36	1.40
2	A	200	HEC	C3C-C2C	-3.63	1.37	1.40
2	B	200	HEC	C3C-C2C	-3.60	1.37	1.40
2	C	200	HEC	C3C-C2C	-3.35	1.37	1.40
3	D	201	TRS	O1-C1	-2.39	1.34	1.42
3	D	201	TRS	C-N	-2.09	1.47	1.50
2	C	200	HEC	C1C-CHC	-2.03	1.34	1.39
2	B	200	HEC	C1C-CHC	-2.02	1.34	1.39
2	C	200	HEC	C3C-C4C	2.05	1.47	1.42
2	B	200	HEC	C3C-C4C	2.12	1.47	1.42
2	D	200	HEC	C3C-C4C	2.23	1.47	1.42
2	D	199	HEC	C4D-CHA	2.32	1.46	1.39
2	D	200	HEC	C4C-NC	2.33	1.39	1.36
2	D	200	HEC	C3B-C4B	2.34	1.48	1.42
2	A	199	HEC	C3C-C4C	2.41	1.48	1.42
2	C	199	HEC	C3C-C4C	2.43	1.48	1.42
2	D	199	HEC	C1A-NA	2.45	1.40	1.36
2	C	200	HEC	C3B-C4B	2.55	1.48	1.42
2	D	199	HEC	CBB-CAB	2.56	1.60	1.49
2	D	199	HEC	C3C-C4C	2.58	1.48	1.42
2	B	199	HEC	C3C-C4C	2.60	1.48	1.42
2	C	199	HEC	CBB-CAB	2.65	1.60	1.49
2	A	199	HEC	CBB-CAB	2.71	1.60	1.49
2	B	200	HEC	C3B-C4B	2.79	1.49	1.42
2	A	200	HEC	C3B-C4B	2.81	1.49	1.42
2	C	199	HEC	C1A-NA	2.92	1.40	1.36
2	B	199	HEC	CBB-CAB	3.26	1.62	1.49
2	C	199	HEC	CMB-C2B	3.29	1.58	1.51
2	B	199	HEC	C1A-NA	3.36	1.41	1.36
2	A	199	HEC	C1A-NA	3.41	1.41	1.36
2	B	199	HEC	CMB-C2B	3.52	1.59	1.51
2	D	199	HEC	CMB-C2B	3.58	1.59	1.51
2	A	199	HEC	CMB-C2B	3.61	1.59	1.51

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	199	HEC	CBB-CAB-C3B	-8.39	108.70	127.35
2	A	199	HEC	CBB-CAB-C3B	-8.12	109.31	127.35
2	C	199	HEC	CBB-CAB-C3B	-7.54	110.60	127.35
2	C	199	HEC	CBC-CAC-C3C	-7.49	110.70	127.35
2	C	200	HEC	CBB-CAB-C3B	-7.48	110.73	127.35
2	D	199	HEC	CBB-CAB-C3B	-7.42	110.86	127.35
2	B	199	HEC	CBC-CAC-C3C	-7.35	111.02	127.35
2	A	199	HEC	CBC-CAC-C3C	-7.22	111.31	127.35
2	A	200	HEC	CBB-CAB-C3B	-7.01	111.78	127.35
2	D	200	HEC	CBB-CAB-C3B	-6.98	111.83	127.35
2	B	200	HEC	CBB-CAB-C3B	-6.96	111.88	127.35
2	A	200	HEC	CBC-CAC-C3C	-6.70	112.47	127.35
4	C	201	GOL	O2-C2-C1	-6.43	79.15	108.65
2	D	199	HEC	CBC-CAC-C3C	-6.37	113.20	127.35
2	D	200	HEC	CBC-CAC-C3C	-5.96	114.11	127.35
2	B	200	HEC	CBC-CAC-C3C	-5.85	114.36	127.35
2	C	200	HEC	CBC-CAC-C3C	-5.76	114.56	127.35
2	B	199	HEC	CMC-C2C-C1C	-3.76	122.14	128.36
3	D	201	TRS	C3-C-C1	-3.72	102.72	110.78
2	C	199	HEC	CMC-C2C-C1C	-3.69	122.25	128.36
2	A	199	HEC	CMC-C2C-C1C	-3.67	122.29	128.36
2	C	200	HEC	CMB-C2B-C1B	-3.65	122.32	128.36
2	B	200	HEC	CMB-C2B-C1B	-3.61	122.39	128.36
2	A	200	HEC	CMB-C2B-C1B	-3.51	122.56	128.36
2	D	200	HEC	CMB-C2B-C1B	-3.49	122.58	128.36
4	A	201	GOL	O2-C2-C1	-3.44	92.90	108.65
2	D	199	HEC	CAD-CBD-CGD	-3.40	106.52	112.75
2	B	199	HEC	CMB-C2B-C1B	-3.40	122.75	128.36
2	B	199	HEC	CAA-C2A-C1A	-3.34	123.38	127.01
2	D	199	HEC	CMC-C2C-C1C	-3.14	123.17	128.36
2	D	200	HEC	CMC-C2C-C1C	-3.14	123.17	128.36
2	C	199	HEC	CAA-C2A-C1A	-3.05	123.69	127.01
2	A	200	HEC	CMC-C2C-C1C	-3.04	123.33	128.36
2	C	199	HEC	CAD-CBD-CGD	-3.04	107.17	112.75
2	D	199	HEC	CMB-C2B-C1B	-2.93	123.51	128.36
2	C	200	HEC	CMC-C2C-C1C	-2.93	123.51	128.36
2	A	199	HEC	CAA-C2A-C1A	-2.89	123.87	127.01
2	B	200	HEC	CMC-C2C-C1C	-2.81	123.71	128.36
2	D	199	HEC	CAA-C2A-C1A	-2.80	123.97	127.01
2	A	199	HEC	CMB-C2B-C1B	-2.78	123.76	128.36
2	A	199	HEC	CAD-CBD-CGD	-2.62	107.94	112.75
2	B	199	HEC	CAD-CBD-CGD	-2.50	108.16	112.75
3	D	201	TRS	C2-C-C1	-2.48	105.42	110.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	201	GOL	O2-C2-C3	-2.43	97.52	108.65
2	C	199	HEC	CBA-CAA-C2A	-2.42	108.19	112.53
2	D	199	HEC	CBA-CAA-C2A	-2.26	108.48	112.53
2	B	199	HEC	CBA-CAA-C2A	-2.24	108.52	112.53
2	C	199	HEC	CMB-C2B-C1B	-2.22	124.70	128.36
2	C	199	HEC	C3C-C4C-NC	-2.04	107.10	110.94
2	B	199	HEC	C3C-C4C-NC	-2.02	107.12	110.94
2	B	199	HEC	CMD-C2D-C3D	2.02	129.47	125.24
2	A	199	HEC	CAD-C3D-C4D	2.04	129.22	127.01
4	A	201	GOL	C3-C2-C1	2.06	119.20	111.12
3	D	201	TRS	C3-C-C2	2.12	115.37	110.78
2	B	199	HEC	CMA-C3A-C2A	2.12	129.67	125.24
2	C	199	HEC	C3B-C4B-NB	2.16	115.03	110.94
2	D	199	HEC	C3B-C4B-NB	2.18	115.06	110.94
3	D	201	TRS	O2-C2-C	2.18	115.60	111.18
2	B	199	HEC	C3B-C4B-NB	2.29	115.27	110.94
2	D	200	HEC	CBD-CAD-C3D	2.31	116.67	112.53
2	C	199	HEC	CMA-C3A-C2A	2.44	130.35	125.24
2	B	200	HEC	CBD-CAD-C3D	2.49	116.98	112.53
2	D	199	HEC	CMA-C3A-C2A	2.54	130.56	125.24
2	A	199	HEC	CMA-C3A-C2A	2.58	130.64	125.24
4	C	201	GOL	C3-C2-C1	3.15	123.47	111.12
2	A	200	HEC	CBD-CAD-C3D	3.62	119.01	112.53
3	D	201	TRS	C3-C-N	4.40	116.10	108.09
3	D	201	TRS	O3-C3-C	4.82	120.93	111.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	199	HEC	4	0
2	A	200	HEC	2	0
2	B	199	HEC	1	0
2	B	200	HEC	2	0
4	C	201	GOL	2	0
2	D	200	HEC	1	0
3	D	201	TRS	7	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/190 (100%)	-0.12	3 (1%) 74 75	7, 14, 28, 47	0
1	B	190/190 (100%)	0.32	10 (5%) 30 29	10, 24, 40, 50	0
1	C	190/190 (100%)	0.07	6 (3%) 51 52	8, 14, 27, 54	0
1	D	190/190 (100%)	0.39	17 (8%) 12 12	9, 17, 33, 47	0
All	All	760/760 (100%)	0.16	36 (4%) 35 35	7, 17, 35, 54	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	ALA	5.9
1	D	53	PRO	5.6
1	D	92[A]	VAL	5.2
1	B	25	PRO	4.2
1	D	132	LEU	3.8
1	B	92	VAL	3.7
1	B	113	ASP	3.6
1	D	57	GLU	3.5
1	B	133	ALA	3.5
1	D	56	PRO	3.0
1	D	6	ALA	3.0
1	D	94	TYR	2.8
1	D	112	LEU	2.7
1	D	59	VAL	2.6
1	A	113	ASP	2.6
1	B	57	GLU	2.6
1	A	57	GLU	2.5
1	A	163	ASP	2.5
1	D	5	GLU	2.5
1	D	72	PRO	2.4
1	C	4	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	5	GLU	2.4
1	D	95	ALA	2.4
1	D	55	ALA	2.3
1	C	94	TYR	2.3
1	B	52	THR	2.2
1	D	74	SER	2.2
1	B	54	GLY	2.2
1	B	94	TYR	2.1
1	D	97	PRO	2.1
1	B	1	ALA	2.1
1	D	3	ASP	2.1
1	C	57	GLU	2.1
1	D	64	LEU	2.1
1	B	166	ILE	2.0
1	C	2	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	C	201	6/6	0.86	0.43	18.22	29,55,56,73	0
3	TRS	D	201	8/8	0.81	0.26	11.44	30,36,50,57	0
2	HEC	C	200	43/43	0.99	0.09	0.62	6,9,13,15	0
2	HEC	D	200	43/43	0.99	0.09	0.52	7,9,11,13	0
2	HEC	A	199	43/43	0.99	0.08	0.12	7,9,13,30	0
2	HEC	A	200	43/43	0.99	0.07	-0.02	6,7,13,16	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEC	B	199	43/43	0.99	0.08	-0.07	10,13,18,32	0
2	HEC	C	199	43/43	0.99	0.07	-0.10	6,8,13,30	0
2	HEC	B	200	43/43	0.99	0.08	-0.17	9,13,20,22	0
4	GOL	A	201	6/6	0.95	0.08	-0.21	14,15,18,18	0
2	HEC	D	199	43/43	0.99	0.08	-0.51	8,12,15,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.