



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:37 PM GMT

PDB ID : 4M6W
Title : Crystal structure of the C-terminal segment of FANCM in complex with FAAP24
Authors : Yang, H.; Zhang, T.; Tong, L.; Ding, J.
Deposited on : 2013-08-11
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
2	B	208	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	2102	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fanconi anemia group M protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1569	989	272	296	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1811	MET	-	EXPRESSION TAG	UNP Q8IYD8
A	1812	GLY	-	EXPRESSION TAG	UNP Q8IYD8

- Molecule 2 is a protein called Fanconi anemia-associated protein of 24 kDa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	1489	952	258	272	7	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	EXPRESSION TAG	UNP Q9BTP7
B	216	LEU	-	EXPRESSION TAG	UNP Q9BTP7
B	217	GLU	-	EXPRESSION TAG	UNP Q9BTP7
B	218	HIS	-	EXPRESSION TAG	UNP Q9BTP7
B	219	HIS	-	EXPRESSION TAG	UNP Q9BTP7
B	220	HIS	-	EXPRESSION TAG	UNP Q9BTP7
B	221	HIS	-	EXPRESSION TAG	UNP Q9BTP7
B	222	HIS	-	EXPRESSION TAG	UNP Q9BTP7
B	223	HIS	-	EXPRESSION TAG	UNP Q9BTP7

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	3	Total	O	0	0
			3	3		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fanconi anemia group M protein



- Molecule 2: Fanconi anemia-associated protein of 24 kDa



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	79.17Å 98.11Å 140.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.90 39.97 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.3 (40.00-2.90) 97.4 (39.97-2.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.08 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.254 , 0.299 0.254 , 0.270	Depositor DCC
R_{free} test set	593 reflections (5.13%)	DCC
Wilson B-factor (Å ²)	60.5	Xtriage
Anisotropy	0.605	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 12152 reflections (0.008%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3081	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1590	0.57	0/2140
2	B	0.38	0/1511	0.54	0/2042
All	All	0.38	0/3101	0.56	0/4182

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1569	0	1588	20	0
2	B	1489	0	1552	10	0
3	A	10	0	0	2	0
3	B	5	0	0	0	0
4	A	5	0	0	0	0
4	B	3	0	0	0	0
All	All	3081	0	3140	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1961:VAL:HG22	1:A:1962:PRO:HD2	1.71	0.71
1:A:1860:ARG:HD3	1:A:1957:VAL:HG21	1.77	0.66
1:A:1920:SER:O	1:A:1924:THR:HG23	1.99	0.62
1:A:1821:LEU:HB2	1:A:1856:ILE:HB	1.87	0.57
1:A:1818:THR:HG23	1:A:1953:GLN:HE21	1.72	0.55
1:A:1827:ILE:HD11	1:A:1847:VAL:HG11	1.88	0.54
2:B:174:ALA:HB3	2:B:175:PRO:HD3	1.89	0.53
1:A:1947:GLU:HB3	2:B:110:LEU:CD2	2.39	0.53
2:B:22:ASN:HA	2:B:46:LEU:HD21	1.92	0.51
1:A:1893:ARG:NH1	3:A:2102:SO4:O4	2.33	0.51
2:B:119:VAL:HG13	2:B:121:SER:O	2.12	0.49
2:B:193:GLY:O	2:B:197:GLN:NE2	2.46	0.48
1:A:1860:ARG:HG3	1:A:1892:GLU:HB2	1.97	0.47
1:A:1853:CYS:HB3	1:A:1887:LEU:HD21	1.97	0.46
2:B:167:PRO:HA	2:B:168:GLY:HA2	1.71	0.46
1:A:1895:CYS:SG	1:A:1933:LEU:HD23	2.56	0.45
1:A:1924:THR:HB	2:B:143:ASN:HB2	1.99	0.45
1:A:1955:LYS:NZ	3:A:2102:SO4:O4	2.49	0.45
1:A:1976:TYR:HD2	1:A:1991:CYS:HG	1.66	0.44
1:A:1818:THR:HG21	1:A:1949:SER:HB3	2.00	0.43
1:A:1819:CYS:HB3	1:A:1959:ILE:HG23	2.01	0.43
1:A:1818:THR:O	1:A:1843:LEU:HB2	2.18	0.43
1:A:1926:ILE:HD11	2:B:117:LEU:HD21	2.01	0.42
1:A:1998:LYS:HB2	2:B:211:PHE:HB3	2.02	0.42
1:A:1895:CYS:SG	1:A:1933:LEU:CD2	3.09	0.41
2:B:163:VAL:HG21	2:B:178:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/221 (87%)	179 (93%)	13 (7%)	0	100	100
2	B	186/208 (89%)	174 (94%)	10 (5%)	2 (1%)	17	51
All	All	378/429 (88%)	353 (93%)	23 (6%)	2 (0%)	34	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	24	LYS
2	B	35	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/201 (90%)	156 (87%)	24 (13%)	5	14
2	B	167/182 (92%)	149 (89%)	18 (11%)	8	24
All	All	347/383 (91%)	305 (88%)	42 (12%)	6	18

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1816	LYS
1	A	1828	THR
1	A	1831	LEU
1	A	1843	LEU
1	A	1845	VAL
1	A	1850	LEU
1	A	1857	VAL
1	A	1860	ARG
1	A	1896	VAL
1	A	1918	TYR
1	A	1921	LEU
1	A	1924	THR
1	A	1931	ARG
1	A	1932	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1944	LEU
1	A	1948	LEU
1	A	1961	VAL
1	A	1973	LEU
1	A	2005	LEU
1	A	2008	ILE
1	A	2015	THR
1	A	2016	HIS
1	A	2017	GLN
1	A	2022	ILE
2	B	30	LEU
2	B	35	GLN
2	B	39	LYS
2	B	44	ASP
2	B	47	THR
2	B	55	ARG
2	B	66	LEU
2	B	94	THR
2	B	104	LEU
2	B	136	GLN
2	B	138	LYS
2	B	142	LYS
2	B	145	LEU
2	B	153	LEU
2	B	154	LEU
2	B	172	VAL
2	B	173	LYS
2	B	197	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1953	GLN
2	B	197	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	2101	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	A	2102	-	4,4,4	0.32	0	6,6,6	0.12	0
3	SO4	B	301	-	4,4,4	0.31	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	2101	-	-	0/0/0/0	0/0/0/0
3	SO4	A	2102	-	-	0/0/0/0	0/0/0/0
3	SO4	B	301	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2102	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/221 (89%)	0.19	4 (2%) 68 64	45, 59, 86, 101	0
2	B	190/208 (91%)	0.50	17 (8%) 12 7	48, 74, 101, 125	0
All	All	388/429 (90%)	0.34	21 (5%) 29 23	45, 68, 98, 125	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	68	ALA	7.9
2	B	140	PRO	3.4
2	B	70	ASN	3.2
1	A	2005	LEU	3.2
2	B	96	MET	3.0
2	B	35	GLN	2.8
2	B	81	ASN	2.8
2	B	213	GLN	2.7
2	B	202	ALA	2.7
2	B	69	GLY	2.5
2	B	54	ASN	2.4
2	B	181	PHE	2.4
1	A	1815	GLY	2.4
2	B	139	GLU	2.3
2	B	38	ILE	2.2
1	A	1962	PRO	2.1
2	B	198	VAL	2.1
2	B	197	GLN	2.1
1	A	1970	SER	2.1
2	B	212	THR	2.0
2	B	72	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	2101	5/5	0.98	0.17	-0.44	80,80,82,82	0
3	SO4	B	301	5/5	0.96	0.11	-1.63	102,103,104,105	0
3	SO4	A	2102	5/5	0.77	0.48	-	52,52,53,54	5

6.5 Other polymers [i](#)

There are no such residues in this entry.