



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:40 PM GMT

PDB ID : 4MAB  
Title : Resolving Cys to Ala variant of Salmonella Alkyl Hydroperoxide Reductase C  
in its substrate-ready conformation  
Authors : Perkins, A.; Nelson, K.J.; Williams, J.R.; Poole, L.B.; Karplus, P.A.  
Deposited on : 2013-08-15  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	<div> <div>16%</div> <div>84%</div> <div>11%</div> </div>
1	B	186	<div> <div>24%</div> <div>94%</div> <div>6%</div> </div>
1	C	186	<div> <div>25%</div> <div>95%</div> <div>5%</div> </div>
1	D	186	<div> <div>20%</div> <div>96%</div> <div>.</div> </div>
1	E	186	<div> <div>36%</div> <div>95%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	K	D	201	-	-	-	X
3	GOL	A	202[B]	-	-	-	X
3	GOL	A	203	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14454 atoms, of which 7024 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkyl hydroperoxide reductase subunit C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	165	Total	C	H	N	O	S	0	1	0
			2555	826	1257	216	253	3			
1	B	186	Total	C	H	N	O	S	0	11	0
			2977	965	1470	248	290	4			
1	C	186	Total	C	H	N	O	S	0	3	0
			2876	928	1421	241	283	3			
1	D	186	Total	C	H	N	O	S	0	3	0
			2877	928	1422	241	283	3			
1	E	186	Total	C	H	N	O	S	0	4	0
			2877	928	1422	241	283	3			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	ALA	CYS	ENGINEERED MUTATION	UNP P0A251
B	165	ALA	CYS	ENGINEERED MUTATION	UNP P0A251
C	165	ALA	CYS	ENGINEERED MUTATION	UNP P0A251
D	165	ALA	CYS	ENGINEERED MUTATION	UNP P0A251
E	165	ALA	CYS	ENGINEERED MUTATION	UNP P0A251

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		
2	A	1	Total	K	0	0
			1	1		
2	D	1	Total	K	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	1
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		
3	D	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	66	Total	O	0	1
			66	66		
5	B	61	Total	O	0	1
			61	61		
5	C	47	Total	O	0	0
			47	47		
5	D	48	Total	O	0	2
			48	48		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	10	Total	O	0	0
			10	10		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alkyl hydroperoxide reductase subunit C




- Molecule 1: Alkyl hydroperoxide reductase subunit C




- Molecule 1: Alkyl hydroperoxide reductase subunit C



- Molecule 1: Alkyl hydroperoxide reductase subunit C



● Molecule 1: Alkyl hydroperoxide reductase subunit C





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.23Å 172.42Å 136.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.15 – 1.90 29.15 – 1.90	Depositor EDS
% Data completeness (in resolution range)	81.8 (29.15-1.90) 81.8 (29.15-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.23 (at 1.91Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.198 , 0.239 0.209 , 0.241	Depositor DCC
$R_{free}$ test set	4793 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.6	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 96108 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14454	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.95	0/1328	0.93	0/1800
1	B	0.94	0/1553	0.92	3/2107 (0.1%)
1	C	0.86	0/1489	0.90	2/2018 (0.1%)
1	D	0.81	0/1489	0.84	1/2018 (0.0%)
1	E	0.63	0/1489	0.77	0/2018
All	All	0.85	0/7348	0.87	6/9961 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	98	ASP	CB-CG-OD2	-7.94	111.16	118.30
1	C	105	ARG	NE-CZ-NH2	-7.62	116.49	120.30
1	B	31	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	D	105	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	B	55	ASP	CB-CG-OD2	-5.42	113.42	118.30
1	B	31	ARG	NE-CZ-NH2	-5.09	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1298	1257	1252	6	0
1	B	1507	1470	1460	8	0
1	C	1455	1421	1417	7	0
1	D	1455	1422	1417	7	0
1	E	1455	1422	1417	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	12	16	16	1	0
3	C	6	8	8	0	0
3	D	6	8	8	0	0
4	D	1	0	0	1	0
5	A	66	0	0	0	0
5	B	61	0	0	1	0
5	C	47	0	0	0	0
5	D	48	0	0	0	0
5	E	10	0	0	0	0
All	All	7430	7024	6995	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:46[B]:CYS:SG	5:B:1052:HOH:O	2.43	0.75
1:B:31:ARG:HD3	1:B:66:ASP:OD2	1.97	0.64
1:A:128:GLY:HA3	3:A:203:GOL:H2	1.91	0.52
1:D:1:SER:HB2	4:D:202:CL:CL	2.47	0.52
1:E:125:ASP:HB2	1:E:126:PRO:CD	2.42	0.50
1:C:180:LEU:HD13	1:E:19:GLU:HA	1.95	0.49
1:A:164:VAL:HG23	1:A:165:ALA:H	1.80	0.47
1:A:39:PRO:HD2	1:A:46:CYS:SG	2.56	0.46
1:A:42:PHE:O	1:B:183:VAL:HG13	2.15	0.46
1:C:125:ASP:HB2	1:C:126:PRO:CD	2.46	0.46
1:B:46[B]:CYS:N	1:B:47[B]:PRO:HD2	2.32	0.45
1:C:167:ALA:HB2	1:D:49:GLU:HG2	1.99	0.44
1:E:120:ALA:HA	1:E:134:GLU:O	2.17	0.43
1:B:109:ASN:OD1	1:B:118:ASP:HB2	2.19	0.43
1:E:32:TRP:CE3	1:E:125:ASP:HA	2.54	0.42
1:A:157:VAL:HG11	1:B:139:GLY:HA3	2.02	0.42
1:B:79:LYS:HE2	1:D:180:LEU:O	2.20	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:164[A]:VAL:HG21	1:D:45:VAL:HG23	2.02	0.42
1:B:125:ASP:HB2	1:B:126:PRO:CD	2.49	0.42
1:C:42:PHE:O	1:D:183:VAL:HG13	2.20	0.41
1:C:45:VAL:HG23	1:D:164[B]:VAL:HG21	2.02	0.40
1:E:10:PHE:O	1:E:24:THR:HA	2.21	0.40
1:A:59:GLU:OE2	1:A:148:LEU:HD21	2.22	0.40
1:C:45:VAL:CG2	1:D:164[B]:VAL:HG21	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/186 (88%)	158 (97%)	4 (2%)	1 (1%)	30	17
1	B	192/186 (103%)	187 (97%)	5 (3%)	0	100	100
1	C	184/186 (99%)	179 (97%)	5 (3%)	0	100	100
1	D	184/186 (99%)	179 (97%)	5 (3%)	0	100	100
1	E	184/186 (99%)	179 (97%)	5 (3%)	0	100	100
All	All	907/930 (98%)	882 (97%)	24 (3%)	1 (0%)	56	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/153 (90%)	137 (100%)	0	100	100
1	B	160/153 (105%)	160 (100%)	0	100	100
1	C	153/153 (100%)	152 (99%)	1 (1%)	88	88
1	D	153/153 (100%)	153 (100%)	0	100	100
1	E	153/153 (100%)	151 (99%)	2 (1%)	76	73
All	All	756/765 (99%)	753 (100%)	3 (0%)	93	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	22	GLU
1	E	86	GLU
1	E	156	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	202[B]	-	5,5,5	0.41	0	5,5,5	0.73	0
3	GOL	A	203	-	5,5,5	0.46	0	5,5,5	0.90	0
3	GOL	C	201	-	5,5,5	0.36	0	5,5,5	0.32	0
3	GOL	D	203	-	5,5,5	0.40	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	202[B]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	203	-	-	0/4/4/4	0/0/0/0
3	GOL	C	201	-	-	0/4/4/4	0/0/0/0
3	GOL	D	203	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	203	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	165/186 (88%)	0.89	29 (17%)	2 2	31, 43, 74, 117	1 (0%)
1	B	186/186 (100%)	1.13	44 (23%)	1 1	29, 44, 81, 100	24 (12%)
1	C	186/186 (100%)	1.15	46 (24%)	1 1	32, 51, 84, 106	24 (12%)
1	D	186/186 (100%)	1.19	37 (19%)	1 1	35, 50, 84, 109	24 (12%)
1	E	186/186 (100%)	1.90	67 (36%)	0 0	46, 73, 107, 138	24 (12%)
All	All	909/930 (97%)	1.26	223 (24%)	1 1	29, 52, 95, 138	97 (10%)

All (223) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	176	LEU	10.7
1	D	182	LEU	10.2
1	E	171	GLU	8.5
1	D	179	SER	8.5
1	B	182	LEU	8.2
1	C	182	LEU	7.9
1	E	169	TRP	7.8
1	E	172	GLY	7.7
1	E	164[B]	VAL	7.6
1	D	183	VAL	7.6
1	E	36	PHE	7.5
1	E	162	GLY	7.3
1	E	177	ALA	7.3
1	C	183	VAL	7.1
1	B	170	LYS	7.0
1	D	169	TRP	6.7
1	E	174	ALA	6.7
1	D	171	GLU	6.6
1	E	170	LYS	6.6
1	E	70	VAL	6.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	180	LEU	6.3
1	D	178	PRO	6.2
1	E	178	PRO	6.1
1	C	170	LYS	6.0
1	D	174	ALA	5.9
1	E	156	TYR	5.8
1	C	174	ALA	5.8
1	E	161	PRO	5.7
1	C	186	ILE	5.7
1	E	183	VAL	5.7
1	B	183	VAL	5.6
1	D	172	GLY	5.5
1	A	45	VAL	5.4
1	D	180	LEU	5.4
1	E	180	LEU	5.4
1	B	180	LEU	5.4
1	B	174	ALA	5.4
1	D	167	ALA	5.3
1	D	184	GLY	5.3
1	B	185	LYS	5.3
1	E	159	ALA	5.3
1	E	44	PHE	5.3
1	D	175	THR	5.3
1	A	164	VAL	5.3
1	D	170	LYS	5.3
1	B	179	SER	5.2
1	C	172	GLY	5.2
1	B	171	GLU	5.1
1	E	69	SER	5.1
1	D	186	ILE	5.1
1	B	176	LEU	5.1
1	B	162	GLY	5.0
1	E	34	VAL	5.0
1	E	182	LEU	5.0
1	D	181	ASP	5.0
1	D	177	ALA	5.0
1	A	122	PHE	4.9
1	A	121	THR	4.8
1	A	70	VAL	4.7
1	E	35	PHE	4.7
1	B	169	TRP	4.7
1	E	71	SER	4.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	177	ALA	4.6
1	A	120	ALA	4.6
1	D	162	GLY	4.6
1	A	36	PHE	4.6
1	E	121	THR	4.5
1	A	35	PHE	4.5
1	B	161	PRO	4.5
1	E	173	GLU	4.5
1	B	177	ALA	4.5
1	E	62	LYS	4.5
1	C	185	LYS	4.4
1	D	185	LYS	4.4
1	E	157	VAL	4.4
1	B	178	PRO	4.4
1	E	184	GLY	4.3
1	E	181	ASP	4.3
1	E	179	SER	4.3
1	E	45	VAL	4.3
1	E	67	VAL	4.3
1	E	38	TYR	4.2
1	E	46	CYS	4.2
1	C	165[A]	ALA	4.2
1	E	37	PHE	4.1
1	E	68	TYR	4.0
1	C	169	TRP	4.0
1	B	46[A]	CYS	4.0
1	B	168	LYS	4.0
1	C	171	GLU	4.0
1	E	185	LYS	3.9
1	C	37	PHE	3.8
1	A	38	TYR	3.8
1	C	121	THR	3.8
1	A	46	CYS	3.8
1	E	148	LEU	3.8
1	A	68	TYR	3.7
1	A	37	PHE	3.7
1	B	186	ILE	3.7
1	B	172	GLY	3.7
1	A	34	VAL	3.6
1	C	70	VAL	3.6
1	D	161	PRO	3.6
1	A	44	PHE	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	122	PHE	3.6
1	A	133	ILE	3.6
1	E	165[B]	ALA	3.5
1	B	121	THR	3.4
1	E	175	THR	3.4
1	C	36	PHE	3.4
1	C	161	PRO	3.3
1	E	168	LYS	3.3
1	E	176	LEU	3.3
1	D	36	PHE	3.3
1	D	34	VAL	3.2
1	C	181	ASP	3.2
1	C	122	PHE	3.2
1	C	62	LYS	3.2
1	B	173	GLU	3.2
1	E	58	GLU	3.2
1	C	156	TYR	3.2
1	E	96	ILE	3.1
1	D	120	ALA	3.1
1	D	165[B]	ALA	3.1
1	B	37	PHE	3.1
1	B	86	GLU	3.1
1	C	176	LEU	3.0
1	C	34	VAL	3.0
1	C	164[A]	VAL	3.0
1	E	160	HIS	3.0
1	D	86	GLU	3.0
1	E	86	GLU	3.0
1	A	69	SER	3.0
1	A	96	ILE	3.0
1	B	35	PHE	3.0
1	B	70	VAL	3.0
1	C	168	LYS	3.0
1	C	45	VAL	3.0
1	B	165[B]	ALA	2.9
1	D	121	THR	2.9
1	E	113	ASP	2.9
1	B	38	TYR	2.9
1	A	165	ALA	2.9
1	A	67	VAL	2.9
1	A	123	VAL	2.9
1	B	133	ILE	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	122	PHE	2.8
1	B	68	TYR	2.8
1	B	44[A]	PHE	2.7
1	E	186	ILE	2.7
1	E	120	ALA	2.7
1	A	71	SER	2.7
1	B	96	ILE	2.7
1	D	133	ILE	2.7
1	C	120	ALA	2.7
1	E	112	GLU	2.6
1	B	69	SER	2.6
1	A	132	ALA	2.6
1	B	71	SER	2.6
1	E	33	SER	2.6
1	C	44	PHE	2.6
1	A	135	VAL	2.6
1	E	146	ASP	2.6
1	B	36	PHE	2.6
1	D	168	LYS	2.6
1	C	162	GLY	2.6
1	E	135	VAL	2.5
1	C	73	ASP	2.5
1	A	134	GLU	2.5
1	D	45	VAL	2.5
1	D	38	TYR	2.5
1	C	63	LEU	2.5
1	C	69	SER	2.5
1	C	35	PHE	2.5
1	A	94	ALA	2.5
1	C	86	GLU	2.5
1	D	173	GLU	2.5
1	B	166[B]	PRO	2.5
1	C	38	TYR	2.5
1	E	73	ASP	2.5
1	D	135	VAL	2.4
1	C	179	SER	2.4
1	D	46	CYS	2.4
1	B	159	ALA	2.4
1	C	184	GLY	2.4
1	C	167	ALA	2.4
1	B	184	GLY	2.4
1	C	58	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	94	ALA	2.4
1	E	123	VAL	2.4
1	C	173	GLU	2.3
1	E	63	LEU	2.3
1	B	120	ALA	2.3
1	C	71	SER	2.3
1	E	163	GLU	2.3
1	C	133	ILE	2.3
1	E	20	PHE	2.3
1	A	74	THR	2.3
1	D	122	PHE	2.3
1	A	33	SER	2.3
1	D	148	LEU	2.3
1	A	92	LYS	2.3
1	E	167[B]	ALA	2.3
1	B	97	GLY	2.2
1	A	107	PHE	2.2
1	B	175	THR	2.2
1	B	167	ALA	2.2
1	C	74	THR	2.2
1	B	45[A]	VAL	2.2
1	E	92	LYS	2.2
1	B	135	VAL	2.2
1	E	134	GLU	2.1
1	E	153	ALA	2.1
1	C	68	TYR	2.1
1	C	67	VAL	2.1
1	E	90	LYS	2.1
1	E	59	GLU	2.1
1	C	46	CYS	2.1
1	D	40	ALA	2.0
1	E	136	THR	2.0
1	D	166[B]	PRO	2.0
1	E	41	ASP	2.0
1	B	181	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	K	D	201	1/1	0.71	0.62	9.45	103,103,103,103	0
3	GOL	A	202[B]	6/6	0.82	0.62	6.51	45,68,106,121	14
3	GOL	A	203	6/6	0.69	0.47	5.56	71,77,116,121	0
3	GOL	C	201	6/6	0.78	0.21	1.72	14,14,85,86	0
2	K	B	201	1/1	0.85	0.25	1.41	79,79,79,79	0
3	GOL	D	203	6/6	0.78	0.16	0.01	14,14,87,88	0
4	CL	D	202	1/1	0.90	0.08	-2.22	68,68,68,68	0
2	K	A	201	1/1	0.69	0.20	-	98,98,98,98	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.