



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:43 PM GMT

PDB ID : 4MD8
Title : Crystal Structure of full-length symmetric CK2 holoenzyme with mutated alpha subunit (F121E)
Authors : Lolli, G.; Ranchio, A.; Battistutta, R.
Deposited on : 2013-08-22
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 76% 15% • 7%
1	B	215	 74% 15% • 9%
1	C	215	 76% 13% • 8%
1	D	215	 74% 15% • 9%
2	E	391	 67% 14% • • 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	391	<p>2% 66% 14% •• 15%</p>
2	G	391	<p>% 66% 15% •• 15%</p>
2	H	391	<p>% 65% 15% •• 15%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 17617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	199	1622	1040	273	294	15	0	0	0
1	B	195	1592	1024	268	285	15	0	0	0
1	C	197	1605	1031	271	288	15	0	0	0
1	D	195	1592	1024	268	285	15	0	0	0

- Molecule 2 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	332	2802	1791	495	505	11	0	0	0
2	F	332	2802	1791	495	505	11	0	0	0
2	G	332	2802	1791	495	505	11	0	0	0
2	H	331	2796	1788	494	503	11	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	121	GLU	PHE	ENGINEERED MUTATION	UNP P68400
F	121	GLU	PHE	ENGINEERED MUTATION	UNP P68400
G	121	GLU	PHE	ENGINEERED MUTATION	UNP P68400
H	121	GLU	PHE	ENGINEERED MUTATION	UNP P68400


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Zn 1	0	0
3	A	1	Total 1	Zn 1	0	0
3	D	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0


• Molecule 2: Casein kinase II subunit alpha



• Molecule 2: Casein kinase II subunit alpha




• Molecule 2: Casein kinase II subunit alpha



• Molecule 2: Casein kinase II subunit alpha





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	141.52Å 57.62Å 185.69Å 90.00° 102.60° 90.00°	Depositor
Resolution (Å)	181.21 – 3.30 47.65 – 3.30	Depositor EDS
% Data completeness (in resolution range)	98.2 (181.21-3.30) 98.5 (47.65-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.04 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.222 , 0.249 0.221 , 0.248	Depositor DCC
R_{free} test set	2230 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	57.0	Xtriage
Anisotropy	0.411	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 27.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	1 of 44213 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	17617	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	2/1671 (0.1%)	0.89	6/2265 (0.3%)
1	B	0.73	1/1641 (0.1%)	0.85	4/2223 (0.2%)
1	C	0.69	1/1654 (0.1%)	0.79	4/2242 (0.2%)
1	D	0.65	1/1641 (0.1%)	0.79	4/2223 (0.2%)
2	E	0.80	2/2876 (0.1%)	0.89	7/3889 (0.2%)
2	F	0.72	1/2876 (0.0%)	0.88	9/3889 (0.2%)
2	G	0.72	3/2876 (0.1%)	0.84	6/3889 (0.2%)
2	H	0.72	3/2870 (0.1%)	0.87	8/3881 (0.2%)
All	All	0.73	14/18105 (0.1%)	0.86	48/24501 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1
2	G	0	1
2	H	0	1
All	All	0	3

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	12	TRP	CD2-CE2	6.17	1.48	1.41
1	D	9	TRP	CD2-CE2	6.01	1.48	1.41
2	F	216	TRP	CD2-CE2	5.83	1.48	1.41
2	H	24	TRP	CD2-CE2	5.70	1.48	1.41
1	C	12	TRP	CD2-CE2	5.66	1.48	1.41

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	47	ARG	NE-CZ-NH1	11.94	126.27	120.30
1	D	47	ARG	NE-CZ-NH2	-11.42	114.59	120.30
2	H	278	ARG	NE-CZ-NH1	11.37	125.99	120.30
1	B	47	ARG	NE-CZ-NH1	11.11	125.86	120.30
1	A	47	ARG	NE-CZ-NH2	10.80	125.70	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	120	ASP	Peptide
2	G	118	ASN	Peptide
2	H	119	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1622	0	1534	27	0
1	B	1592	0	1517	28	0
1	C	1605	0	1524	26	0
1	D	1592	0	1517	30	2
2	E	2802	0	2745	71	0
2	F	2802	0	2745	90	0
2	G	2802	0	2745	68	2
2	H	2796	0	2740	78	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	17617	0	17067	394	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 394 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:118:ASN:HD22	2:F:163:MET:CE	1.53	1.22
2:H:121:GLU:HA	2:H:121:GLU:OE1	1.36	1.16
2:E:275:ARG:HH11	2:E:275:ARG:HG3	1.13	1.11
2:F:118:ASN:ND2	2:F:163:MET:HE3	1.67	1.09
2:F:124:LEU:HD11	2:F:128:LEU:HD21	1.30	1.09

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:148:SER:OG	2:G:271:ASP:OD2[2_546]	2.06	0.14
1:D:17:ARG:NE	2:G:271:ASP:O[2_546]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/215 (91%)	190 (97%)	5 (3%)	0	100	100
1	B	191/215 (89%)	186 (97%)	5 (3%)	0	100	100
1	C	193/215 (90%)	187 (97%)	6 (3%)	0	100	100
1	D	191/215 (89%)	185 (97%)	5 (3%)	1 (0%)	34	71
2	E	330/391 (84%)	310 (94%)	20 (6%)	0	100	100
2	F	330/391 (84%)	305 (92%)	25 (8%)	0	100	100
2	G	330/391 (84%)	302 (92%)	26 (8%)	2 (1%)	30	68
2	H	329/391 (84%)	304 (92%)	24 (7%)	1 (0%)	46	81
All	All	2089/2424 (86%)	1969 (94%)	116 (6%)	4 (0%)	52	85

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	37	ASP
1	D	67	ASN
2	G	22	GLU
2	G	121	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	174/191 (91%)	168 (97%)	6 (3%)	44 77
1	B	171/191 (90%)	166 (97%)	5 (3%)	50 80
1	C	172/191 (90%)	164 (95%)	8 (5%)	32 70
1	D	171/191 (90%)	164 (96%)	7 (4%)	37 74
2	E	305/347 (88%)	286 (94%)	19 (6%)	23 61
2	F	305/347 (88%)	285 (93%)	20 (7%)	21 59
2	G	305/347 (88%)	284 (93%)	21 (7%)	19 57
2	H	304/347 (88%)	283 (93%)	21 (7%)	19 57
All	All	1907/2152 (89%)	1800 (94%)	107 (6%)	26 65

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	47	ARG
2	F	172	ARG
2	H	122	LYS
2	F	52	GLU
2	F	123	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	18	HIS
2	F	118	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	123	GLN
2	F	40	GLN
2	F	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	199/215 (92%)	-0.35	0 100 100	23, 40, 76, 104	0
1	B	195/215 (90%)	-0.21	2 (1%) 84 80	27, 58, 98, 140	0
1	C	197/215 (91%)	-0.07	2 (1%) 84 80	38, 59, 93, 153	0
1	D	195/215 (90%)	-0.04	2 (1%) 84 80	38, 65, 104, 152	0
2	E	332/391 (84%)	-0.30	1 (0%) 94 94	29, 44, 78, 113	0
2	F	332/391 (84%)	0.06	6 (1%) 71 65	44, 69, 108, 134	0
2	G	332/391 (84%)	0.10	3 (0%) 85 82	46, 73, 104, 123	0
2	H	331/391 (84%)	0.04	5 (1%) 76 71	50, 70, 106, 138	0
All	All	2113/2424 (87%)	-0.08	21 (0%) 84 80	23, 62, 101, 153	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	118	ASN	3.3
2	H	126	GLN	3.0
1	C	69	SER	3.0
1	D	59	ASP	2.9
2	F	125	TYR	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ZN	A	301	1/1	1.00	0.07	-2.40	29,29,29,29	0
3	ZN	B	301	1/1	0.99	0.08	-2.48	32,32,32,32	0
3	ZN	C	301	1/1	0.99	0.07	-2.54	47,47,47,47	0
3	ZN	D	301	1/1	1.00	0.08	-3.40	40,40,40,40	0

6.5 Other polymers

There are no such residues in this entry.