



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:40 PM GMT

PDB ID : 4MDD
Title : Crystal Structure of the Glucocorticoid Receptor Bound to a Non-steroidal Antagonist Reveals Repositioning and Partial Disordering of Activation Function Helix 12
Authors : Coghlan, M.J.; Luz, J.G.
Deposited on : 2013-08-22
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>6%</div> <div>89% 7% .</div> </div>
1	B	258	<div> <div>6%</div> <div>77% 8% 15%</div> </div>
2	C	15	<div> <div>13%</div> <div>80% 7% 13%</div> </div>
2	D	15	<div> <div>27%</div> <div>60% 20% 20%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucocorticoid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	0	0
			1989	1278	327	367	17			
1	B	220	Total	C	N	O	S	0	0	0
			1780	1144	297	324	15			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	520	SER	-	EXPRESSION TAG	UNP P04150
A	521	SER	-	EXPRESSION TAG	UNP P04150
A	525	SER	LEU	ENGINEERED MUTATION	UNP P04150
A	528	SER	LEU	ENGINEERED MUTATION	UNP P04150
A	535	ALA	LEU	ENGINEERED MUTATION	UNP P04150
A	538	THR	VAL	ENGINEERED MUTATION	UNP P04150
A	602	TYR	PHE	ENGINEERED MUTATION	UNP P04150
A	638	ASP	CYS	ENGINEERED MUTATION	UNP P04150
A	684	ALA	GLU	ENGINEERED MUTATION	UNP P04150
A	688	ALA	GLU	ENGINEERED MUTATION	UNP P04150
A	712	SER	TRP	ENGINEERED MUTATION	UNP P04150
B	520	SER	-	EXPRESSION TAG	UNP P04150
B	521	SER	-	EXPRESSION TAG	UNP P04150
B	525	SER	LEU	ENGINEERED MUTATION	UNP P04150
B	528	SER	LEU	ENGINEERED MUTATION	UNP P04150
B	535	ALA	LEU	ENGINEERED MUTATION	UNP P04150
B	538	THR	VAL	ENGINEERED MUTATION	UNP P04150
B	602	TYR	PHE	ENGINEERED MUTATION	UNP P04150
B	638	ASP	CYS	ENGINEERED MUTATION	UNP P04150
B	684	ALA	GLU	ENGINEERED MUTATION	UNP P04150
B	688	ALA	GLU	ENGINEERED MUTATION	UNP P04150
B	712	SER	TRP	ENGINEERED MUTATION	UNP P04150

- Molecule 2 is a protein called Nuclear receptor corepressor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	13	Total	C	N	O	S	0	0	0
			98	63	17	17	1			
2	D	12	Total	C	N	O	S	0	0	0
			90	57	16	16	1			

- Molecule 3 is N-[2-{[BENZYL(METHYL)AMINO]METHYL}-3-(4-FLUORO-2-METHOXYPHENYL)-5-(PROPAN-2-YL)-1H-INDOL-7-YL]METHANESULFONAMIDE (three-letter code: 29M) (formula: C₂₈H₃₂FN₃O₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	0
			36	28	1	3	3	1		
3	B	1	Total	C	F	N	O	S	0	0
			36	28	1	3	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	66	Total	O	0	0
			66	66		
4	B	43	Total	O	0	0
			43	43		
4	C	2	Total	O	0	0
			2	2		
4	D	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucocorticoid receptor




- Molecule 1: Glucocorticoid receptor



- Molecule 2: Nuclear receptor corepressor 1



- Molecule 2: Nuclear receptor corepressor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	72.54Å 72.54Å 229.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.80 – 2.40 19.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (19.80-2.40) 99.6 (19.80-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.41 (at 2.41Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.221 , 0.258 0.224 , 0.261	Depositor DCC
R_{free} test set	870 reflections (3.19%)	DCC
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.2	EDS
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 28177 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4141	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 29M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2031	0.64	0/2747
1	B	0.47	0/1817	0.63	0/2456
2	C	0.46	0/97	0.68	0/127
2	D	0.50	0/89	0.58	0/116
All	All	0.47	0/4034	0.63	0/5446

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1989	0	1999	6	0
1	B	1780	0	1778	8	0
2	C	98	0	113	0	0
2	D	90	0	99	2	0
3	A	36	0	32	3	0
3	B	36	0	32	0	0
4	A	66	0	0	0	0
4	B	43	0	0	1	0
4	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	1	0	0	0	0
All	All	4141	0	4053	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:591:ASP:HA	1:B:594:THR:HG22	1.80	0.62
1:B:578:ALA:HA	1:B:581:ILE:HD12	1.87	0.57
3:A:801:29M:H2	3:A:801:29M:H26	1.87	0.56
1:B:593:MET:HE3	2:D:8:ARG:HA	1.87	0.56
1:A:743:LYS:HA	1:A:747:ILE:HD12	1.87	0.56
1:A:577:TRP:CZ2	1:A:581:ILE:HD11	2.42	0.54
1:A:578:ALA:HA	1:A:581:ILE:HD12	1.90	0.52
1:B:662:GLU:OE2	1:B:704:ARG:HD2	2.15	0.47
3:A:801:29M:C1	3:A:801:29M:H26	2.46	0.45
1:B:720:LYS:HG2	1:B:775:HIS:CD2	2.52	0.44
1:B:611:ARG:NH2	4:B:911:HOH:O	2.50	0.44
1:B:588:HIS:HD2	1:B:590:ASP:HB2	1.82	0.44
1:A:740:PHE:CZ	1:A:745:MET:HE3	2.54	0.42
1:A:610:TRP:HB2	1:A:663:TYR:CE1	2.55	0.42
1:B:578:ALA:HA	1:B:581:ILE:CD1	2.49	0.41
2:D:8:ARG:O	2:D:12:MET:HG2	2.21	0.41
1:A:639:MET:HE2	3:A:801:29M:H32	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/258 (95%)	241 (98%)	4 (2%)	0	100	100
1	B	216/258 (84%)	213 (99%)	3 (1%)	0	100	100
2	C	11/15 (73%)	11 (100%)	0	0	100	100
2	D	10/15 (67%)	10 (100%)	0	0	100	100
All	All	482/546 (88%)	475 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/234 (95%)	215 (97%)	7 (3%)	46	68
1	B	196/234 (84%)	187 (95%)	9 (5%)	33	51
2	C	10/12 (83%)	9 (90%)	1 (10%)	9	14
2	D	9/12 (75%)	8 (89%)	1 (11%)	8	10
All	All	437/492 (89%)	419 (96%)	18 (4%)	37	57

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	613	TYR
1	A	631	GLU
1	A	641	ASP
1	A	642	GLN
1	A	753	LEU
1	A	760	GLN
1	A	766	ASN
1	B	543	VAL
1	B	556	THR
1	B	613	TYR
1	B	636	LEU
1	B	650	SER
1	B	674	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	689	ILE
1	B	731	ASN
1	B	739	THR
2	C	3	LEU
2	D	5	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	597	GLN
1	A	760	GLN
1	B	586	ASN
1	B	597	GLN
1	B	615	GLN
1	B	711	ASN
1	B	775	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	29M	A	801	-	36,39,39	1.62	4 (11%)	47,57,57	1.81	7 (14%)
3	29M	B	801	-	36,39,39	1.52	4 (11%)	47,57,57	1.54	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	29M	A	801	-	-	0/22/23/23	0/4/4/4
3	29M	B	801	-	-	0/22/23/23	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	29M	C8-C5	-4.07	1.45	1.50
3	B	801	29M	C8-C5	-3.55	1.46	1.50
3	B	801	29M	C15-C4	-2.72	1.48	1.51
3	A	801	29M	C15-C4	-2.68	1.48	1.51
3	A	801	29M	C2-N2	2.17	1.44	1.42
3	B	801	29M	O-C13	2.70	1.41	1.37
3	B	801	29M	S-N2	6.09	1.71	1.63
3	A	801	29M	S-N2	6.65	1.72	1.63

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	29M	C24-S-N2	-6.21	99.00	106.83
3	A	801	29M	C-C7-C6	-5.04	118.44	121.93
3	A	801	29M	C7-C-C25	-4.65	115.07	121.67
3	B	801	29M	C-C7-C6	-3.93	119.21	121.93
3	B	801	29M	C7-C-C25	-3.44	116.79	121.67
3	B	801	29M	C24-S-N2	-2.79	103.31	106.83
3	B	801	29M	C14-O-C13	-2.77	113.34	117.54
3	A	801	29M	C14-O-C13	-2.63	113.56	117.54
3	B	801	29M	C16-N1-C17	2.08	114.00	110.50
3	A	801	29M	O2-S-N2	2.24	111.91	107.17
3	A	801	29M	C7-C-C1	2.28	121.60	118.13
3	A	801	29M	C3-C2-N2	3.53	120.98	115.77
3	B	801	29M	C3-C2-N2	4.90	123.00	115.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	29M	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/258 (95%)	0.21	16 (6%) 22 22	28, 46, 87, 115	0
1	B	220/258 (85%)	0.19	15 (6%) 20 20	27, 51, 86, 101	0
2	C	13/15 (86%)	0.83	2 (15%) 3 3	46, 62, 79, 86	0
2	D	12/15 (80%)	1.51	4 (33%) 0 1	68, 81, 93, 98	0
All	All	492/546 (90%)	0.25	37 (7%) 17 17	27, 50, 89, 115	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	1	LEU	5.5
2	D	2	GLY	4.5
2	D	3	LEU	4.4
1	A	766	ASN	4.3
1	A	764	TYR	4.3
1	B	769	ILE	4.3
1	A	632	GLN	3.9
1	B	740	PHE	3.8
1	B	741	LEU	3.7
1	A	530	PRO	3.7
1	A	762	PRO	3.6
1	A	705	GLU	3.6
1	A	704	ARG	3.6
1	B	638	ASP	3.3
1	A	765	SER	3.3
1	B	632	GLN	3.3
2	D	4	GLU	3.3
1	A	635	THR	3.2
1	A	763	LYS	3.0
1	A	761	ILE	2.9
1	A	713	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	588	HIS	2.9
2	C	2	GLY	2.8
1	B	640	TYR	2.7
1	A	752	MET	2.7
1	B	633	ARG	2.6
1	A	531	THR	2.5
1	A	634	MET	2.4
1	B	637	PRO	2.2
1	A	638	ASP	2.2
1	B	776	GLN	2.2
1	B	768	ASN	2.1
2	D	13	GLY	2.1
1	B	645	HIS	2.1
1	B	678	ASP	2.1
1	B	592	GLN	2.1
1	B	596	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	29M	B	801	36/36	0.94	0.18	0.54	23,35,51,52	0
3	29M	A	801	36/36	0.94	0.18	0.34	25,32,42,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.