



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:57 PM GMT

PDB ID : 1MG3  
Title : MUTATION OF ALPHA PHE55 OF METHYLAMINE DEHYDROGENASE ALTERS THE REORGANIZATION ENERGY AND ELECTRONIC COUPLING FOR ITS ELECTRON TRANSFER REACTION WITH AMICYANIN  
Authors : Sun, D.; Chen, Z.W.; Mathews, F.S.; Davidson, V.L.  
Deposited on : 2002-08-14  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	<div> <div>2%</div> <div>63% 32% ..</div> </div>
1	E	390	<div> <div>%</div> <div>61% 34% ..</div> </div>
1	I	390	<div> <div>2%</div> <div>64% 31% ..</div> </div>
1	M	390	<div> <div>%</div> <div>64% 31% ..</div> </div>
2	B	131	<div> <div>%</div> <div>63% 31% • 5%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	131	
2	J	131	
2	N	131	
3	C	105	
3	G	105	
3	K	105	
3	O	105	
4	D	155	
4	H	155	
4	L	155	
4	P	155	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PO4	A	408	-	-	-	X
6	PO4	B	402	-	-	-	X
6	PO4	E	406	-	-	-	X
6	PO4	F	401	-	-	X	X
6	PO4	I	405	-	-	-	X
6	PO4	J	403	-	-	X	X
6	PO4	M	407	-	-	-	X
6	PO4	O	404	-	-	X	X
7	NA	H	604	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 24562 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylamine dehydrogenase, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	0	0
			2961	1872	509	572	8			
1	E	382	Total	C	N	O	S	0	0	0
			2961	1872	509	572	8			
1	I	382	Total	C	N	O	S	0	0	0
			2961	1872	509	572	8			
1	M	382	Total	C	N	O	S	0	0	0
			2961	1872	509	572	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	ALA	PHE	ENGINEERED	UNP P29894
A	312	PHE	LEU	SEE REMARK 999	UNP P29894
A	313	VAL	LEU	SEE REMARK 999	UNP P29894
E	55	ALA	PHE	ENGINEERED	UNP P29894
E	312	PHE	LEU	SEE REMARK 999	UNP P29894
E	313	VAL	LEU	SEE REMARK 999	UNP P29894
I	55	ALA	PHE	ENGINEERED	UNP P29894
I	312	PHE	LEU	SEE REMARK 999	UNP P29894
I	313	VAL	LEU	SEE REMARK 999	UNP P29894
M	55	ALA	PHE	ENGINEERED	UNP P29894
M	312	PHE	LEU	SEE REMARK 999	UNP P29894
M	313	VAL	LEU	SEE REMARK 999	UNP P29894

- Molecule 2 is a protein called Methylamine dehydrogenase, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	125	Total	C	N	O	S	0	0	0
			963	596	163	191	13			
2	F	125	Total	C	N	O	S	0	0	0
			963	596	163	191	13			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	125	Total	C	N	O	S	0	0	0
			963	596	163	191	13			
2	N	125	Total	C	N	O	S	0	0	0
			963	596	163	191	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	57	TRW	TRP	MODIFIED RESIDUE	UNP P22619
F	57	TRW	TRP	MODIFIED RESIDUE	UNP P22619
J	57	TRW	TRP	MODIFIED RESIDUE	UNP P22619
N	57	TRW	TRP	MODIFIED RESIDUE	UNP P22619

- Molecule 3 is a protein called Amicyanin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	105	Total	C	N	O	S	0	0	0
			807	516	133	152	6			
3	G	105	Total	C	N	O	S	0	0	0
			807	516	133	152	6			
3	K	105	Total	C	N	O	S	0	0	0
			807	516	133	152	6			
3	O	105	Total	C	N	O	S	0	0	0
			807	516	133	152	6			

- Molecule 4 is a protein called CYTOCHROME C-L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	147	Total	C	N	O	S	0	0	0
			1144	724	182	230	8			
4	H	147	Total	C	N	O	S	0	0	0
			1144	724	182	230	8			
4	L	147	Total	C	N	O	S	0	0	0
			1144	724	182	230	8			
4	P	147	Total	C	N	O	S	0	0	0
			1144	724	182	230	8			

- Molecule 5 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	Cu	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	O	1	Total 1	Cu 1	0	0
5	C	1	Total 1	Cu 1	0	0
5	K	1	Total 1	Cu 1	0	0

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	F	1	Total 5	O 4	P 1	0	0
6	B	1	Total 5	O 4	P 1	0	0
6	J	1	Total 5	O 4	P 1	0	0
6	O	1	Total 5	O 4	P 1	0	0
6	I	1	Total 5	O 4	P 1	0	0
6	E	1	Total 5	O 4	P 1	0	0
6	M	1	Total 5	O 4	P 1	0	0
6	A	1	Total 5	O 4	P 1	0	0

- | Mol | Chain | Residues | Atoms           | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 7   | P     | 1        | Total Na<br>1 1 | 0       | 0       |
| 7   | D     | 1        | Total Na<br>1 1 | 0       | 0       |
| 7   | L     | 1        | Total Na<br>1 1 | 0       | 0       |
| 7   | H     | 1        | Total Na<br>1 1 | 0       | 0       |

- 
- Chemical structure of HEM (Heme) showing a central iron atom coordinated by four nitrogen atoms in a porphyrin-like ring, with various side chains and a central heme group.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	H	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	L	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	P	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- 
- WORLD WIDE  
PDB  
PROTEIN DATA BANK


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	123	Total 123	O 123	0	0
9	B	20	Total 20	O 20	0	0
9	C	9	Total 9	O 9	0	0
9	D	13	Total 13	O 13	0	0
9	E	113	Total 113	O 113	0	0
9	F	51	Total 51	O 51	0	0
9	G	35	Total 35	O 35	0	0
9	H	21	Total 21	O 21	0	0
9	I	140	Total 140	O 140	0	0
9	J	49	Total 49	O 49	0	0
9	K	36	Total 36	O 36	0	0
9	L	22	Total 22	O 22	0	0
9	M	139	Total 139	O 139	0	0
9	N	20	Total 20	O 20	0	0
9	O	13	Total 13	O 13	0	0
9	P	38	Total 38	O 38	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methylamine dehydrogenase, heavy chain




- Molecule 1: Methylamine dehydrogenase, heavy chain




- Molecule 1: Methylamine dehydrogenase, heavy chain









- Molecule 2: Methylamine dehydrogenase, light chain




- Molecule 3: Amicyanin




- Molecule 3: Amicyanin



- Molecule 3: Amicyanin




- Molecule 3: Amicyanin







• Molecule 4: CYTOCHROME C-L



• Molecule 4: CYTOCHROME C-L



• Molecule 4: CYTOCHROME C-L



• Molecule 4: CYTOCHROME C-L



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.52Å 188.37Å 127.37Å 90.00° 98.84° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 49.05 – 2.24	Depositor EDS
% Data completeness (in resolution range)	81.5 (50.00-2.40) 72.1 (49.05-2.24)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 2.24Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.193 , 0.246 0.193 , 0.242	Depositor DCC
$R_{free}$ test set	11668 reflections (11.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.7	Xtriage
Anisotropy	0.526	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 46.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 127220 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	24562	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PO4, TRW, CU, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/3037	0.64	0/4139
1	E	0.35	0/3037	0.64	0/4139
1	I	0.36	0/3037	0.66	0/4139
1	M	0.35	0/3037	0.65	0/4139
2	B	0.37	0/964	0.62	0/1315
2	F	0.37	0/964	0.61	0/1315
2	J	0.37	0/964	0.61	0/1315
2	N	0.36	0/964	0.61	0/1315
3	C	0.31	0/828	0.54	0/1124
3	G	0.37	0/828	0.63	0/1124
3	K	0.36	0/828	0.60	0/1124
3	O	0.36	0/828	0.57	0/1124
4	D	0.38	1/1179 (0.1%)	0.65	1/1605 (0.1%)
4	H	0.36	0/1179	0.67	1/1605 (0.1%)
4	L	0.36	0/1179	0.66	1/1605 (0.1%)
4	P	0.38	1/1179 (0.1%)	0.66	1/1605 (0.1%)
All	All	0.36	2/24032 (0.0%)	0.64	4/32732 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	61	HIS	CE1-NE2	5.48	1.45	1.32
4	D	61	HIS	CE1-NE2	5.22	1.44	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	61	HIS	ND1-CG-CD2	8.13	120.19	108.80
4	D	61	HIS	ND1-CG-CD2	8.05	120.07	108.80
4	L	61	HIS	ND1-CG-CD2	8.00	120.00	108.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	61	HIS	ND1-CG-CD2	7.93	119.90	108.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2961	0	2843	141	0
1	E	2961	0	2843	120	0
1	I	2961	0	2843	127	0
1	M	2961	0	2843	128	0
2	B	963	0	876	50	0
2	F	963	0	876	46	0
2	J	963	0	876	44	0
2	N	963	0	876	54	0
3	C	807	0	794	31	0
3	G	807	0	794	15	0
3	K	807	0	794	15	0
3	O	807	0	794	22	0
4	D	1144	0	1038	32	0
4	H	1144	0	1038	37	0
4	L	1144	0	1038	38	0
4	P	1144	0	1038	42	0
5	C	1	0	0	0	0
5	G	1	0	0	0	0
5	K	1	0	0	0	0
5	O	1	0	0	0	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
6	E	5	0	0	1	0
6	F	5	0	0	3	0
6	I	5	0	0	1	0
6	J	5	0	0	3	0
6	M	5	0	0	1	0
6	O	5	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	1	0	0	0	0
7	H	1	0	0	0	0
7	L	1	0	0	0	0
7	P	1	0	0	0	0
8	D	43	0	30	0	0
8	H	43	0	30	2	0
8	L	43	0	30	2	0
8	P	43	0	30	3	0
9	A	123	0	0	5	0
9	B	20	0	0	3	0
9	C	9	0	0	0	0
9	D	13	0	0	1	0
9	E	113	0	0	4	0
9	F	51	0	0	0	0
9	G	35	0	0	0	0
9	H	21	0	0	0	0
9	I	140	0	0	5	0
9	J	49	0	0	1	0
9	K	36	0	0	0	0
9	L	22	0	0	2	0
9	M	139	0	0	10	0
9	N	20	0	0	2	0
9	O	13	0	0	0	0
9	P	38	0	0	0	0
All	All	24562	0	22324	842	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 842 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:ARG:HB2	1:A:20:ARG:HH21	1.19	1.07
2:J:21:GLN:HE22	1:M:11:GLN:HG3	1.28	0.98
2:B:21:GLN:HE22	1:E:11:GLN:HG2	1.26	0.96
1:E:173:LYS:HE2	1:E:173:LYS:HA	1.50	0.93
1:I:35:ARG:H	2:N:45:ASN:HD22	1.16	0.92

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/390 (97%)	352 (93%)	26 (7%)	2 (0%)	34	48
1	E	380/390 (97%)	353 (93%)	24 (6%)	3 (1%)	24	35
1	I	380/390 (97%)	357 (94%)	20 (5%)	3 (1%)	24	35
1	M	380/390 (97%)	360 (95%)	19 (5%)	1 (0%)	46	63
2	B	122/131 (93%)	110 (90%)	12 (10%)	0	100	100
2	F	122/131 (93%)	109 (89%)	13 (11%)	0	100	100
2	J	122/131 (93%)	109 (89%)	13 (11%)	0	100	100
2	N	122/131 (93%)	108 (88%)	14 (12%)	0	100	100
3	C	103/105 (98%)	91 (88%)	11 (11%)	1 (1%)	19	28
3	G	103/105 (98%)	98 (95%)	5 (5%)	0	100	100
3	K	103/105 (98%)	99 (96%)	4 (4%)	0	100	100
3	O	103/105 (98%)	88 (85%)	10 (10%)	5 (5%)	3	1
4	D	145/155 (94%)	126 (87%)	15 (10%)	4 (3%)	6	5
4	H	145/155 (94%)	129 (89%)	14 (10%)	2 (1%)	14	19
4	L	145/155 (94%)	133 (92%)	10 (7%)	2 (1%)	14	19
4	P	145/155 (94%)	133 (92%)	6 (4%)	6 (4%)	3	3
All	All	3000/3124 (96%)	2755 (92%)	216 (7%)	29 (1%)	19	28

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	131	ALA
3	O	18	ASP
1	A	102	ILE
4	D	68	LYS
1	E	342	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/311 (99%)	292 (95%)	15 (5%)	31	48
1	E	307/311 (99%)	292 (95%)	15 (5%)	31	48
1	I	307/311 (99%)	290 (94%)	17 (6%)	27	42
1	M	307/311 (99%)	292 (95%)	15 (5%)	31	48
2	B	104/106 (98%)	100 (96%)	4 (4%)	40	60
2	F	104/106 (98%)	100 (96%)	4 (4%)	40	60
2	J	104/106 (98%)	99 (95%)	5 (5%)	31	49
2	N	104/106 (98%)	100 (96%)	4 (4%)	40	60
3	C	85/85 (100%)	81 (95%)	4 (5%)	32	50
3	G	85/85 (100%)	80 (94%)	5 (6%)	24	38
3	K	85/85 (100%)	82 (96%)	3 (4%)	43	64
3	O	85/85 (100%)	81 (95%)	4 (5%)	32	50
4	D	118/125 (94%)	114 (97%)	4 (3%)	44	65
4	H	118/125 (94%)	110 (93%)	8 (7%)	20	31
4	L	118/125 (94%)	109 (92%)	9 (8%)	16	25
4	P	118/125 (94%)	110 (93%)	8 (7%)	20	31
All	All	2456/2508 (98%)	2332 (95%)	124 (5%)	30	48

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	H	128	PRO
1	I	242	TRP
3	O	48	ARG
4	H	134	LEU
1	I	46	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
4	H	14	ASN
1	I	82	ASN
2	N	45	ASN
1	I	11	GLN
1	I	288	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	TRW	B	57	2	22,25,25	2.48	7 (31%)	19,34,34	2.08	7 (36%)
2	TRW	F	57	2	22,25,25	2.52	7 (31%)	19,34,34	1.98	6 (31%)
2	TRW	J	57	2	22,25,25	2.51	7 (31%)	19,34,34	2.00	8 (42%)
2	TRW	N	57	2	22,25,25	2.52	7 (31%)	19,34,34	2.00	7 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRW	B	57	2	-	0/8/11/11	0/3/3/3
2	TRW	F	57	2	-	0/8/11/11	0/3/3/3
2	TRW	J	57	2	-	0/8/11/11	0/3/3/3
2	TRW	N	57	2	-	0/8/11/11	0/3/3/3

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	57	TRW	O7-CZ2	-3.27	1.23	1.35
2	B	57	TRW	O7-CZ2	-3.13	1.24	1.35
2	F	57	TRW	O7-CZ2	-2.85	1.25	1.35
2	N	57	TRW	O7-CZ2	-2.78	1.25	1.35
2	B	57	TRW	C4-C3	2.21	1.43	1.38

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	57	TRW	C6-C1-C2	-3.32	114.45	119.06
2	J	57	TRW	C6-C1-C2	-3.26	114.53	119.06
2	B	57	TRW	C6-C1-C2	-3.21	114.60	119.06
2	N	57	TRW	C6-C1-C2	-3.17	114.65	119.06
2	B	57	TRW	O7-CZ2-CE2	-2.55	115.22	119.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	57	TRW	4	0
2	F	57	TRW	3	0
2	J	57	TRW	3	0
2	N	57	TRW	3	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PO4	A	408	-	4,4,4	0.87	0	6,6,6	0.27	0
6	PO4	B	402	-	4,4,4	1.07	0	6,6,6	0.27	0
8	HEM	D	200	4	30,50,50	2.75	6 (20%)	24,82,82	2.99	10 (41%)
6	PO4	E	406	-	4,4,4	1.06	0	6,6,6	0.27	0
6	PO4	F	401	-	4,4,4	1.08	0	6,6,6	0.27	0
8	HEM	H	200	4	30,50,50	2.63	7 (23%)	24,82,82	2.89	9 (37%)
6	PO4	I	405	-	4,4,4	1.09	0	6,6,6	0.27	0
6	PO4	J	403	-	4,4,4	1.07	0	6,6,6	0.27	0
8	HEM	L	200	4	30,50,50	2.84	7 (23%)	24,82,82	2.84	9 (37%)
6	PO4	M	407	-	4,4,4	1.08	0	6,6,6	0.27	0
6	PO4	O	404	-	4,4,4	0.98	0	6,6,6	0.27	0
8	HEM	P	200	4	30,50,50	2.75	7 (23%)	24,82,82	2.93	12 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PO4	A	408	-	-	0/0/0/0	0/0/0/0
6	PO4	B	402	-	-	0/0/0/0	0/0/0/0
8	HEM	D	200	4	-	0/10/54/54	0/0/8/8
6	PO4	E	406	-	-	0/0/0/0	0/0/0/0
6	PO4	F	401	-	-	0/0/0/0	0/0/0/0
8	HEM	H	200	4	-	0/10/54/54	0/0/8/8
6	PO4	I	405	-	-	0/0/0/0	0/0/0/0
6	PO4	J	403	-	-	0/0/0/0	0/0/0/0
8	HEM	L	200	4	-	0/10/54/54	0/0/8/8
6	PO4	M	407	-	-	0/0/0/0	0/0/0/0
6	PO4	O	404	-	-	0/0/0/0	0/0/0/0
8	HEM	P	200	4	-	0/10/54/54	0/0/8/8

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	L	200	HEM	C3B-C4B	-10.09	1.42	1.51
8	D	200	HEM	C3B-C4B	-9.25	1.43	1.51
8	H	200	HEM	C3B-C4B	-8.85	1.44	1.51
8	P	200	HEM	C3B-C4B	-8.84	1.44	1.51
8	D	200	HEM	C2D-C3D	-6.31	1.35	1.54

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	D	200	HEM	C3B-CAB-CBB	-6.85	113.96	124.46
8	H	200	HEM	C3B-CAB-CBB	-6.83	113.97	124.46
8	P	200	HEM	C3B-CAB-CBB	-6.63	114.28	124.46
8	L	200	HEM	C3B-CAB-CBB	-5.88	115.43	124.46
8	D	200	HEM	C3C-CAC-CBC	-5.76	115.62	124.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	406	PO4	1	0
6	F	401	PO4	3	0
8	H	200	HEM	2	0
6	I	405	PO4	1	0
6	J	403	PO4	3	0
8	L	200	HEM	2	0
6	M	407	PO4	1	0
6	O	404	PO4	2	0
8	P	200	HEM	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	382/390 (97%)	-0.30	8 (2%) 67 66	10, 26, 48, 90	0
1	E	382/390 (97%)	-0.32	5 (1%) 79 79	9, 28, 50, 87	0
1	I	382/390 (97%)	-0.39	6 (1%) 74 74	10, 24, 49, 78	0
1	M	382/390 (97%)	-0.46	2 (0%) 91 91	9, 23, 41, 72	0
2	B	124/131 (94%)	-0.30	1 (0%) 87 87	18, 26, 42, 79	0
2	F	124/131 (94%)	-0.32	2 (1%) 74 74	13, 20, 37, 82	0
2	J	124/131 (94%)	-0.37	1 (0%) 87 87	13, 20, 34, 80	0
2	N	124/131 (94%)	-0.29	1 (0%) 87 87	14, 25, 40, 83	0
3	C	105/105 (100%)	0.90	23 (21%) 1 1	30, 53, 75, 86	0
3	G	105/105 (100%)	-0.41	3 (2%) 55 54	12, 28, 47, 62	0
3	K	105/105 (100%)	-0.40	2 (1%) 70 69	10, 24, 40, 62	0
3	O	105/105 (100%)	0.20	9 (8%) 13 13	23, 39, 67, 77	0
4	D	147/155 (94%)	-0.17	1 (0%) 89 88	22, 40, 66, 84	0
4	H	147/155 (94%)	-0.08	4 (2%) 58 57	18, 38, 70, 79	0
4	L	147/155 (94%)	-0.15	2 (1%) 78 77	18, 39, 63, 74	0
4	P	147/155 (94%)	-0.34	4 (2%) 58 57	14, 32, 60, 82	0
All	All	3032/3124 (97%)	-0.26	74 (2%) 62 61	9, 27, 60, 90	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	63	GLY	4.9
1	E	5	GLU	4.7
1	I	208	THR	4.6
2	F	131	SER	4.5
1	E	386	GLY	4.4

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	TRW	J	57	23/23	0.96	0.14	-	13,20,34,39	0
2	TRW	F	57	23/23	0.98	0.11	-	9,17,31,35	0
2	TRW	B	57	23/23	0.95	0.14	-	16,26,38,39	0
2	TRW	N	57	23/23	0.96	0.13	-	12,22,37,42	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	PO4	E	406	5/5	0.83	0.35	15.08	84,87,91,94	0
6	PO4	I	405	5/5	0.93	0.26	12.13	78,78,82,83	0
6	PO4	M	407	5/5	0.83	0.24	9.96	80,83,89,90	0
6	PO4	A	408	5/5	0.83	0.24	7.80	72,75,79,88	0
6	PO4	F	401	5/5	0.90	0.27	7.46	86,86,88,89	0
6	PO4	J	403	5/5	0.72	0.31	6.01	84,85,89,90	0
6	PO4	O	404	5/5	0.87	0.29	3.33	89,91,93,94	0
7	NA	H	604	1/1	0.95	0.18	2.43	23,23,23,23	0
6	PO4	B	402	5/5	0.94	0.22	2.12	74,74,76,76	0
7	NA	L	602	1/1	0.98	0.14	1.88	21,21,21,21	0
7	NA	D	603	1/1	0.78	0.17	1.10	40,40,40,40	0
8	HEM	D	200	43/43	0.96	0.15	0.45	29,38,54,68	0
8	HEM	L	200	43/43	0.98	0.13	0.43	18,28,35,46	0
7	NA	P	601	1/1	0.98	0.11	0.23	10,10,10,10	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	HEM	H	200	43/43	0.97	0.13	0.23	20,29,32,35	0
8	HEM	P	200	43/43	0.98	0.11	-0.30	11,23,28,31	0
5	CU	K	107	1/1	1.00	0.09	-0.58	31,31,31,31	0
5	CU	C	107	1/1	0.99	0.07	-2.04	56,56,56,56	0
5	CU	G	107	1/1	1.00	0.07	-2.28	33,33,33,33	0
5	CU	O	107	1/1	0.99	0.08	-2.89	44,44,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.