



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 03:05 AM BST

PDB ID : 2MGT  
Title : Zinc induced dimer of the metal binding domain 1-16 of human amyloid beta-peptide with Alzheimer's disease pathogenic English mutation H6R  
Authors : Polshakov, V.; Istrate, A.; Kozin, S.  
Deposited on : 2013-11-06

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 40%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	18	
1	B	18	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:16, B:1-B:16 (32)	1.21	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 7, 8, 10, 13, 16, 20
2	3, 4, 11, 15, 18
3	2, 9, 12, 19
4	6, 17
Single-model clusters	5; 14

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 539 atoms, of which 252 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Amyloid beta A4 protein.

Mol	Chain	Residues	Atoms					Trace
1	A	18	Total	C	H	N	O	1
			269	86	126	29	28	
1	B	18	Total	C	H	N	O	1
			269	86	126	29	28	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ACE	-	ACETYLATION	UNP P05067
A	6	ARG	HIS	ENGINEERED MUTATION	UNP P05067
A	17	NH2	-	AMIDATION	UNP P05067
B	0	ACE	-	ACETYLATION	UNP P05067
B	6	ARG	HIS	ENGINEERED MUTATION	UNP P05067
B	17	NH2	-	AMIDATION	UNP P05067

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Amyloid beta A4 protein



- Molecule 1: Amyloid beta A4 protein




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




### 4.2.2 Score per residue for model 2

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




### 4.2.3 Score per residue for model 3

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




### 4.2.4 Score per residue for model 4

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.5 Score per residue for model 5

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.6 Score per residue for model 6

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.7 Score per residue for model 7

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.8 Score per residue for model 8

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.9 Score per residue for model 9

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.10 Score per residue for model 10

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein






#### 4.2.11 Score per residue for model 11

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein



#### 4.2.12 Score per residue for model 12

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.13 Score per residue for model 13

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.14 Score per residue for model 14

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.15 Score per residue for model 15

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.16 Score per residue for model 16

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.17 Score per residue for model 17

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.18 Score per residue for model 18

- Molecule 1: Amyloid beta A4 protein




- Molecule 1: Amyloid beta A4 protein




#### 4.2.19 Score per residue for model 19

- Molecule 1: Amyloid beta A4 protein



- Molecule 1: Amyloid beta A4 protein




#### 4.2.20 Score per residue for model 20

- Molecule 1: Amyloid beta A4 protein

Chain A:  72% 17% 11%



- Molecule 1: Amyloid beta A4 protein

Chain B:  78% 6% 6% 11%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
GROMACS	refinement	4.5.4
CPMD	geometry optimization	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mgt_cs.str
Number of chemical shift lists	3
Total number of shifts	528
Number of shifts mapped to atoms	528
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	40%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.72±0.25	0±1/142 (0.1±0.4%)	1.13±0.05	0±0/188 (0.1±0.2%)
1	B	0.73±0.23	0±1/142 (0.2±0.6%)	1.13±0.07	0±0/188 (0.1±0.2%)
All	All	0.77	11/5680 (0.2%)	1.13	9/7520 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.7±0.7
1	B	0.0±0.0	0.5±0.6
All	All	0	22

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	3	GLU	CD-OE2	-14.06	1.10	1.25	18	1
1	A	3	GLU	CD-OE1	11.37	1.38	1.25	18	1
1	B	11	GLU	CD-OE1	-10.35	1.14	1.25	1	1
1	B	11	GLU	CD-OE2	9.77	1.36	1.25	1	2
1	A	11	GLU	CD-OE2	-9.61	1.15	1.25	1	1
1	B	3	GLU	CD-OE2	-8.75	1.16	1.25	3	2
1	A	11	GLU	CD-OE1	8.53	1.35	1.25	1	1
1	B	3	GLU	CD-OE1	-8.11	1.16	1.25	12	2

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	ARG	NE-CZ-NH1	5.84	123.22	120.30	20	4
1	A	5	ARG	NE-CZ-NH1	5.55	123.08	120.30	4	1
1	B	6	ARG	NE-CZ-NH1	5.33	122.96	120.30	5	3
1	B	5	ARG	NE-CZ-NH1	5.05	122.82	120.30	18	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	10	TYR	Peptide	11
1	B	10	TYR	Peptide,Sidechain	7
1	B	5	ARG	Sidechain	1
1	A	2	ALA	Peptide	1
1	A	5	ARG	Sidechain	1

## 6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	139	121	121	0±0
1	B	139	121	121	0±0
All	All	5580	4840	4840	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:14:HIS:CE1	1:B:15:GLN:HE21	0.45	2.30	17	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	16/18 (89%)	13±1 (84±6%)	2±1 (15±6%)	0±0 (1±3%)	20	66
1	B	16/18 (89%)	12±1 (76±8%)	4±1 (23±8%)	0±0 (1±2%)	26	73
All	All	640/720 (89%)	512 (80%)	121 (19%)	7 (1%)	23	69

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	5	ARG	3
1	B	14	HIS	2
1	B	6	ARG	1
1	A	8	SER	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	14/14 (100%)	13±0 (90±3%)	1±0 (10±3%)	15	60
1	B	14/14 (100%)	13±1 (90±4%)	1±1 (10±4%)	15	60
All	All	560/560 (100%)	506 (90%)	54 (10%)	15	60

All 12 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	TYR	20
1	B	10	TYR	20
1	A	1	ASP	3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	B	5	ARG	2
1	A	5	ARG	2
1	B	7	ASP	1
1	A	16	LYS	1
1	A	15	GLN	1
1	B	11	GLU	1
1	B	15	GLN	1
1	B	1	ASP	1
1	B	14	HIS	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 40% for the well-defined parts and 40% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2mgt\_cs.str

Chemical shift list name: *Assigned\_chemical\_shifts\_2*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	172
Number of shifts mapped to atoms	172
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	16	$-0.17 \pm 0.11$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	12	$0.23 \pm 0.35$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	11	$-1.38 \pm 0.49$	Should be applied

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 37%, i.e. 159 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	59/160 (37%)	32/64 (50%)	16/64 (25%)	11/32 (34%)
Sidechain	78/208 (38%)	54/124 (44%)	24/68 (35%)	0/16 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	22/62 (35%)	14/34 (41%)	8/24 (33%)	0/4 (0%)
Overall	159/430 (37%)	100/222 (45%)	48/156 (31%)	11/52 (21%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 37%, i.e. 159 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	59/160 (37%)	32/64 (50%)	16/64 (25%)	11/32 (34%)
Sidechain	78/208 (38%)	54/124 (44%)	24/68 (35%)	0/16 (0%)
Aromatic	22/62 (35%)	14/34 (41%)	8/24 (33%)	0/4 (0%)
Overall	159/430 (37%)	100/222 (45%)	48/156 (31%)	11/52 (21%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



## 7.2 Chemical shift list 2

File name: 2mgt\_cs.str

Chemical shift list name: *Assigned\_chemical\_shifts\_3*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	174
Number of shifts mapped to atoms	174
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	16	$-0.22 \pm 0.10$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	14	$0.28 \pm 0.19$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	11	$-1.48 \pm 0.34$	Should be applied

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 37%, i.e. 161 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	59/160 (37%)	32/64 (50%)	16/64 (25%)	11/32 (34%)
Sidechain	80/208 (38%)	54/124 (44%)	26/68 (38%)	0/16 (0%)
Aromatic	22/62 (35%)	14/34 (41%)	8/24 (33%)	0/4 (0%)
Overall	161/430 (37%)	100/222 (45%)	50/156 (32%)	11/52 (21%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 37%, i.e. 161 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	59/160 (37%)	32/64 (50%)	16/64 (25%)	11/32 (34%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Sidechain	80/208 (38%)	54/124 (44%)	26/68 (38%)	0/16 (0%)
Aromatic	22/62 (35%)	14/34 (41%)	8/24 (33%)	0/4 (0%)
Overall	161/430 (37%)	100/222 (45%)	50/156 (32%)	11/52 (21%)


#### 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



### 7.3 Chemical shift list 3

File name: 2mgt\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.3.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	182
Number of shifts mapped to atoms	182

Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.3.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	16	$-0.16 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	15	$0.14 \pm 0.10$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	16	$-1.24 \pm 0.42$	Should be applied

### 7.3.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 40%, i.e. 171 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	64/160 (40%)	32/64 (50%)	16/64 (25%)	16/32 (50%)
Sidechain	81/208 (39%)	54/124 (44%)	27/68 (40%)	0/16 (0%)
Aromatic	26/62 (42%)	14/34 (41%)	12/24 (50%)	0/4 (0%)
Overall	171/430 (40%)	100/222 (45%)	55/156 (35%)	16/52 (31%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 40%, i.e. 171 atoms were assigned a chemical shift out of a possible 430. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	64/160 (40%)	32/64 (50%)	16/64 (25%)	16/32 (50%)
Sidechain	81/208 (39%)	54/124 (44%)	27/68 (40%)	0/16 (0%)
Aromatic	26/62 (42%)	14/34 (41%)	12/24 (50%)	0/4 (0%)
Overall	171/430 (40%)	100/222 (45%)	55/156 (35%)	16/52 (31%)

### 7.3.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

### 7.3.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

