



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 02:47 AM BST

PDB ID : 2MJC  
Title : Zn-binding domain of eukaryotic translation initiation factor 3, subunit G  
Authors : Al-Abdul-Wahid, M.; Menade, M.; Xie, J.; Kozlov, G.; Gehring, K.  
Deposited on : 2014-01-03

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 78%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	34	

## 2 Ensemble composition and analysis ⓘ

This entry contains 15 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:26 (16)	0.22	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 14, 15
2	10, 13

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 384 atoms, of which 191 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 3 subunit G.

Mol	Chain	Residues	Atoms						Trace
1	A	24	Total	C	H	N	O	S	0
			383	120	191	36	33	3	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	PRO	-	EXPRESSION TAG	UNP O75821
A	3	LEU	-	EXPRESSION TAG	UNP O75821
A	4	GLY	-	EXPRESSION TAG	UNP O75821
A	5	SER	-	EXPRESSION TAG	UNP O75821

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Eukaryotic translation initiation factor 3 subunit G



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: Eukaryotic translation initiation factor 3 subunit G



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	structure solution	
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mjc_cs.str
Number of chemical shift lists	1
Total number of shifts	345
Number of shifts mapped to atoms	345
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	78%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	127	120	121	2±2
All	All	1920	1800	1815	29

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 18 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:14:ARG:O	1:A:14:ARG:NE	0.52	2.42	5	1
1:A:21:TRP:O	1:A:23:THR:N	0.51	2.43	13	2
1:A:14:ARG:C	1:A:14:ARG:NE	0.47	2.67	5	1
1:A:13:CYS:SG	1:A:14:ARG:N	0.47	2.87	3	2
1:A:11:VAL:HG23	1:A:21:TRP:CD1	0.47	2.45	3	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	16/34 (47%)	10±1 (60±7%)	5±1 (32±7%)	1±1 (8±5%)	3	16
All	All	240/510 (47%)	145 (60%)	77 (32%)	18 (8%)	3	16

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	21	TRP	12
1	A	13	CYS	5
1	A	26	PRO	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	15/30 (50%)	15±0 (97±3%)	0±0 (3±3%)	56	92
All	All	225/450 (50%)	219 (97%)	6 (3%)	56	92

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	23	THR	2
1	A	20	HIS	2
1	A	14	ARG	1
1	A	17	LYS	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 78% for the well-defined parts and 80% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2mjc\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	345
Number of shifts mapped to atoms	345
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	56

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	34	$0.00 \pm 0.00$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	29	$1.04 \pm 0.50$	Should be applied
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	31	$-0.52 \pm 0.66$	None needed (imprecise)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 156 atoms were assigned a chemical shift out of a possible 199. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	62/78 (79%)	31/31 (100%)	16/32 (50%)	15/15 (100%)
Sidechain	78/103 (76%)	50/62 (81%)	28/34 (82%)	0/7 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	16/18 (89%)	8/10 (80%)	7/7 (100%)	1/1 (100%)
Overall	156/199 (78%)	89/103 (86%)	51/73 (70%)	16/23 (70%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	27	TYR	CE2	34.12	124.68 – 111.18	-62.1
1	A	27	TYR	CD2	49.20	140.11 – 125.31	-56.4
1	A	21	TRP	CZ2	30.73	121.76 – 106.66	-55.3
1	A	21	TRP	CZ3	38.03	129.20 – 113.60	-53.4
1	A	21	TRP	CE3	36.83	129.06 – 111.96	-48.9
1	A	20	HIS	CE1	25.61	149.70 – 125.30	-45.9
1	A	21	TRP	CH2	40.90	133.06 – 114.56	-44.8
1	A	21	TRP	CD1	42.03	136.18 – 116.78	-43.5
1	A	22	THR	CG2	50.86	27.15 – 15.95	26.2
1	A	23	THR	CG2	50.74	27.15 – 15.95	26.1
1	A	30	THR	CG2	49.78	27.15 – 15.95	25.2
1	A	28	LYS	CG	52.03	30.67 – 19.17	23.6
1	A	20	HIS	CD2	44.37	137.40 – 103.40	-22.4
1	A	10	ILE	CG2	45.27	24.63 – 10.43	19.5
1	A	11	VAL	CG1	48.23	28.40 – 14.60	19.4
1	A	15	ILE	CD1	46.01	21.91 – 5.01	19.3
1	A	5	SER	CB	36.00	71.24 – 56.34	-18.7
1	A	12	SER	CB	36.52	71.24 – 56.34	-18.3
1	A	33	PRO	CA	35.28	71.13 – 55.53	-18.0
1	A	2	PRO	CA	35.35	71.13 – 55.53	-17.9
1	A	26	PRO	CA	35.88	71.13 – 55.53	-17.6
1	A	11	VAL	CG2	48.44	29.20 – 13.40	17.2
1	A	23	THR	CB	40.88	78.10 – 61.30	-17.2
1	A	15	ILE	CG2	41.53	24.63 – 10.43	16.9
1	A	30	THR	CB	41.92	78.10 – 61.30	-16.5
1	A	22	THR	CB	42.00	78.10 – 61.30	-16.5
1	A	3	LEU	CD2	51.46	32.60 – 15.60	16.1
1	A	10	ILE	CD1	40.62	21.91 – 5.01	16.1
1	A	31	LEU	CD2	51.20	32.60 – 15.60	15.9
1	A	19	ASP	CA	24.84	64.88 – 44.48	-14.6
1	A	34	MET	CE	44.93	26.97 – 7.37	14.2
1	A	12	SER	CA	29.11	69.25 – 48.25	-14.1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	29	ASP	CA	26.99	64.88 – 44.48	-13.6
1	A	5	SER	CA	30.42	69.25 – 48.25	-13.5
1	A	31	LEU	CA	27.09	66.36 – 44.96	-13.3
1	A	24	ARG	CA	26.03	68.35 – 45.25	-13.3
1	A	3	LEU	CA	27.29	66.36 – 44.96	-13.3
1	A	9	LYS	CA	28.38	67.97 – 45.97	-13.0
1	A	20	HIS	CA	25.99	68.24 – 44.74	-13.0
1	A	6	LYS	CA	28.45	67.97 – 45.97	-13.0
1	A	35	GLN	CA	29.34	67.31 – 45.91	-12.7
1	A	34	MET	CA	27.57	67.38 – 44.88	-12.7
1	A	17	LYS	CA	30.68	67.97 – 45.97	-11.9
1	A	28	LYS	CA	30.70	67.97 – 45.97	-11.9
1	A	27	TYR	CA	28.90	70.88 – 45.38	-11.5
1	A	14	ARG	CA	30.67	68.35 – 45.25	-11.3
1	A	21	TRP	CA	28.74	70.54 – 44.84	-11.3
1	A	10	ILE	CA	33.21	75.08 – 48.18	-10.6
1	A	30	THR	CA	34.44	75.37 – 49.07	-10.6
1	A	23	THR	CA	35.62	75.37 – 49.07	-10.1
1	A	11	VAL	CA	33.84	76.93 – 48.03	-9.9
1	A	15	ILE	CA	36.41	75.08 – 48.18	-9.4
1	A	22	THR	CA	37.99	75.37 – 49.07	-9.2
1	A	16	CYS	CA	30.67	75.07 – 41.27	-8.1
1	A	25	CYS	CA	30.89	75.07 – 41.27	-8.1
1	A	13	CYS	CA	31.86	75.07 – 41.27	-7.8

### 7.1.5 Random Coil Index (RCI) plots

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

