



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:39 AM GMT

PDB ID : 3MJE
Title : Structure of A-type Ketoreductases from Modular Polyketide Synthase
Authors : Zheng, J.; Taylor, C.A.; Piasecki, S.K.; Keatinge-Clay, A.T.
Deposited on : 2010-04-12
Resolution : 1.36 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2199 (1.40-1.32)
Clashscore	102246	2337 (1.40-1.32)
Ramachandran outliers	100387	2280 (1.40-1.32)
Sidechain outliers	100360	2279 (1.40-1.32)
RSRZ outliers	91569	2199 (1.40-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	<div> <div>9%</div> <div>73%</div> <div>19%</div> <div>• • 5%</div> </div>
1	B	496	<div> <div>8%</div> <div>71%</div> <div>19%</div> <div>• • 6%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AmphB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	470	Total	C	N	O	S	0	0	0
			3422	2137	629	647	9			
1	B	468	Total	C	N	O	S	0	0	0
			3406	2128	625	644	9			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q93NW7
A	-19	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-18	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-17	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-16	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-15	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-14	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-13	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-12	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-11	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-10	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-9	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-8	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-7	LEU	-	EXPRESSION TAG	UNP Q93NW7
A	-6	VAL	-	EXPRESSION TAG	UNP Q93NW7
A	-5	PRO	-	EXPRESSION TAG	UNP Q93NW7
A	-4	ARG	-	EXPRESSION TAG	UNP Q93NW7
A	-3	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-2	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-1	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	0	MET	-	EXPRESSION TAG	UNP Q93NW7
B	-20	MET	-	EXPRESSION TAG	UNP Q93NW7
B	-19	GLY	-	EXPRESSION TAG	UNP Q93NW7
B	-18	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-17	SER	-	EXPRESSION TAG	UNP Q93NW7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-15	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-14	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-13	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-12	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-11	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-10	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-9	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-8	GLY	-	EXPRESSION TAG	UNP Q93NW7
B	-7	LEU	-	EXPRESSION TAG	UNP Q93NW7
B	-6	VAL	-	EXPRESSION TAG	UNP Q93NW7
B	-5	PRO	-	EXPRESSION TAG	UNP Q93NW7
B	-4	ARG	-	EXPRESSION TAG	UNP Q93NW7
B	-3	GLY	-	EXPRESSION TAG	UNP Q93NW7
B	-2	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-1	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	0	MET	-	EXPRESSION TAG	UNP Q93NW7

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total	O	0	0
			227	227		
4	B	247	Total	O	0	0
			247	247		

R457	P466	V469	S470	L471	L472	SER	ASP	GLU
------	------	------	------	------	------	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.65Å 63.76Å 71.74Å 72.99° 67.21° 89.79°	Depositor
Resolution (Å)	62.76 – 1.36 31.38 – 1.36	Depositor EDS
% Data completeness (in resolution range)	93.3 (62.76-1.36) 83.7 (31.38-1.36)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 1.36Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.202 , 0.225 0.200 , 0.221	Depositor DCC
R_{free} test set	9658 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	12.8	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 51.7	EDS
Estimated twinning fraction	0.059 for h,-k,h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 191224 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7404	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.58	42/3490 (1.2%)	1.55	49/4764 (1.0%)
1	B	1.55	36/3473 (1.0%)	1.57	70/4741 (1.5%)
All	All	1.56	78/6963 (1.1%)	1.56	119/9505 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

The worst 5 of 78 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	GLU	CB-CG	-15.02	1.23	1.52
1	A	166	GLU	CG-CD	-13.15	1.32	1.51
1	B	469	VAL	CB-CG2	-10.96	1.29	1.52
1	B	284	GLU	CB-CG	-10.91	1.31	1.52
1	A	389	SER	CA-CB	-10.57	1.37	1.52

The worst 5 of 119 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	382	ARG	NE-CZ-NH2	-21.13	109.74	120.30
1	B	382	ARG	NE-CZ-NH1	20.25	130.43	120.30
1	A	328	ARG	NE-CZ-NH2	-19.54	110.53	120.30
1	B	382	ARG	CD-NE-CZ	17.09	147.53	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ARG	NE-CZ-NH2	15.42	128.01	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	228	GLY	Mainchain,Peptide
1	A	280	ALA	Mainchain
1	B	72	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3422	0	3387	51	0
1	B	3406	0	3375	60	0
2	A	48	0	26	6	0
2	B	48	0	26	8	0
3	B	6	0	8	0	0
4	A	227	0	0	5	0
4	B	247	0	0	12	0
All	All	7404	0	6822	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:476:NDP:C5N	4:A:587:HOH:O	1.84	1.21
2:A:476:NDP:H5N	4:A:587:HOH:O	1.38	1.19
2:B:476:NDP:C5N	4:B:533:HOH:O	1.93	1.15
1:B:262:LEU:O	1:B:266:LEU:HD13	1.53	1.08
2:B:476:NDP:H5N	4:B:533:HOH:O	1.55	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/496 (94%)	454 (97%)	11 (2%)	1 (0%)	52	22
1	B	464/496 (94%)	452 (97%)	12 (3%)	0	100	100
All	All	930/992 (94%)	906 (97%)	23 (2%)	1 (0%)	56	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/351 (94%)	314 (95%)	15 (5%)	33	4
1	B	327/351 (93%)	311 (95%)	16 (5%)	31	4
All	All	656/702 (93%)	625 (95%)	31 (5%)	32	4

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	410	ARG
1	B	23	ARG
1	B	401	MET
1	B	-2	SER
1	B	48	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	HIS
1	A	414	GLN
1	B	245	HIS
1	A	310	HIS
1	B	310	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	476	-	42,52,52	1.66	10 (23%)	55,80,80	2.42	17 (30%)
2	NDP	B	476	-	42,52,52	1.51	8 (19%)	55,80,80	2.27	14 (25%)
3	GOL	B	477	-	5,5,5	0.93	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	476	-	-	0/30/77/77	0/5/5/5
2	NDP	B	476	-	-	0/30/77/77	0/5/5/5
3	GOL	B	477	-	-	0/4/4/4	0/0/0/0

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	476	NDP	C4N-C5N	-3.74	1.41	1.49
2	B	476	NDP	C4N-C5N	-3.14	1.42	1.49
2	A	476	NDP	PN-O2N	-2.31	1.45	1.54
2	B	476	NDP	C3B-C4B	-2.25	1.46	1.53
2	B	476	NDP	PN-O2N	-2.02	1.46	1.54

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	476	NDP	N3A-C2A-N1A	-11.46	120.12	128.89
2	B	476	NDP	N3A-C2A-N1A	-9.63	121.52	128.89
2	B	476	NDP	C4B-O4B-C1B	-5.83	103.32	109.72
2	A	476	NDP	C1B-N9A-C4A	-4.66	119.91	126.94
2	A	476	NDP	O4B-C1B-C2B	-4.54	98.38	106.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	476	NDP	6	0
2	B	476	NDP	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	470/496 (94%)	0.71	44 (9%) 11 11	7, 14, 28, 42	0
1	B	468/496 (94%)	0.66	38 (8%) 15 14	6, 13, 28, 42	0
All	All	938/992 (94%)	0.69	82 (8%) 13 12	6, 14, 28, 42	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	399	VAL	8.4
1	A	413	ARG	7.4
1	B	410	ARG	7.4
1	A	410	ARG	6.8
1	B	401	MET	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NDP	A	476	48/48	0.91	0.22	1.64	18,23,36,38	0
2	NDP	B	476	48/48	0.93	0.17	1.10	14,19,23,27	0
3	GOL	B	477	6/6	0.92	0.10	0.37	15,17,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.