



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 10:39 AM GMT

PDB ID : 3MQ3  
Title : Crystal structure of native bovine PDP1c  
Authors : Guo, Y.; Ernst, S.R.; Carroll, D.W.; Hackert M.L.  
Deposited on : 2010-04-27  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 3068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate dehydrogenase phosphatase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	378	2967	1885	516	556	10	0	0	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	99	Total O 99 99	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate dehydrogenase phosphatase 1



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.27Å    75.27Å    172.64Å 90.00°    90.00°    120.00°	Depositor
Resolution (Å)	27.00 – 2.80 26.01 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.0 (27.00-2.80) 97.2 (26.01-2.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	6.43 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
$R$ , $R_{free}$	0.224 , 0.284 0.226 , 0.278	Depositor DCC
$R_{free}$ test set	716 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.9	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 66.7	EDS
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.34$	Xtriage
Outliers	0 of 14130 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	3068	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.91	0/3031	0.89	3/4119 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	253	ARG	NE-CZ-NH2	6.06	123.33	120.30
1	A	253	ARG	NE-CZ-NH1	-5.62	117.49	120.30
1	A	344	LEU	CA-CB-CG	5.24	127.36	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	312	LYS	Peptide
1	A	317	ASN	Peptide
1	A	435	LEU	Peptide
1	A	50	ALA	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2967	0	2858	144	0
2	A	2	0	0	0	0
3	A	99	0	0	11	0
All	All	3068	0	2858	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

All (144) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:435:LEU:CD1	1:A:436:PRO:HD3	1.39	1.49
1:A:435:LEU:HD22	1:A:436:PRO:N	1.36	1.35
1:A:435:LEU:HD13	1:A:436:PRO:CD	1.57	1.32
1:A:435:LEU:HD22	1:A:436:PRO:CA	1.66	1.25
1:A:50:ALA:CB	1:A:51:PRO:HD2	1.64	1.24
1:A:50:ALA:HB3	1:A:51:PRO:CD	1.79	1.12
1:A:50:ALA:CB	1:A:51:PRO:CD	2.30	1.09
1:A:435:LEU:HD22	1:A:436:PRO:CD	1.82	1.09
1:A:224:MET:HE2	1:A:238:THR:HA	1.28	1.09
1:A:438:GLU:HA	1:A:438:GLU:OE2	1.58	1.01
1:A:435:LEU:HD21	1:A:436:PRO:HG3	1.44	0.99
1:A:294:GLN:O	1:A:298:ILE:HG12	1.63	0.98
1:A:11:PRO:HA	3:A:548:HOH:O	1.66	0.95
1:A:54:ASP:OD2	1:A:448:THR:HG22	1.67	0.94
1:A:435:LEU:CD2	1:A:436:PRO:N	2.30	0.94
1:A:435:LEU:CD2	1:A:436:PRO:CG	2.48	0.92
1:A:435:LEU:CD2	1:A:436:PRO:CD	2.47	0.92
1:A:124:HIS:HD2	1:A:126:ASN:H	1.11	0.92
1:A:435:LEU:CD2	1:A:436:PRO:CA	2.49	0.89
1:A:435:LEU:CD2	1:A:436:PRO:HG3	2.02	0.89
1:A:435:LEU:HD22	1:A:436:PRO:CG	2.04	0.87
1:A:435:LEU:CG	1:A:436:PRO:HD3	2.05	0.87
1:A:50:ALA:HB3	1:A:51:PRO:HD2	0.89	0.85
1:A:435:LEU:CD1	1:A:436:PRO:CD	2.30	0.85

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:MET:CE	1:A:238:THR:HA	2.07	0.84
1:A:11:PRO:HB2	1:A:14:VAL:H	1.42	0.83
1:A:124:HIS:CD2	1:A:126:ASN:H	1.96	0.83
1:A:431:LYS:HD3	1:A:432:MET:HB2	1.62	0.81
1:A:435:LEU:H	1:A:436:PRO:CD	1.97	0.78
1:A:325:THR:HG21	3:A:498:HOH:O	1.84	0.77
1:A:435:LEU:HD22	1:A:436:PRO:HA	1.65	0.76
1:A:250:GLU:CD	1:A:253:ARG:NH1	2.43	0.72
1:A:435:LEU:H	1:A:435:LEU:HD13	1.54	0.72
1:A:184:ASP:H	1:A:190:ASN:HD22	1.37	0.72
1:A:75:HIS:HE1	1:A:284:ASP:OD1	1.72	0.71
1:A:75:HIS:CE1	1:A:284:ASP:OD1	2.43	0.71
1:A:276:LEU:HD22	1:A:278:PRO:HD2	1.72	0.71
1:A:365:TYR:HE1	1:A:405:GLN:HG3	1.55	0.70
1:A:50:ALA:HB1	1:A:51:PRO:CD	2.21	0.70
1:A:435:LEU:H	1:A:436:PRO:HD3	1.55	0.70
1:A:435:LEU:CD2	1:A:436:PRO:HA	2.19	0.70
1:A:214:HIS:HD2	3:A:469:HOH:O	1.75	0.68
1:A:11:PRO:HB2	1:A:13:GLN:H	1.58	0.67
1:A:94:VAL:HA	1:A:97:LEU:HD22	1.77	0.66
1:A:432:MET:SD	3:A:563:HOH:O	2.54	0.65
1:A:406:ASN:HD22	1:A:409:THR:H	1.45	0.65
1:A:54:ASP:HB3	1:A:448:THR:HG21	1.78	0.65
1:A:435:LEU:HD22	1:A:435:LEU:C	2.13	0.64
1:A:435:LEU:HD22	1:A:436:PRO:CB	2.28	0.64
1:A:250:GLU:CD	1:A:253:ARG:HH12	2.00	0.64
1:A:75:HIS:HD2	3:A:470:HOH:O	1.82	0.63
1:A:435:LEU:CG	1:A:436:PRO:CD	2.72	0.62
1:A:352:THR:HG22	1:A:353:MET:HG3	1.81	0.62
1:A:73:ASP:H	1:A:202:THR:HG22	1.65	0.62
1:A:238:THR:HG22	1:A:238:THR:O	2.00	0.61
1:A:124:HIS:HD2	1:A:126:ASN:N	1.92	0.61
1:A:438:GLU:CA	1:A:438:GLU:OE2	2.42	0.61
1:A:277:MET:HE2	3:A:518:HOH:O	2.00	0.61
1:A:435:LEU:HD13	1:A:435:LEU:N	2.16	0.60
1:A:273:LEU:HD13	1:A:316:PRO:HB2	1.84	0.60
1:A:435:LEU:HD13	1:A:436:PRO:HD3	0.65	0.60
1:A:229:GLU:HG3	3:A:506:HOH:O	2.00	0.60
1:A:317:ASN:N	3:A:550:HOH:O	2.34	0.60
1:A:353:MET:HE1	1:A:361:ILE:HD12	1.83	0.60
1:A:45:GLN:HE21	1:A:448:THR:HB	1.67	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:GLU:HG3	1:A:188:PHE:HE1	1.67	0.59
1:A:406:ASN:ND2	1:A:409:THR:H	2.01	0.59
1:A:333:HIS:HE1	1:A:339:ASP:OD2	1.85	0.59
1:A:251:VAL:HG12	1:A:255:LYS:HE2	1.85	0.59
1:A:99:HIS:CE1	1:A:103:LEU:CD2	2.85	0.59
1:A:132:GLU:HG3	1:A:188:PHE:CE1	2.37	0.58
1:A:202:THR:HG21	1:A:221:SER:OG	2.03	0.58
1:A:298:ILE:O	1:A:299:GLU:OE2	2.21	0.58
1:A:11:PRO:HB2	1:A:14:VAL:N	2.16	0.57
1:A:297:VAL:O	1:A:299:GLU:N	2.38	0.56
1:A:80:CYS:O	1:A:84:VAL:HG13	2.06	0.55
1:A:276:LEU:CD2	1:A:278:PRO:HD2	2.36	0.54
1:A:75:HIS:HE1	1:A:284:ASP:CG	2.10	0.54
1:A:99:HIS:CE1	1:A:103:LEU:HD21	2.42	0.54
1:A:43:SER:HA	1:A:449:ILE:O	2.09	0.53
1:A:250:GLU:OE2	1:A:253:ARG:NH1	2.42	0.53
1:A:435:LEU:N	1:A:436:PRO:CD	2.65	0.53
1:A:432:MET:C	3:A:563:HOH:O	2.47	0.53
1:A:353:MET:CE	1:A:361:ILE:HD12	2.39	0.53
1:A:435:LEU:CD1	1:A:436:PRO:CG	2.87	0.53
1:A:124:HIS:CD2	1:A:125:PRO:HD2	2.45	0.52
1:A:298:ILE:O	1:A:299:GLU:CD	2.48	0.52
1:A:106:GLU:HA	1:A:106:GLU:OE1	2.08	0.52
1:A:244:ASN:C	1:A:244:ASN:HD22	2.13	0.52
1:A:116:LEU:H	1:A:116:LEU:CD2	2.21	0.52
1:A:365:TYR:HE1	1:A:405:GLN:CG	2.22	0.52
1:A:442:MET:HA	1:A:442:MET:CE	2.40	0.51
1:A:133:ALA:HA	1:A:189:LEU:HD21	1.93	0.51
1:A:23:TYR:CG	1:A:24:SER:N	2.79	0.51
1:A:47:PRO:HA	1:A:52:ILE:HD12	1.93	0.51
1:A:184:ASP:N	1:A:190:ASN:HD22	2.07	0.51
1:A:433:LEU:O	1:A:435:LEU:HD13	2.11	0.51
1:A:285:VAL:HG13	1:A:289:TRP:CD2	2.46	0.50
1:A:47:PRO:CA	1:A:52:ILE:HD12	2.41	0.50
1:A:442:MET:HE2	1:A:442:MET:HA	1.94	0.49
1:A:435:LEU:N	1:A:435:LEU:CD1	2.77	0.48
1:A:116:LEU:N	1:A:116:LEU:CD2	2.77	0.48
1:A:433:LEU:O	1:A:435:LEU:CD1	2.62	0.47
1:A:435:LEU:N	1:A:436:PRO:HD3	2.19	0.47
1:A:406:ASN:ND2	1:A:408:ALA:H	2.12	0.47
1:A:169:PHE:CG	1:A:328:PRO:HG3	2.49	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:ARG:HB3	1:A:350:TRP:NE1	2.30	0.45
1:A:435:LEU:CD2	1:A:436:PRO:CB	2.92	0.45
1:A:336:ARG:HB3	1:A:338:GLN:OE1	2.17	0.45
1:A:49:ASN:O	1:A:50:ALA:O	2.33	0.45
1:A:317:ASN:C	1:A:317:ASN:ND2	2.70	0.45
1:A:116:LEU:HD22	1:A:116:LEU:N	2.31	0.45
1:A:22:GLU:O	1:A:22:GLU:HG3	2.17	0.45
1:A:22:GLU:CG	1:A:22:GLU:O	2.64	0.45
1:A:367:THR:HG23	1:A:367:THR:O	2.18	0.44
1:A:352:THR:HG21	1:A:415:ALA:HA	1.98	0.44
1:A:416:VAL:HG12	1:A:416:VAL:O	2.11	0.44
1:A:435:LEU:CD2	1:A:435:LEU:C	2.82	0.44
1:A:75:HIS:HE1	1:A:284:ASP:OD2	2.00	0.44
1:A:107:ASN:O	1:A:108:ALA:C	2.55	0.44
1:A:435:LEU:HD21	1:A:436:PRO:CG	2.19	0.44
1:A:325:THR:HB	1:A:327:GLU:H	1.82	0.43
1:A:54:ASP:HB3	1:A:448:THR:CG2	2.46	0.43
1:A:255:LYS:HE3	3:A:505:HOH:O	2.18	0.43
1:A:411:LEU:HD23	1:A:411:LEU:HA	1.92	0.43
1:A:23:TYR:O	1:A:24:SER:CB	2.67	0.43
1:A:219:GLY:H	1:A:243:HIS:HD2	1.67	0.43
1:A:99:HIS:HE1	1:A:103:LEU:HD21	1.82	0.42
1:A:441:ARG:HA	1:A:444:ARG:O	2.19	0.42
1:A:271:ARG:HD2	3:A:512:HOH:O	2.19	0.42
1:A:73:ASP:N	1:A:202:THR:HG22	2.34	0.41
1:A:12:PRO:O	1:A:13:GLN:CB	2.68	0.41
1:A:163:GLU:O	1:A:167:ASN:ND2	2.53	0.41
1:A:49:ASN:HD21	1:A:441:ARG:HH21	1.69	0.41
1:A:365:TYR:CE1	1:A:405:GLN:HG3	2.44	0.41
1:A:91:TYR:CE2	1:A:171:ARG:HG3	2.55	0.41
1:A:251:VAL:O	1:A:255:LYS:HG3	2.21	0.41
1:A:412:ILE:HG13	1:A:449:ILE:HD12	2.03	0.41
1:A:50:ALA:HB1	1:A:51:PRO:HD3	2.01	0.41
1:A:132:GLU:CD	1:A:132:GLU:H	2.23	0.41
1:A:191:TYR:HA	1:A:191:TYR:HD1	1.73	0.41
1:A:254:LEU:HB3	1:A:266:VAL:HG11	2.02	0.40
1:A:60:THR:OG1	1:A:122:HIS:HE1	2.04	0.40
1:A:44:ASN:HB3	1:A:412:ILE:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	366/467 (78%)	342 (93%)	16 (4%)	8 (2%)	8 28

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	50	ALA
1	A	298	ILE
1	A	24	SER
1	A	107	ASN
1	A	435	LEU
1	A	316	PRO
1	A	437	GLU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	314/404 (78%)	280 (89%)	34 (11%)	8 23

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	SER
1	A	27	VAL
1	A	46	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	69	LEU
1	A	84	VAL
1	A	97	LEU
1	A	100	GLU
1	A	103	LEU
1	A	115	LEU
1	A	116	LEU
1	A	145	TRP
1	A	171	ARG
1	A	194	LEU
1	A	202	THR
1	A	213	LEU
1	A	224	MET
1	A	235	SER
1	A	237	VAL
1	A	238	THR
1	A	244	ASN
1	A	254	LEU
1	A	268	LYS
1	A	272	LEU
1	A	285	VAL
1	A	286	LYS
1	A	314	ILE
1	A	338	GLN
1	A	367	THR
1	A	406	ASN
1	A	432	MET
1	A	435	LEU
1	A	438	GLU
1	A	441	ARG
1	A	448	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	49	ASN
1	A	75	HIS
1	A	99	HIS
1	A	122	HIS
1	A	124	HIS
1	A	167	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	190	ASN
1	A	214	HIS
1	A	228	GLN
1	A	243	HIS
1	A	244	ASN
1	A	317	ASN
1	A	333	HIS
1	A	356	GLN
1	A	405	GLN
1	A	406	ASN
1	A	410	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	378/467 (80%)	-0.19	9 (2%) 62 50	20, 38, 72, 91	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	462	TYR	4.5
1	A	356	GLN	3.4
1	A	100	GLU	3.3
1	A	438	GLU	2.8
1	A	99	HIS	2.8
1	A	107	ASN	2.7
1	A	368	GLY	2.5
1	A	312	LYS	2.5
1	A	256	LEU	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	A	501	1/1	0.98	0.15	0.17	34,34,34,34	0
2	MN	A	502	1/1	0.97	0.13	-0.39	32,32,32,32	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.