



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:47 PM GMT

PDB ID : 4MR8
Title : Crystal structure of the extracellular domain of human GABA(B) receptor bound to the antagonist CGP35348
Authors : Geng, Y.; Bush, M.; Mosyak, L.; Wang, F.; Fan, Q.R.
Deposited on : 2013-09-17
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	 3% 83% 13% • •
2	B	433	 3% 83% 9% • 7%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid type B receptor subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	3271	2090	551	616	14	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	460	ASP	-	EXPRESSION TAG	UNP Q9UBS5
A	461	TYR	-	EXPRESSION TAG	UNP Q9UBS5
A	462	LYS	-	EXPRESSION TAG	UNP Q9UBS5
A	463	ASP	-	EXPRESSION TAG	UNP Q9UBS5
A	464	ASP	-	EXPRESSION TAG	UNP Q9UBS5
A	465	ASP	-	EXPRESSION TAG	UNP Q9UBS5
A	466	ASP	-	EXPRESSION TAG	UNP Q9UBS5
A	467	LYS	-	EXPRESSION TAG	UNP Q9UBS5

- Molecule 2 is a protein called Gamma-aminobutyric acid type B receptor subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	403	3214	2052	543	604	15	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	467	ASP	-	EXPRESSION TAG	UNP O75899
B	468	TYR	-	EXPRESSION TAG	UNP O75899
B	469	LYS	-	EXPRESSION TAG	UNP O75899
B	470	ASP	-	EXPRESSION TAG	UNP O75899
B	471	ASP	-	EXPRESSION TAG	UNP O75899
B	472	ASP	-	EXPRESSION TAG	UNP O75899
B	473	ASP	-	EXPRESSION TAG	UNP O75899
B	474	LYS	-	EXPRESSION TAG	UNP O75899

- Molecule 3 is a polymer of unknown type called SUGAR (5-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	5	Total	C	N	O	0	0
			60	34	2	24		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is (S)-(3-AMINOPROPYL)(DIETHOXYMETHYL)PHOSPHINIC ACID (three-letter code: 2BW) (formula: $C_8H_{20}NO_4P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			14	8	1	4	1		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	3	Total	C	N	O	0	0
			38	22	2	14		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	241	Total	O	0	0
			241	241		
7	B	233	Total	O	0	0
			233	233		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-aminobutyric acid type B receptor subunit 1



- Molecule 2: Gamma-aminobutyric acid type B receptor subunit 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.98Å 112.75Å 73.45Å 90.00° 97.80° 90.00°	Depositor
Resolution (Å)	59.67 – 2.15 59.67 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.1 (59.67-2.15) 99.1 (59.67-2.15)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.14Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.209 , 0.223 0.216 , 0.230	Depositor DCC
R_{free} test set	3081 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.1	EDS
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 61657 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7085	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2BW, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3354	0.63	0/4547
2	B	0.44	0/3287	0.64	1/4449 (0.0%)
All	All	0.43	0/6641	0.64	1/8996 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	389	ASN	N-CA-C	-5.04	97.39	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3271	0	3193	47	1
2	B	3214	0	3160	25	1
3	A	60	0	52	0	0
4	A	14	0	13	1	0
5	A	14	0	20	0	0
6	B	38	0	34	1	0
7	A	241	0	0	6	0
7	B	233	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7085	0	6472	72	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:ASN:HB2	7:B:831:HOH:O	1.60	1.01
1:A:462:LYS:HD3	1:A:462:LYS:H	1.24	1.00
1:A:462:LYS:HD3	1:A:462:LYS:N	1.80	0.93
6:B:501:NAG:H83	7:B:790:HOH:O	1.68	0.92
2:B:389:ASN:OD1	2:B:391:THR:HB	1.74	0.87
1:A:269:LYS:HD3	1:A:452:ILE:CG2	2.09	0.81
1:A:368:SER:O	1:A:377:ARG:HG2	1.85	0.76
1:A:294:THR:OG1	1:A:297:GLU:HG3	1.86	0.76
1:A:234:VAL:HG21	1:A:260:GLU:HB3	1.70	0.73
1:A:377:ARG:HG3	1:A:378:LEU:N	2.05	0.69
1:A:269:LYS:HD3	1:A:452:ILE:HG22	1.73	0.69
1:A:269:LYS:HD3	1:A:452:ILE:HG21	1.76	0.67
1:A:368:SER:C	1:A:377:ARG:HG2	2.14	0.66
2:B:244:LYS:HD2	2:B:270:GLU:HB3	1.77	0.65
7:A:829:HOH:O	2:B:112:LYS:HE3	1.98	0.64
1:A:153:SER:HB3	1:A:170:HIS:CE1	2.33	0.63
1:A:324:MET:SD	1:A:332:LYS:HD2	2.38	0.63
1:A:462:LYS:CD	1:A:462:LYS:N	2.58	0.62
1:A:53:VAL:HG13	1:A:123:ILE:HG12	1.82	0.61
1:A:368:SER:O	1:A:377:ARG:CG	2.48	0.61
1:A:391:GLN:HE21	1:A:394:ARG:HH22	1.49	0.60
1:A:235:LYS:HD3	7:A:747:HOH:O	2.00	0.60
2:B:388:PHE:CD1	2:B:395:LEU:HD23	2.38	0.59
1:A:88:ILE:HG12	4:A:506:NAG:H82	1.85	0.58
1:A:241:ASP:OD2	1:A:461:TYR:CE1	2.60	0.55
2:B:65:THR:HG23	2:B:68:VAL:HG23	1.88	0.55
1:A:302:VAL:O	1:A:305:HIS:HD2	1.90	0.55
1:A:53:VAL:CG1	1:A:123:ILE:HG12	2.37	0.54
1:A:462:LYS:CD	1:A:462:LYS:H	1.99	0.54
1:A:283:TRP:HA	1:A:286:ILE:HD12	1.90	0.53
2:B:389:ASN:OD1	2:B:391:THR:CB	2.53	0.52
1:A:377:ARG:HG3	1:A:378:LEU:H	1.73	0.52
1:A:85:ARG:HB2	1:A:88:ILE:HD12	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:HIS:HD2	7:A:677:HOH:O	1.95	0.50
1:A:142:MET:HB3	2:B:115:LYS:HD2	1.94	0.50
1:A:142:MET:HE1	7:A:782:HOH:O	2.12	0.49
1:A:198:THR:HG22	1:A:225:SER:OG	2.12	0.49
1:A:224:GLN:HB3	1:A:233:PRO:HB3	1.95	0.49
2:B:188:LYS:HE3	2:B:447:ASP:O	2.13	0.49
2:B:109:ASP:HB3	2:B:112:LYS:HB2	1.95	0.48
1:A:258:PHE:HA	1:A:261:VAL:HB	1.95	0.48
1:A:211:GLU:O	1:A:215:GLU:HG3	2.13	0.48
1:A:197:GLN:HB2	1:A:249:PHE:HB3	1.96	0.48
2:B:129:MET:HB2	2:B:152:VAL:O	2.14	0.48
2:B:320:GLU:OE1	2:B:427:LYS:HB2	2.14	0.47
1:A:226:PHE:HB3	1:A:233:PRO:HG2	1.96	0.47
2:B:207:ARG:HH11	2:B:207:ARG:HG3	1.79	0.47
1:A:259:CYS:HA	1:A:298:MET:CE	2.44	0.47
2:B:389:ASN:CG	2:B:391:THR:HB	2.35	0.47
2:B:206:GLN:HE21	2:B:210:GLU:HG2	1.80	0.46
1:A:302:VAL:O	1:A:305:HIS:CD2	2.68	0.46
1:A:368:SER:HB2	7:A:737:HOH:O	2.15	0.46
2:B:197:ARG:NH2	2:B:247:ASP:O	2.49	0.46
1:A:321:ILE:HG12	1:A:402:GLU:O	2.16	0.46
2:B:199:GLY:HA3	2:B:248:VAL:HG11	1.97	0.45
1:A:178:PRO:HA	1:A:181:VAL:HG22	1.99	0.45
1:A:351:PRO:HB3	1:A:404:VAL:HG23	1.98	0.45
2:B:303:LEU:HB2	2:B:306:ASN:ND2	2.33	0.44
1:A:327:GLN:HB3	7:A:805:HOH:O	2.18	0.44
1:A:409:VAL:HG23	1:A:417:MET:HB2	1.98	0.44
2:B:376:LEU:CD1	2:B:385:ILE:HG13	2.49	0.43
2:B:286:GLU:O	2:B:289:TRP:HB3	2.18	0.43
1:A:66:PRO:O	1:A:69:GLN:HG3	2.18	0.42
2:B:388:PHE:HD1	2:B:395:LEU:HD23	1.83	0.42
2:B:160:THR:HA	2:B:161:PRO:HD3	1.96	0.41
1:A:234:VAL:HA	1:A:237:LEU:HD12	2.02	0.41
2:B:89:ARG:NH2	2:B:99:LEU:O	2.52	0.41
2:B:261:ALA:O	2:B:307:LEU:HD11	2.21	0.41
2:B:54:PRO:HG2	2:B:97:TYR:CD2	2.55	0.41
1:A:235:LYS:HE3	1:A:239:ARG:HD2	2.02	0.40
1:A:259:CYS:CA	1:A:298:MET:HE2	2.50	0.40
1:A:191:LYS:HB2	1:A:459:ASP:HB2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:ASP:OD1	2:B:301:ARG:NH1[2_444]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/420 (96%)	389 (96%)	13 (3%)	1 (0%)	52	51
2	B	397/433 (92%)	386 (97%)	11 (3%)	0	100	100
All	All	800/853 (94%)	775 (97%)	24 (3%)	1 (0%)	56	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	457	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/362 (98%)	349 (99%)	5 (1%)	74	80
2	B	349/375 (93%)	342 (98%)	7 (2%)	63	67
All	All	703/737 (95%)	691 (98%)	12 (2%)	68	74

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	249	PHE
1	A	278	TRP
1	A	378	LEU
1	A	391	GLN
1	A	462	LYS
2	B	204	ASP
2	B	301	ARG
2	B	349	VAL
2	B	370	GLN
2	B	385	ILE
2	B	391	THR
2	B	435	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	144	ASN
1	A	161	GLN
1	A	196	GLN
1	A	292	ASN
1	A	305	HIS
1	A	327	GLN
1	A	340	HIS
1	A	391	GLN
2	B	150	ASN
2	B	192	HIS
2	B	206	GLN
2	B	306	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	501	1,3	14,14,15	0.81	0	15,19,21	2.25	4 (26%)
3	NAG	A	502	3	14,14,15	0.77	0	15,19,21	1.50	4 (26%)
3	BMA	A	503	3	11,11,12	0.81	0	14,15,17	2.15	3 (21%)
3	BMA	A	504	3	11,11,12	1.05	1 (9%)	14,15,17	1.59	2 (14%)
3	FUC	A	505	3	10,10,11	1.21	0	14,14,16	1.73	3 (21%)
6	NAG	B	501	2,6	14,14,15	0.91	1 (7%)	15,19,21	1.14	1 (6%)
6	NAG	B	502	6	14,14,15	0.69	0	15,19,21	1.19	1 (6%)
6	FUC	B	503	6	10,10,11	1.20	1 (10%)	14,14,16	1.10	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	501	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	502	3	-	0/6/23/26	0/1/1/1
3	BMA	A	503	3	-	0/2/19/22	0/1/1/1
3	BMA	A	504	3	-	0/2/19/22	0/1/1/1
3	FUC	A	505	3	-	0/0/17/20	0/1/1/1
6	NAG	B	501	2,6	-	0/6/23/26	0/1/1/1
6	NAG	B	502	6	-	0/6/23/26	0/1/1/1
6	FUC	B	503	6	-	0/0/17/20	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	503	FUC	C2-C3	2.21	1.55	1.52
3	A	504	BMA	C2-C3	2.30	1.55	1.52
6	B	501	NAG	C1-C2	2.53	1.56	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	BMA	O6-C6-C5	-5.01	94.78	111.33
3	A	503	BMA	O3-C3-C2	-4.21	102.39	110.00
6	B	502	NAG	C2-N2-C7	-2.94	119.26	123.04
3	A	501	NAG	O4-C4-C3	-2.56	104.56	110.34
3	A	502	NAG	C4-C3-C2	-2.55	107.26	111.23
3	A	502	NAG	C2-N2-C7	-2.54	119.78	123.04
6	B	501	NAG	O4-C4-C3	-2.51	104.69	110.34
3	A	502	NAG	C6-C5-C4	-2.49	106.88	113.02
3	A	501	NAG	C4-C3-C2	-2.30	107.66	111.23
3	A	503	BMA	O2-C2-C3	-2.19	105.72	110.12
3	A	502	NAG	C8-C7-N2	-2.09	112.10	116.11
6	B	503	FUC	C1-C2-C3	2.21	112.15	109.54
3	A	501	NAG	C2-N2-C7	2.29	125.98	123.04
3	A	505	FUC	C1-C2-C3	2.71	112.74	109.54
3	A	504	BMA	C1-O5-C5	2.91	115.94	112.25
3	A	505	FUC	O5-C1-C2	3.08	115.86	110.86
3	A	504	BMA	O5-C1-C2	3.30	116.21	110.86
3	A	505	FUC	C1-O5-C5	3.87	118.35	112.38
3	A	501	NAG	C1-O5-C5	6.99	121.12	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	501	NAG	1	0

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	506	1	14,14,15	0.76	0	15,19,21	1.16	1 (6%)
5	2BW	A	507	-	9,13,13	1.96	1 (11%)	8,16,16	1.43	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	506	1	-	0/6/23/26	0/1/1/1
5	2BW	A	507	-	-	0/8/17/17	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	507	2BW	P-O4	5.73	1.60	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	507	2BW	P-C6-C7	-3.35	109.18	114.48
4	A	506	NAG	C2-N2-C7	-2.38	119.97	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	506	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/420 (96%)	0.57	14 (3%) 49 59	29, 50, 88, 107	0
2	B	403/433 (93%)	0.44	12 (2%) 54 64	28, 45, 76, 107	0
All	All	810/853 (94%)	0.50	26 (3%) 51 61	28, 48, 86, 107	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	461	TYR	6.1
2	B	388	PHE	4.6
2	B	378	ALA	4.4
1	A	222	PHE	3.7
2	B	387	ASP	3.4
2	B	53	SER	3.2
2	B	93	LEU	3.1
2	B	454	ASP	3.0
1	A	339	ARG	2.8
1	A	269	LYS	2.6
1	A	419	TRP	2.5
2	B	453	ASN	2.4
1	A	445	TRP	2.4
1	A	129	CYS	2.4
2	B	97	TYR	2.4
1	A	239	ARG	2.3
2	B	326	GLN	2.3
1	A	417	MET	2.3
2	B	452	ILE	2.3
2	B	303	LEU	2.3
1	A	404	VAL	2.2
2	B	451	ILE	2.2
1	A	443	LEU	2.1
1	A	212	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	270	LYS	2.1
1	A	458	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	502	14/15	0.93	0.17	-	83,87,89,89	0
3	BMA	A	504	11/12	0.75	0.21	-	98,101,103,105	0
3	BMA	A	503	11/12	0.71	0.15	-	91,93,96,96	0
3	FUC	A	505	10/11	0.67	0.35	-	89,90,91,91	0
6	NAG	B	502	14/15	0.88	0.14	-	81,85,88,88	0
6	FUC	B	503	10/11	0.94	0.13	-	77,77,79,79	0
6	NAG	B	501	14/15	0.86	0.13	-	71,75,77,78	0
3	NAG	A	501	14/15	0.86	0.15	-	77,81,83,86	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	A	506	14/15	0.87	0.15	-0.03	71,75,77,78	0
5	2BW	A	507	14/14	0.98	0.13	-0.81	31,33,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.