wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 03:40 AM BST

PDB ID	:	2MT9
Title	:	Solution structure of holo_FldB
Authors	:	Jin, C.; Fu, W.; Ye, Q.
Deposited on	:	2014-08-15

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	173	

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:1-A:121, A:125-A:173 (170)	0.21	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 4, 7, 8, 10, 14, 15, 16, 18
2	2, 3, 5, 6, 12, 17, 19
3	9, 11
Single-model clusters	13; 20

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2754 atoms, of which 1335 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Flavodoxin-2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	173	2704	893	1316	212	275	8	0

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$).



Mol	Chain	Residues	Atoms					
			Total	C	H	N	O	P
2	A	1	50	17	19	4	9	1

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Flavodoxin-2

Chain A:



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 4. Colouring as in section 4.1 above.

- Molecule 1: Flavodoxin-2

Chain A:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
AMBER	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mt9_cs.str
Number of chemical shift lists	1
Total number of shifts	2094
Number of shifts mapped to atoms	2094
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.66±0.00	0±0/1397 (0.0±0.0%)	1.01±0.02	2±1/1897 (0.1±0.1%)
All	All	0.66	0/27940 (0.0%)	1.01	43/37940 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.5±0.7
All	All	0	10

There are no bond-length outliers.

5 of 11 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	96	TRP	CB-CG-CD2	8.02	137.03	126.60	3	2
1	A	96	TRP	CB-CG-CD1	-6.82	118.14	127.00	6	2
1	A	172	TYR	CB-CG-CD2	-6.12	117.33	121.00	13	1
1	A	43	MET	CA-CB-CG	5.88	123.30	113.30	10	6
1	A	119	TRP	CB-CG-CD2	5.86	134.22	126.60	11	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	93	TYR	Sidechain	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	156	ARG	Sidechain	3
1	A	172	TYR	Sidechain	1
1	A	22	ARG	Sidechain	1
1	A	171	HIS	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1363	1298	1298	6±2
2	A	31	19	19	0±0
All	All	27880	26340	26340	129

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

5 of 33 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:93:TYR:CE2	1:A:96:TRP:HB2	0.65	2.27	20	1
1:A:84:LEU:HD12	1:A:105:HIS:CD2	0.63	2.28	16	1
1:A:95:GLU:HB2	1:A:96:TRP:CE3	0.57	2.35	3	2
1:A:81:ILE:HG21	1:A:168:MET:HG2	0.54	1.80	9	18
1:A:24:ILE:HD13	1:A:158:GLN:HA	0.53	1.80	10	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	168/173 (97%)	155±2 (92±1%)	11±2 (7±1%)	2±1 (1±0%)	20 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	3360/3460 (97%)	3093 (92%)	223 (7%)	44 (1%)	20 66

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	126	PHE	19
1	A	12	CYS	12
1	A	57	ASP	10
1	A	98	LEU	2
1	A	97	PHE	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	146/148 (99%)	125±2 (86±2%)	21±2 (14±2%)	8 47
All	All	2920/2960 (99%)	2497 (86%)	423 (14%)	8 47

5 of 47 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	42	LEU	20
1	A	5	LEU	20
1	A	133	ILE	20
1	A	90	GLN	20
1	A	53	ILE	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	FMN	A	201	-	31,33,33	1.91±0.03	2±0 (6±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	FMN	A	201	-	32,50,50	2.34±0.05	1±0 (3±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	201	-	-	0±0,18,18,18	0±0,3,3,3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	201	FMN	C4A-N5	6.17	1.42	1.33	16	20
2	A	201	FMN	C5A-N5	6.08	1.44	1.35	9	20

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
2	A	201	FMN	C2-N1-C10	12.44	124.00	113.39	2	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 89% for the entire structure.

7.1 Chemical shift list 1

File name: 2mt9_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2094
Number of shifts mapped to atoms	2094
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	29

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	173	-0.31 ± 0.11	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	156	0.08 ± 0.11	None needed (< 0.5 ppm)
$^{13}C'$	169	-0.05 ± 0.14	None needed (< 0.5 ppm)
^{15}N	166	0.29 ± 0.24	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1837 atoms were assigned a chemical shift out of a possible 2057. 26 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	831/838 (99%)	332/334 (99%)	336/340 (99%)	163/164 (99%)
Sidechain	874/1008 (87%)	524/585 (90%)	337/396 (85%)	13/27 (48%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	132/211 (63%)	70/110 (64%)	56/92 (61%)	6/9 (67%)
Overall	1837/2057 (89%)	926/1029 (90%)	729/828 (88%)	182/200 (91%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	22	ARG	NE	116.65	92.63 – 76.73	20.1
1	A	56	TRP	HE3	11.36	9.33 – 5.33	10.1
1	A	142	LEU	HB3	-1.67	3.34 – -0.26	-8.9
1	A	95	GLU	HB2	0.45	3.08 – 0.98	-7.5
1	A	95	GLU	HB3	0.38	3.10 – 0.90	-7.4
1	A	98	LEU	HD11	-1.06	2.16 – -0.64	-6.5
1	A	98	LEU	HD12	-1.06	2.16 – -0.64	-6.5
1	A	98	LEU	HD13	-1.06	2.16 – -0.64	-6.5
1	A	90	GLN	HB3	0.30	3.37 – 0.67	-6.4
1	A	98	LEU	HG	-0.57	3.16 – -0.14	-6.3
1	A	35	LEU	HD23	-0.98	2.14 – -0.66	-6.1
1	A	35	LEU	HD21	-0.98	2.14 – -0.66	-6.1
1	A	35	LEU	HD22	-0.98	2.14 – -0.66	-6.1
1	A	142	LEU	HD13	-0.88	2.16 – -0.64	-5.8
1	A	142	LEU	HD12	-0.88	2.16 – -0.64	-5.8
1	A	142	LEU	HD11	-0.88	2.16 – -0.64	-5.8
1	A	10	SER	H	11.64	11.23 – 5.33	5.7
1	A	148	ASN	HD22	9.91	9.59 – 4.69	5.7
1	A	121	THR	HG21	-0.13	2.29 – -0.01	-5.5
1	A	121	THR	HG22	-0.13	2.29 – -0.01	-5.5
1	A	121	THR	HG23	-0.13	2.29 – -0.01	-5.5
1	A	142	LEU	HG	-0.26	3.16 – -0.14	-5.4
1	A	148	ASN	ND2	125.05	124.24 – 101.34	5.4
1	A	14	THR	H	11.40	11.34 – 5.14	5.1
1	A	143	ALA	HB2	0.09	2.61 – 0.11	-5.1
1	A	143	ALA	HB3	0.09	2.61 – 0.11	-5.1
1	A	143	ALA	HB1	0.09	2.61 – 0.11	-5.1
1	A	107	LYS	HE3	1.94	3.86 – 1.96	-5.1
1	A	143	ALA	H	5.19	11.19 – 5.19	-5.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: