wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 03:35 AM BST

PDB ID	:	2MU3
Title	:	Spider wrapping silk fibre architecture arising from its modular soluble protein precursor
Authors	:	Xu, L.; Tremblay, M.; Meng, Q.; Liu, X.; Rainey, J.K.; Lefevre, T.; Sarker, M.; Orrell, K.E.; Leclerc, J.; Pezolet, M.; Auger, M.
Deposited on	:	2014-09-03

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	199	54% 9% • 37%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:12-A:77, A:86-A:144 (125)	0.53	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 6, 11, 12, 13, 15, 16, 17, 18, 20
2	2, 5, 7, 8, 9, 14
Single-model clusters	10; 19

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2632 atoms, of which 1296 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Aciniform spidroin 1.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	199	2632	809	1296	240	287	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Aciniform spidroin 1

Chain A: 54% 9% • 37%

A1 G2 P3 Q4 G5 G6 F7 G8 A9 T10 G11 S14 L17 I18 V21 A22 N23 N27 T33 I43 S46 R50 Q53 S57 N64 P78 A79 G80 S81 D82 T83 S84 A85 L100 N104 S114 V120 S145 F146 L147 S148 T149 S150

S151 S152 S153 A154 S155 Y156 S157 Q158 A159 S160 A161 S162 S163 T164 S165 G166 A167 G168 Y169 T170 G171 P172 S173 G174 P175 S176 T177 G178 P179 S180 G181 Y182 P183 G184 P185 L186 G187 G188 G189 A190 P191 F192 G193 Q194 S195 G196 F197 G198 G199

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 18. Colouring as in section 4.1 above.

- Molecule 1: Aciniform spidroin 1

Chain A: 46% 15% • 37%

A1 G2 P3 Q4 G5 G6 F7 G8 A9 T10 G11 S14 L17 I18 S19 R20 V21 A22 N23 A24 S29 T30 L31 R32 T33 S40 I43 S46 R50 Q53 S54 L55 A56 S57 N64 N65 L77 P78 A79 G80 S81 D82 T83 S84 A85 Y86 A87 Q88 A89 F90

L94 L100 N104 S114 V120 V131 D132 S144 S145 F146 L147 S148 T149 S150 S151 S152 S153 A154 S155 Y156 S157 Q158 A159 S160 A161 S162 S163 T164 S165 G166 A167 G168 Y169 T170 G171 P172 S173 G174 P175 S176 T177 G178 P179 S180 G181 Y182 P183 G184 P185 L186 G187 G188 G189

A190 P191 F192 G193 Q194 S195 G196 F197 G198 G199

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
NMRpipe	refinement	1.0
NMRpipe	refinement	1.0
NMRpipe	refinement	1.0
NMRpipe	refinement	1.0
NMRpipe	refinement	1.0
NMRpipe	refinement	1.0
ARIA	refinement	2
ARIA	structure solution	2
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mu3_cs.str
Number of chemical shift lists	1
Total number of shifts	1991
Number of shifts mapped to atoms	1991
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	873	888	888	14±4
All	All	17460	17760	17760	277

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 126 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:52:ALA:HB2	1:A:129:ILE:HG23	0.66	1.67	6	3
1:A:61:VAL:HA	1:A:65:ASN:OD1	0.63	1.93	1	7
1:A:43:ILE:HG13	1:A:44:ALA:N	0.62	2.09	5	2
1:A:59:LEU:HD11	1:A:65:ASN:ND2	0.62	2.09	5	1
1:A:50:ARG:NH1	1:A:123:ALA:HA	0.61	2.10	12	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	125/199 (63%)	96±3 (77±2%)	23±3 (18±3%)	6±2 (5±1%)	5 28
All	All	2500/3980 (63%)	1926 (77%)	454 (18%)	120 (5%)	5 28

5 of 23 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	33	THR	17
1	A	114	SER	14
1	A	46	SER	13
1	A	57	SER	13
1	A	21	VAL	13

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	94/139 (68%)	90±1 (96±1%)	4±1 (4±1%)	42 84
All	All	1880/2780 (68%)	1803 (96%)	77 (4%)	42 84

5 of 31 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	53	GLN	9
1	A	100	LEU	8
1	A	23	ASN	6
1	A	37	THR	6
1	A	104	ASN	5

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: 2mu3_cs.str

Chemical shift list name: *assigned_chem_shift_list*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1991
Number of shifts mapped to atoms	1991
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	198	-0.39 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	164	0.12 ± 0.06	None needed (< 0.5 ppm)
$^{13}C'$	196	-0.23 ± 0.10	None needed (< 0.5 ppm)
^{15}N	187	0.23 ± 0.15	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1210 atoms were assigned a chemical shift out of a possible 1348. 0 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	621/625 (99%)	248/250 (99%)	249/250 (100%)	124/125 (99%)
Sidechain	589/688 (86%)	350/394 (89%)	218/256 (85%)	21/38 (55%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	0/35 (0%)	0/19 (0%)	0/16 (0%)	0/0 (—%)
Overall	1210/1348 (90%)	598/663 (90%)	467/522 (89%)	145/163 (89%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: